



Endocrinología y Nutrición



195 - DHA-RICH FISH OIL SUPPLEMENTATION IMPROVES BODY COMPOSITION WITHOUT INFLUENCE OF PPAR-GAMMA PRO12ALA POLYMORPHISM IN PATIENTS WITH TYPE 2 DIABETES

A. Mansoori^a and F. Koohdani^b

^aAhvaz Junishapur University of Medical Sciences. Irán. ^bTehran University of Medical Sciences. Irán.

Resumen

Introduction: The aims of this research were to investigate (1) the impact of DHA-rich fish oil supplementation on body composition, plasma adiponectin level, and PPAR γ gene expression; (2) whether the effect of DHA-rich fish oil supplementation on the aforementioned variables is modulated by PPAR γ Pro12Ala polymorphism.

Methods: We genotyped PPAR γ Pro12Ala polymorphism in subjects with T2DM. Ala carriers and non-Ala carriers were randomly assigned to DHA-rich fish oil or placebo intake for 8 weeks.

Results: Glycemic control was not affected by the intervention. The supplementation with DHA-rich fish oil decreased WC ($p \leq 0.001$), body fat mass ($p = 0.01$), body fat percent ($p = 0.04$), and visceral fat rating ($p = 0.02$) as well as trunk fat mass ($p = 0.04$). Weight, BMI, fat free mass, Adiponectin level and PPAR γ gene expression changes showed no significant difference. No gene-diet interaction was found on body composition, adiponectin level and PPAR γ gene expression.

Conclusions: DHA-rich fish oil supplementation favorably modulated body composition in patients with type 2 diabetes, and could be useful to reduce visceral obesity. However, the PPAR γ Pro12Ala polymorphism did not influence the changes in the desired variables.