



# Endocrinología y Nutrición



## 329 - The relationship between daily dietary nutrient intakes and total kidney cyst volume on patients with autosomal dominant polycystic kidney disease (ADPKD)

Y. Sevim<sup>a</sup>, O. Persil Ozkan<sup>b</sup>, E. Cebeci<sup>c</sup>, H. Merve Bayram<sup>a</sup>, H. Nur Onur<sup>a</sup>, Y. Savas<sup>c</sup>, S. Ozturk<sup>c</sup> and G. Kiziltan<sup>d</sup>

<sup>a</sup>Department of Nutrition and Dietetics. Istanbul Gelisim University. Turquía. <sup>b</sup>Department of Nutrition and Dietetics. Istanbul Arel University. Turquía. <sup>c</sup>Haseki Training and Research Hospital. Clinic of Nephrology. Istanbul. Turquía. <sup>d</sup>Baskent University. Department of Nutrition and Dietetics. Ankara. Turquía.

### Resumen

ADPKD is one of the most common inherited and a multi-systemic disease which is responsible for about 10% of cases of end-stage renal disease (ESRD). Cardiovascular problems are a major cause of morbidity and mortality in patients with ADPKD. This study was conducted to determine the relationship between daily dietary nutrient intakes and kidney volume. The study was carried out on 22 male and 38 female patients, with a mean age of  $48.6 \pm 11.3$  years, who were diagnosed with ADPKD. In this study the nutrient intakes of patients were estimated by the 24h dietary records obtained on 3 consecutive days. The mean eGFR was calculated to be  $54.8 \pm 36.46$  ml/dk/1.73 m<sup>2</sup> and the mean kidney volume was  $1012 \pm 776.8$ ml. Patients total dietary energy and protein intake was found to be  $25.57 \pm 9.32$  kcal/kg/day,  $0.87 \pm 0.32$  g/kg/day (respectively). Patients' percentage of energy from carbohydrates, protein, total fat and saturated fat was determined as  $49 \pm 7\%$ ,  $14 \pm 2\%$ ,  $36 \pm 6\%$  and  $11.6 \pm 3.6\%$  (respectively). Patients', especially women, mean dietary folat intake was much lower than the DRI (Daily Recommended Intake). Patients' mean dietary calcium and potassium intakes were insufficient. Patients' mean dietary sodium intake was higher than the DRI recommendation. It was found that when patients were divided in to two groups, by their median kidney volume (743 ml), there was no difference in their carbohydrates, protein and total fat intakes. Dietary calcium, magnesium, potassium, sodium and zinc intakes were lower; phosphorus and iron intakes were similar in patients with higher kidney volume than the patients with smaller kidney volume but these relationships weren't statistically important. As a conclusion, patients' total fat, saturated fat and sodium intakes were higher; dietary folat, calcium and potassium intakes were inadequate. Due to the lack of medical diet therapy recommendations on patients with ADPKD, patients should be followed by a specialist (renal) dietitian to protect malnutrition or hypertension and to prevent further renal injury in order to slow polycystic kidney progression to ESRD.