

MARKERS OF MATURATION AND OPG/RANKL SYSTEM IN HUMAN PRIMARY OSTEOBLASTS CULTURES

M. Giner, M.J. Montoya, M.A. Vázquez, R.M. Moruno, M.J. Miranda y R. Pérez-Cano.

Bone Metabolism Unit, Internal Medicine, University Hospital Virgen Macarena. Seville, Spain.

C-TERMINAL PARATHYROID HORMONE-RELATED PROTEIN (107-139) INHIBITS APOPTOSIS THROUGH VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR-2 IN HUMAN OSTEOBLASTIC CELLS

V. Alonso¹, A.R. de Gortázar¹, R. Berruguete², P. Esbrit¹ and M.V. Álvarez-Arroyo².

¹Bone and Mineral Metabolism Laboratory and ²Fundación Jiménez Díaz-UTE (Capio Group). MADRID, Spain

CHARACTERIZATION OF AN EXPERIMENTAL MODEL OF OSTEOPOROSIS IN RABBITS INDUCED BY COMBINED OVARECTOMY AND CORTICOSTEROIDS

S. Castañeda¹, E. Calvo², R. Largo², C. de la Piedra², M. Díaz-Curiel², M. Bellido² y G. Herrero-Beaumont².

¹Rheumatology Unit, ²Bone and Mineral Metabolism Laboratory. Hospital de la Princesa, Fundación Jiménez Díaz, Madrid, Spain.

A HIGH CTx LEVEL IS ASSOCIATED WITH A HIGHER RATE OF BONE LOSS IN EARLY POSTMENOPAUSAL WOMEN

L. Navarro², J.A. Blázquez¹, M. Cháfer², F. Mateos¹, E. Almar³, J. del Pino⁴.

¹Departments of Internal Medicine1 and Clinical Chemistry. ²University Hospital, Albacete, Spain. ³Epidemiology, Primary Care, Albacete, Spain. ⁴Department of Medicine. University Hospital, Salamanca, Spain.

Nuestro agradecimiento a

Lilly, Roche/GSK, Sanofi-Aventis/Amirall y Nycomed por hacer viable esta iniciativa.