



REVISTA MÉDICA DEL
HOSPITAL GENERAL
DE MÉXICO

www.elsevier.es/hgmx



CLINICAL CASE

Septic arthritis of the acromioclavicular joint due to *Streptococcus agalactiae*. Case report

C.A.J. Mangas-Loría^a, M.A. Fuentes-Nucamendi^b, A.F. Sánchez-Chávez^{a,*},
C. Monreal-Chairez^a, C. Ramos-Córdova^a



^a Orthopaedics, Hospital General de México "Dr. Eduardo Liceaga", Mexico City, Mexico

^b Orthopaedics, Spinal Column Unit, Orthopaedics Service, Hospital General de México "Dr. Eduardo Liceaga", Mexico City, Mexico

Received 19 September 2016; accepted 26 October 2016

Available online 23 November 2016

KEYWORDS

Septic arthritis;
Acromioclavicular
joint;
*Streptococcus
agalactiae*

Abstract Septic arthritis of the acromioclavicular joint is a rare entity: only 30 cases have been reported in the literature since 1985. We present the case of a 53-year-old diabetic male, with septic arthritis of one acromioclavicular joint due to *Streptococcus agalactiae*. Current condition characterised by neck pain, limited movement of the right shoulder; hyperthermia, hyperaemia and increased volume in the acromioclavicular joint. Upon physical examination, increased volume was found from the proximal third of the deltoid to the middle third of the clavicle, pain on palpation, localised hyperthermia, limited range of motion. X-ray with enlargement of soft tissues, presence of subcutaneous gas and increased space in the acromioclavicular joint compared with the contralateral. Ceftriaxone and Clindamycin were administered at therapeutic doses and an acromioclavicular arthrotomy was performed, obtaining 10 ml of purulent material from the sub-deltoid and 2 ml from the joint. Five days later, *Streptococcus agalactiae* was reported. Clinical improvement was observed and it was decided to discharge the patient. © 2016 Sociedad Médica del Hospital General de México. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Artritis séptica;
Articulación
acromioclavicular;
*Streptococcus
agalactiae*

Artritis séptica acromioclavicular por *Streptococcus agalactiae*. Reporte de caso

Resumen La artritis séptica acromioclavicular (ASAC) es una entidad poco frecuente, desde 1985 se han reportado 30 casos en la literatura. Se presenta el caso de un paciente masculino, diabético, de 53 años, con ASAC monoarticular por *Streptococcus agalactiae*. Padecimiento actual caracterizado por dolor en cuello, limitación al movimiento de hombro derecho,

* Corresponding author at: Lago Zirahuen 49, Colonia Anáhuac II, Delegación Miguel Hidalgo, CP 11320 Mexico City, Mexico.
E-mail address: Andresdoc19@hotmail.com (A.F. Sánchez-Chávez).

hipertermia, hiperemia y aumento de volumen en articulación acromioclavicular. A la exploración física, con aumento de volumen en tercio proximal de deltoides hasta tercio medio de clavícula, dolor a la palpación, hipertermia localizada, arcos de movilidad limitados. Rx con aumento de partes blandas, presencia de gas subcutáneo y aumento de espacio articular acromioclavicular en comparación con la contralateral. Se administra Ceftriaxona y Clindamicina a dosis terapéuticas y se realiza artrotomía acromioclavicular derecha, se obtiene 10cc de material purulento subdeltóideo y 2cc articular. Se reporta al 5º día *Streptococcus agalactiae*. Presenta mejoría clínica y se decide egreso.

© 2016 Sociedad Médica del Hospital General de México. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Septic arthritis of the acromioclavicular joint is a rare entity; 30 cases have been reported in the literature since 1985.¹

The most frequent etiological agent at any age is *Staphylococcus aureus* in 75% of cases, followed by gram-negative in 20% of cases. Chronic presentation is usually due to microbacteria and filamentous fungi.^{1,2}

Once the germ is found in the synovial membrane, it begins to reproduce and various virulence factors are produced such as extracellular toxins, enzymes, adhesins, proteins with bacterial cell walls, which trigger a flood of inflammation through T cells, B cells and macrophages. As a consequence, pro-inflammatory molecules including TNF-α, IL-6 and IL-8, immunomodulatory and inflammatory cytokines are produced by monocytes, macrophages and synovial fibroblasts so that in a period of 24–48 h, intra-articular effusion occurs with a polymorphonuclear leucocyte count of up to 50,000 per cubic millimetre, glucose reduction and an increase in proteins in the synovial fluid.^{3,4}

Commonly affected joints are the knee (50%), hip (20%), shoulder (8%), ankle (7%), wrist (7%); Martinez-Morillo et al., reported a series of 101 cases of which 6% affected the acromioclavicular joint.^{2,5,6}

The incidence of septic arthritis of the acromioclavicular joint is 2–10/100,000 in the general population; if the patient has an articular prosthesis, the incidence increases to 30–60/100,000, ranging in age from 17 to 79 with an average age of 54 years and is more prevalent in males 5:1.^{6–9}

Treatment is medical-surgical and is considered urgent due to the potential for serious consequences. Its natural evolution leads to the destruction of the articular cartilage and the adjacent bone.

Up until now, only 30 cases of septic arthritis of the acromioclavicular joint have been reported.¹⁰ The purpose of this document is to present the case of a patient with septic arthritis of one acromioclavicular joint caused by *Streptococcus agalactiae*.

Case report

Male, 53 years of age, type 2 diabetic, condition evolving for 15 days, characterised by neck pain, limited shoulder

movement, hyperthermia, hyperaemia and increased volume in the right acromioclavicular joint. Symptom management, presenting partial improvement. Evolution with an increase in symptomatology and fever.

The physical examination shows shoulder asymmetry with an increase in volume from the deltoid region to the middle third of the right clavicle (Fig. 1a), hyperthermia and pain on palpation 10/10 VAS in the acromioclavicular joint, limited range of motion in the joint, secondary to pain and without distal neurovascular compromise.

Lab tests: leukocytes: 16,800 mm³, neutrophils: 75.50%, glucose: 341 mg/dl.

Comparative AP radiography of shoulder (19/09/2014) (Fig. 2a and b): Right: with enlargement of soft tissue, presence of subcutaneous gas and increase in the acromioclavicular joint space (9 mm) compared to the contralateral. Left: Acromioclavicular space of 1.41 mm. No bone lesion data.

Intravenous antibiotic therapy initiated with 1 g of Ceftriaxone every 12 h and 600 mg of clindamycin every 8 h. Arthrotomy on the right acromioclavicular joint, surgical lavage and debridement were performed (Fig. 1b), obtaining 10 ml of subdeltoid purulent and 2 mm of articular exudate.

Culture report: *Streptococcus agalactiae*.

The same antibiotic regimen was continued for proper evolution until completion at the end of the 10 days. Evolution with improvement and decrease in leucocyte numbers (16,800 mm³ → 9300 mm³) and ESR (31 mm³/h → 10 mm³/h). Discharged with oral antibiotic (500 mg ciprofloxacin orally every 8 h and 300 mg clindamycin every 8 h) for 4 weeks.

Follow-up 2 weeks after surgery, asymptomatic. The last control was at 9 months where the X-ray showed a lytic acromioclavicular space of 11 mm and full range of motion in the right shoulder (Fig. 3).

Discussion

This is a rare pathology given the limited space and characteristics of the joint, and may even be underdiagnosed because it is not easy to distinguish a septic glenohumeral disorder from a simple physical examination.^{2,3,8,11,12}

In addition, it must be differentiated from traumatic synovitis, abscesses, acute rheumatic fever, joint infection




Figure 1 (a) Increase in volume, hyperthermia and hyperaemia from the deltoid area to the middle third of the clavicle. (b) 10 cm³ of subdeltoid and 2 cc³ of articular purulent material were obtained.




Figure 2 Comparative shoulder X-rays: (a) Right shoulder X-ray: volume increase in soft tissues, with radiolucent images compatible with subcutaneous emphysema; irregular acromioclavicular bone edges, right acromioclavicular joint 9.61 mm. (b) Left shoulder X-ray: Acromioclavicular space 1.41 mm without any bone injury data.

due to microbacteria, cellulitis, acute osteomyelitis and haemophilia.^{3,9,13}

Septic arthritis of the acromioclavicular joint tends to appear in immunocompromised patients, either infected with HIV, rheumatoid arthritis, renal insufficiency, multiple myeloma, chronic steroid use, hepatic cirrhosis^{2,8}; as well as a history of venipuncture, local trauma, articular puncture, prior shoulder surgery, septic arthritis in another joint, periprosthetic infections and ulcers on diabetic feet.^{8,9,14}

The patient being studied had a history of chronic evolution as an insulin-dependent diabetic, who did not adhere well to treatment and did not use glucose-lowering drugs for 1 week prior to his admission. He also presented with plantar infections 6 and 12 months prior.

It is described that in the absence of trauma and articular manipulation, septic arthritis must be considered a consequence of haematogenous dissemination,^{8,9} as is likely in the case of our patient.

Patients present initially with general symptoms such as general malaise, diffuse pain, hyperthermia; after a few

days, the focus is on increased volume, local oedema that limits shoulder and even neck movement, pain upon superficial palpation of the anterior shoulder, O'Brien manoeuvre or active compression of the positive acromioclavicular joint, with evolution ranging between less than 24 h and up to 12 days.

The vast majority were previously treated with analgesics and anti-inflammatories, as in this case, leading to the delay in diagnosis and treatment 15 days after onset.

Chronic cases are attributed to microbacteria and can take up to a year to diagnose.¹³

Image studies are very useful for diagnosis.

A simple X-ray of the shoulder does not usually reveal bone changes in the initial stages,^{6,9,12} however, it is an easy, inexpensive, and quick study that does not merit patient preparation, is not user-dependent and provides a lot of data, both positive and negative, as in this case, where upon admittance, he already presented with changes that were very suggestive of the pathology and vital for regulating treatment.



Figure 3 Follow-up after 9 months: Right shoulder range of motion: abduction: 0°–180°; abduction: 0°–90°, without pain; flexion: 0°–170°; extension: 0°–40°.

Ultrasound is a tool for the early diagnosis and treatment of septic arthritis of the acromioclavicular joint,¹² it can detect an increase in echogenicity, distension of the joint cavity, bone irregularity, as well as guiding punctures, similarly, septic arthritis of the acromioclavicular joint can be ruled out if the joint space is less than 3 mm.^{15,16}

Magnetic resonance imaging (MRI) can detect septic joint changes in the first 24 h of onset,⁸ changes compatible with oedema, joint effusion, thickening of the cartilage and, as with ultrasound, septic arthritis of the acromioclavicular joint can be ruled out if the joint space is less than 3 mm in a coronal cut.¹²

A firm suspicion, knowledge of the pathology and a focused physical examination are vital for diagnosing septic arthritis of the acromioclavicular joint² and tests such as MRI and ultrasound may not always be available.³

Antibiotic treatment should initially be empirical and focus on the most common causal agent (*S. aureus*), and subsequently be adapted according to the cultivated agent; similarly, treatment provided should be directed at improving the general conditions with emphasis on the underlying pathologies and as a priority factor, drainage of the affected joint should be performed.

In the current case, the isolated agent is rare, Group B *Streptococcus agalactiae* (GBS), which is a commensal bacterium that is part of the intestinal and vaginal tract in 15–30% of healthy adults, but is one of the most significant pathogens in newborns. Endogenous factors such as diabetes alter the normal composition of the intestinal microbiota; these factors can partly explain why GBS is a significant cause of infection in soft tissues and the urinary tract, of sepsis and arthritis in patients with chronic disorders.⁴

In the published literature, we find only the case of a patient with bilateral septic arthritis of the acromioclavicular joints, septic arthritis in the left hip and right knee with a history of CRI, skin lesion and abscess, treated intravenously with Penicillin G Sodium, oral amoxicillin and surgical debridement.⁶

Kyu Cheol Noh et al., comment that septic arthritis of the acromioclavicular joint can be treated conservatively if fever and pain improve with the administration of antibiotics, and report that surgical treatment should be used as an option in the event of ineffective antibiotic therapy, persistent fever, pain, and changes in bone erosion as well as radiographic osteomyelitis.³

We consider that septic arthritis is an orthopaedic surgical emergency, and should be treated as such. The proteolytic enzymes of the leukocytes are released in the joint, which leads to the destruction of cartilage and irreversible joint damage in 48 h.³

The small size of the acromioclavicular joint is associated with a high risk of local spreading and more rapid destruction of the joint, as well as the potential danger of subsequent infection of the glenohumeral joint; given that it commonly presents in immunocompromised patients, an early diagnosis should be made and surgical treatment provided immediately.^{2,12}

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflict of interest

The authors declare that they have no conflict of interests.

References

- Blankstein A, Amsalem JL, Rubinstein E, et al. Septic arthritis of the acromioclavicular joint. *Arch Orthop Trauma Surg.* 1985;103:417–8.
- Iyengar K, Gudena R, Chitgopkar S, et al. Primary septic arthritis of the acromio-clavicular joint: case report and review of literature. *Arch Orthop Trauma Surg.* 2008;129:83–6.
- Noh K, Chung K, Yu H, et al. Arthroscopic treatment of septic arthritis of acromioclavicular joint. *Clin Orthop Surg.* 2010;2:186.
- Landwehr-Kenzel S, Henneke P. Interaction of *Streptococcus agalactiae* and cellular innate immunity in colonization and disease. *Frontiers Immunol.* 2014;5.
- Cone L, Gauto A, Kazi A, et al. Staphylococcal septic arthritis of the small joints of the shoulder girdle. *J Clin Rheumatol.* 2008;14:181–2.
- Martínez-Morillo M, Mateo Soria L, Riveros Frutos A, et al. Artritis séptica de la articulación acromioclavicular: una localización atípica. *Reumatol Clín.* 2014;10:37–42.
- Corey S, Agger W, Saterbak A. Acromioclavicular septic arthritis and sternoclavicular septic arthritis with contiguous pyomyositis. *Clin Orthoped Surg.* 2015;7:131.
- Hammel J, Kwon N. Septic arthritis of the acromioclavicular joint. *J Emerg Med.* 2005;29:425–7.

9. Hong M, Kim Y, Ham H. Acute septic arthritis of the acromioclavicular joint caused by *Haemophilus parainfluenzae*: a rare causative origin. *Clin Rheumatol*. 2015;34:811–4.
10. Hashemi-Sadraei N, Gupta R, Machicado JD, et al. Bilateral acromioclavicular septic arthritis as an initial presentation of *Streptococcus pneumoniae* endocarditis. *Case Rep Infect Dis*. 2014.
11. Chiang AS, Ropiak CR, Bosco JA II, et al. Septic arthritis of the acromioclavicular joint: a report of four cases. *Bull NYU Hosp Joint Dis*. 2007;65:308–11.
12. Bossert M, Prati C, Bertolini E, et al. Septic arthritis of the acromioclavicular joint. *Joint Bone Spine*. 2010;77:466–9.
13. Talbot C, Rhodes B. An atypical mycobacterial infection of the shoulder. *Int J Shoulder Surg*. 2012;6:64–6.
14. Zicat B, Rahme D, Swaraj K, et al. Septic arthritis of the acromioclavicular joint. *Clin Nucl Med*. 2006;31:145–6.
15. Blankstein A. Ultrasonography as a diagnostic modality in acromioclavicular joint pathologies. *Israel Med Assoc J*. 2005;7:28–30.
16. Widman DS, Craig JG, van Holsbeeck MT. Sonographic detection, evaluation and aspiration of infected acromioclavicular joints. *Skeletal Radiol*. 2001;30:388–92.