



REVISTA MÉDICA DEL
HOSPITAL GENERAL
DE MÉXICO

www.elsevier.es/hgmx



ORIGINAL ARTICLE

Changes in functional coupling of 5-HT_{1A} receptor to the G-protein in neocortex temporal tissues of patients with temporal lobe epilepsy



M. Cuellar-Herrera^{a,*}, L. Rocha^b, N. Saleh-Subaie^c, F. Velasco^a, G. Aguado-Carrillo^a, M. Alonso-Vanegas^d, R. Guevara-Guzmán^e, A.L. Velasco^a

^a *Epilepsy Clinic, Hospital General de México Dr. Eduardo Liceaga, Ciudad de México, Mexico*

^b *Department of Pharmacobiology, Center of Research and Advanced Studies, Ciudad de México, Mexico*

^c *Movement and Sleep Disorders Unit, Hospital General Dr. Manuel Gea González, Ciudad de México, Mexico*

^d *National Institute of Neurology and Neurosurgery "Manuel Velasco Suarez", Ciudad de México, Mexico*

^e *Departamento de Fisiología, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México, Mexico*

Received 2 March 2017; accepted 13 July 2017

Available online 3 May 2018

KEYWORDS

5-HT_{1A} receptor;
G-protein;
Lateral temporal
neocortex;
Mesial temporal lobe
epilepsy

Abstract

Background: The 5-hydroxytryptamine-1A receptors are known to be involved in the inhibition of seizures in epilepsy.

Objective: The main aim of the present study is to determine the functional coupling to G-protein of the 5-HT_{1A} receptor through 8-OH-DPAT-stimulated [³⁵S]GTPγS binding assay in lateral temporal neocortex tissue of patients with mesial temporal lobe epilepsy (mTLE) and correlate it with clinical data.

Material and methods: The activation of the G protein complex was determined by [³⁵S]GTPγS binding assay. The temporal neocortex tissue was obtained from 5 patients with mTLE during epilepsy surgery and from 5 subjects (autopsies) who died due to an accident and without history of neurological disease.

Results: We found that values of maximal stimulation (E_{max}) of [³⁵S]GTPγS binding significant decreased in the mTLE group (30.8%, $p < .05$) when compared to the autopsy samples (65.9%). The values of potency (pEC_{50}) were similar in both groups. However, we found no significant differences between the E_{max} and age of patient, age of seizure onset, duration of epilepsy and frequency of seizures. Also, the pEC_{50} values revealed no significant correlations with the clinical data.

* Corresponding author at: Dr. Balmis # 148, Col. Doctores, Delegación Cuauhtémoc, Ciudad de México, Mexico.
E-mail address: manolacuellar@yahoo.com.mx (M. Cuellar-Herrera).

PALABRAS CLAVE

Receptor 5-HT_{1A};
 Proteína-G;
 Neocorteza temporal lateral;
 Epilepsia del lóbulo temporal mesial

Conclusions: Our data provide evidence that the lateral temporal neocortex of patients with pharmacoresistant mTLE presents alterations in the functional coupling to G-protein of the 5-HT_{1A} receptor. Such a change could be involved in hyperexcitability in the neocortex of patients with mTLE.

© 2018 Sociedad Médica del Hospital General de México. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Cambios en el acoplamiento funcional del receptor 5-HT_{1A} a la proteína-G en tejido de neocorteza temporal de pacientes con epilepsia del lóbulo temporal

Resumen

Antecedentes: Los receptores 5-hidroxitriptamina-1A (5-HT_{1A}) están implicados en la inhibición de las crisis en epilepsia.

Objetivo: El presente estudio es determinar el acoplamiento funcional a la proteína-G del receptor 5-HT_{1A} a través del ensayo de unión [³⁵S]GTPγS estimulando al receptor por 8-OH-DPAT (agonista del receptor 5-HT_{1A}) en tejido de neocorteza temporal de pacientes con epilepsia del lóbulo temporal mesial (ELTm) y correlacionar con datos clínicos.

Material y métodos: La activación del complejo de la proteína G del receptor de 5-HT_{1A} se determinó por [³⁵S]GTPγS por ensayos de unión. Tejido de neocorteza temporal de 5 pacientes con ELTm obtenido de la cirugía de epilepsia y 5 tejidos de autopsias de sujetos que fallecieron por accidente o por causas diferentes a una enfermedad neurológica.

Resultados: Los resultados demostraron estimulación máxima (E_{max}) significativamente disminuida en el grupo ELTm (30.8%, p < 0,05), en comparación con tejido de muestras de autopsia (65.9%). Los valores de potencia (pEC₅₀) fueron similares en ambos grupos. Sin embargo, no se encontraron diferencias significativas entre la E_{max} con la edad del paciente, la edad de inicio de las crisis, duración con epilepsia y la frecuencia de crisis. Además, los valores de pEC₅₀ no revelaron correlación con los datos clínicos.

Conclusión: Los presente estudio proporcionan evidencia que la neocorteza temporal de pacientes con ELTm fármaco resistente presenta alteraciones en el acoplamiento funcional a la proteína-G del receptor 5-HT_{1A}. Tal cambio podría estar implicado en la hiperexcitabilidad que se observa en la neocorteza de pacientes con ELTm.

© 2018 Sociedad Médica del Hospital General de México. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The G-protein-coupled receptors (GPCRs) are membrane proteins that sense signaling molecules such as hormones, light, odors, taste and neurotransmitters.¹ GPCRs undergo conformational changes after ligand binding, causing the activation of signaling networks, resulting in a cellular response. G proteins are heterotrimers made of α , β and γ subunit. When the receptor stimulates the G protein, the α subunit releases guanosine diphosphate (GDP) and binds to GTP (guanosine triphosphate). The G α subunits are divided in G α_s , G $\alpha_{i/o}$, G $\alpha_{q/11}$, and G $\alpha_{12/13}$ based on sequence homology and functional similarities, each one active a cascade of second messengers¹ (Fig. 1). The G $\alpha_{i/o}$ subunit is interest, because it inhibits adenylyl cyclases (AC) and reduces the concentration of cyclic adenosine monophosphate (cAMP), this leads to mediate receptor-dependent inhibition of various types of adenylyl cyclases, in addition to being expressed more in brain.²

The metabotropic receptors G $\alpha_{i/o}$ -protein-coupled have been considered as promising drug targets in the treatment of several disease, related to excitation disorders such as epilepsy.

Epilepsy is a chronic neurological disorder characterized by increased excessive or synchronous abnormal neuronal activity in the brain.³ Numerous studies imply an imbalance between excitatory and inhibitory neurotransmission.⁴ The neurobiological basis of epilepsy involves alterations of neuronal functions at multiple levels. It is known that changes in neurotransmission systems as receptors coupled to G-proteins are involved in epileptic seizures.⁵⁻⁸ The activation of GPCR receptors specifically those coupled to G $\alpha_{i/o}$ proteins have demonstrated anticonvulsive, antiepileptic and neuroprotective effects in different experimental models of epilepsy.⁹⁻¹² In contrast, brain tissue of patients with epilepsy showed alterations in their density and functional activity.^{13-15,30} These evidences in preclinical models and brain tissue of patients with epilepsy suggest the participation of receptors to G $\alpha_{i/o}$ proteins in epilepsy.




Figure 1 Representative image of the G-protein-coupled receptors (GPCRs) associated with the different G α subunits (G α_s , G $\alpha_{i/o}$, G $\alpha_{q/11}$, and G $\alpha_{12/13}$). AMPc: cyclic adenosine monophosphate; DAG: diacylglycerol; IP₃: inositol triphosphate; +: stimulate; -: inhibit.

On the other hand, the 5-hydroxytryptamine-1A (5-HT_{1A}) receptors are coupled to G $\alpha_{i/o}$ -proteins. It is known that induce inhibitory effect on seizure activity in models animals of epilepsy.^{16–18}

Studies in patients with mesial temporal lobe epilepsy (mTLE) have demonstrated a decrease in 5-HT_{1A} receptor binding in the insula, anterior cingulate, and medial and lateral temporal regions, ipsilateral to the epileptic focus in positron emission tomography (PET).^{19–21} Studies *in vitro* autoradiography on resected of neocortex tissues of surrounding the epileptic focus of patients with mTLE showed a decrease in 5-HT_{1A} receptor binding in layers I-II,²² this could be associated to high excitability and facilitation of seizure activity.

The neocortical temporal tissues have important role in seizure generation and/or propagation.^{23,24} Previous studies describe that the neocortical temporal tissues of patients with mTLE presents a disorganization in synaptic circuits which consists of an increase and decrease of excitatory and inhibitory synapses, respectively.^{5,25}

These studies (PET or autoradiography) provide information on the expression of receptor and their distribution in the brain. However, it does not provide information on the functional coupling of 5-HT_{1A} receptor to the G $\alpha_{i/o}$ protein which could be implicated in the pathophysiology of the epilepsy, recalling that the activation of this G $\alpha_{i/o}$ protein induces inhibition.

The use of the [³⁵S]GTP γ S radioligand binding assay is a technique that allows the evaluation of the functional activity of the receptor occupation of G proteins. The [³⁵S]GTP γ S technique consists of the activation of receptors by selective agonists, promoting the exchange of GDP to GTP which facilitates the binding of sulfur-labeled GTP to the G $\alpha_{i/o}$ subunit, allowing the functional measurement of the receptors.²⁶

Therefore, the aim of the present study was to examine 5-HT_{1A} receptor-induced G-protein activation in lateral temporal neocortex tissue of patients with mTLE with analyses the [³⁵S]GTP γ S binding. Also, it is correlating of 5-HT_{1A}

receptor-induced G-protein functional activation with the subject's age, age of seizure onset, duration of epilepsy and frequency of seizures.

Methods

Patients group

The protocol was approved for scientific committees of the institutions involved in the present research (protocol number DI/15/403/03/32) and each patient signed an informed consent.

The inclusion criteria of protocol were established by the Epilepsy Clinic Program of the General Hospital of Mexico included patients with pharmacoresistant mTLE who had a detailed seizure calendar of at least 3 months prior to surgery in order to confirm the presence of seizures, blood levels of antiepileptic drugs to verify that they were within therapeutic range, magnetic resonance imaging to detect the presence of hippocampal sclerosis or atrophy, four serial electroencephalograms to determine the presence of temporal epileptiform activity and neuropsychological assessment to determine basal cognitive condition and identify behavioral or memory disorders related to seizures.

A standard anterior temporal lobectomy, ipsilateral to the epileptogenic zone, was performed in all patients after 48 h from the last seizure occurrence. Epileptic lateral neocortex temporal tissue was obtained from five patients with pharmacoresistant mTLE history (Table 1). During the surgical procedure, lateral temporal neocortex tissue was collected immediately after resection then quickly frozen in pulverized dry ice and stored at -70°C .

Autopsy samples

Lateral neocortex temporal tissue was obtained from the autopsy of five subjects, who died because of various causes,

Table 1 Clinical data of patients with pharmacoresistant temporal lobe epilepsy and autopsies.

Patients n = 5	Gender	Age (years)	Age of seizure onset (years)	Duration of epilepsy (years)	Frequency of seizures (per month)	AED	Diagnosis or cause of death (PMI in hours)	Side of focus
HUM-181	F	29	8	21	30	PHT/CBZ/VPA	mTLE	Right
HUM-172	M	35	13	22	4	VPA/PHT/LTG	mTLE	Left
HUM-176	F	38	3	35	15	TPM/VPA/CBZ/ PHT/LTG	mTLE	Right
HUM-120	M	45	17	28	48	CBZ/PHT	mTLE	Left
HUM-179	F	44	22	22	17	CBZ/VPA	mTLE	Right
C1	F	71	-	-	-	-	Diabetes (5)	-
C2	M	50	-	-	-	-	Hepatitis (6)	-
C3	F	40	-	-	-	-	Hypovolemic shock (5)	-
C4	F	39	-	-	-	-	Ovarian Cancer (3)	-
C5	M	29	-	-	-	-	Chronic renal failure (4)	-

M, male; F, female; AED, antiepileptic drugs; PHT, phenytoin; CBZ, carbamazepine; VPA, valproic acid; LTG, lamotrigine; TPM, topiramate; OXC, oxcarbazepine; PB, phenobarbital; CZP, clonazepam; RIS, Risperidone.

without evidence of neurological disease. The control autopsy samples were provided of the Institute of Forensic Sciences, in agreement with the Faculty of Medicine of the National Autonomous University of Mexico. Our tissue was dissected at the time of autopsy, with a postmortem interval of 3 to 6 h, immediately stored at -70°C and then manipulated as described below.

Binding assays

Membrane preparation

The lateral temporal neocortex tissues were prepared according to the method previously described²⁷ slight modifications. Briefly, samples (200–500 mg) were homogenized on ice in 50 mM Tris-HCl buffer (pH 7.4) using a Teflon-glass homogenizer. The homogenate was centrifuged at 15,000 rpm for 25 min at 4°C and the resulting pellet was re-suspended in fresh buffer and incubated for 30 min at 35°C . The centrifugation step was repeated, and the final pellet was re-suspended in 50 mM Tris-HCl buffer (pH 7.4) and stored at -70°C until use. Protein levels were determined by the method of Lowry et al. (1951).²⁸

[³⁵S]GTP γ S functional assay

The 5-HT_{1A} receptor-mediated G-protein activation experiments were carried as previously described for Spetea et al. (1998)²⁹ with slight modifications. The membrane fractions ($\approx 10\ \mu\text{g}$ of protein/sample) of lateral neocortex temporal tissue were incubated at 30°C for 60 min in Tris-EGTA buffer (1 M Tris-HCl, 0.2 M EGTA, 1 M MgCl₂, 1 M NaCl, pH 7.4), containing [³⁵S]GTP γ S (0.05 nM) and increasing concentrations (Log 10^{-11} to 10^{-5} M) of 8-OH-DPAT (5-HT_{1A} receptor agonist) in presence of excess GDP (10 mM) in a final volume of 1 ml. Total binding was measured in absence of the tested compound. Non-specific binding was determined in presence of 10 mM unlabeled GTP γ S and subtracted from total binding to calculate the specific binding. The reaction was initiated by adding [³⁵S]GTP γ S and completed with filtration of the samples through Whatman GF/B glass fiber filters; it was washed three times with ice-cold 50 mM Tris-HCl buffer (pH 7.4) using Brandel M-48 Cell Harvester. The dried filters were immersed in a Sigma-Fluor™ scintillation cocktail (Sigma) and then used to determine bound radioactivity as described above. The binding experiments were performed in triplicates.

Data analysis

The data were expressed as mean \pm S.E.M for the subject's age, age of seizure onset, duration of epilepsy and frequency of seizures. G-protein activation was given as percent of the specific [³⁵S]GTP γ S binding observed in absence of receptor ligands (basal activity). The binding data were subjected to non-linear regression analysis of concentration-effect curves performed by Prism (GraphPad Software, Inc.) for determined agonist potency (pEC₅₀) and the maximum stimulation (E_{max}) values. The data were expressed as mean \pm S.E.M. The results from the specific binding were examined statistically by Student's *t*-test to find differences of E_{max} and pEC₅₀ between autopsy samples and mTLE tissue. Pearson's correlation coefficients were calculated to establish a significant




Figure 2 Specific [³⁵S]GTP_γS binding to membranes of samples from autopsies (■) and from patients with mTLE (□) as a function of increasing concentration of the 5-HT_{1A} receptor agonist 8-OH-DPAT. Each point represents the mean ± S.E.M of the percentage of stimulation over basal values. Note that in patients with mTLE, [³⁵S]-GTP_γS binding stimulation by 8-OH-DPAT was significantly different from that of autopsy samples.

correlation ($p < 0.05$) of E_{\max} and pEC_{50} with the obtained values of patient's age, age of seizure onset, duration of epilepsy and frequency of seizures.

Results

Autopsy samples

Autopsy samples were acquired from five subjects (two male and three female) ranging in age from 29 to 71 years (45.8 ± 7.2 years), who died by different causes and without history of neurological disease. The postmortem interval was 3 to 6 h (4.6 ± 0.5 h) (Table 1). 8-OH-DPAT-stimulated [³⁵S]GTP_γS binding was concentration-dependent and saturable. The binding assays revealed a E_{\max} of 65.9% and a potency pEC_{50} value within the micromolar range (-7.3 ± 0.3) (Fig. 2). We found no significant correlation in autopsy samples between the coefficients of the E_{\max} of [³⁵S]GTP_γS binding and pEC_{50} values, with the age ($r = 0.7453$, $r = -0.0989$, respectively) and postmortem interval ($r = 0.7092$, $r = -0.0305$, respectively) (Table 2).

Patients group

The lateral temporal neocortex tissues were obtained from five patients (two male and three female) of age 38.2 ± 2.9 years old (ranged from 29 to 45 years) with 25.6 ± 2.6 years

of epilepsy duration; the age of seizure onset was 12.6 ± 3.3 years and they had 22.8 ± 7.5 seizures per month (Table 1). We found no significant differences in age between mTLE patients and the autopsy samples (38.2 ± 2.9 and 45.8 ± 7.2 years, respectively, $p = 0.1766$). Therefore, we found that E_{\max} values significantly decreased in the mTLE group (30.8%, $p < 0.05$) when compared to autopsy samples (65.9%) (Fig. 2). However, the pEC_{50} values in the micro molar range were similar in both groups (-7.3 ± 0.3 and -7.2 ± 0.1 , autopsy samples and mTLE, respectively). Regarding the correlation analysis, we found no significant differences between the E_{\max} with age of patient, age of seizure onset, duration of epilepsy and frequency of seizures. Also, the pEC_{50} values revealed no significant correlations with clinical data (Table 2).

Discussion

Even though the results showed a decrease of the 5-HT_{1A} receptor-induced G-protein functional activation in mTLE, we found no correlation between the decrease in functional coupling to the 5-HT_{1A} receptor with the patient's age, duration with epilepsy, age at onset with epilepsy, and frequency of seizures.

In the present study, we find limitations that describe below. Previous studies have indicated that 5-HT_{1A} receptor density decreases with age in brain regions such as cortex, hippocampal regions, locus coeruleus and dorsal raphe nuclei.^{30,31} These results suggest that a decrease in receptor density could indirectly produce changes in functional 5-HT_{1A} receptor activation. An important point to mention, we used tissue from mTLE surgical patients and autopsy samples from subjects of similar age, in order to reduce the possible influence of this variable. Also, our results show no significant differences in age between mTLE patients and the autopsy samples neither correlation with 5-HT_{1A} receptor-induced G-protein functional activation. Our results suggest that decrease functional 5-HT_{1A} receptor activation is directly related to changes associated with epilepsy rather than age.

Concerning the postmortem interval of the autopsy samples (4.6 ± 0.5 h) may produce changes that could alter the results, in this context, previous studies indicate that 5-HT_{1A} receptors are preserved within a postmortem interval of 8–92 h.³² Also, we results showed that postmortem interval did not correlate with the 5-HT_{1A} receptor-induced G-protein functional activation. These previous studies and

Table 2 Correlations of [³⁵S]GTP_γS binding parameters between clinical data and data from the autopsy sample from neocortex temporal membranes in patients with temporal lobe epilepsy.

[³⁵ S]GTP _γ S binding assay	Clinical data				Data from the autopsy samples	
	Age of patient	Seizure onset age	Duration of epilepsy	Frequency of seizures	Age of death	Postmortem interval
E_{\max}	0.3906	-0.0991	0.5587	0.1898	0.7453	0.7092
pEC_{50}	-0.6604	-0.0280	-0.6997	-0.7113	-0.0989	-0.0305

Values represent Pearson Correlation Coefficients. E_{\max} values (expressed as maximal stimulation of [³⁵S]GTP_γS binding) and pEC_{50} values (expressed as the negative logarithm of EC_{50}).

results obtained lead us to suggest that tissue from autopsy samples can be used as control.

Another limitation was the fabric storage time at -70°C . It has been reported that storage intervals of 1–82 months do not affect the functional activity of the 5-HT_{1A} receptor.²⁶ In our report the storage time was 4–12 months.

Regarding our results that showed a decrease in functional 5-HT_{1A} receptor activation in patients with mTLE can be explained as a consequence of changes in density of receptor in neocortex. It is known that the cortical layers (I–II) of the neocortex present a dense pattern of these receptors.³³ Rocha et al. (2007)²² showed decreased of 5-HT_{1A} receptor binding in the external cortical layers (I–II) of patients with mTLE. Other studies with PET have shown decrease the 5-HT_{1A} receptor binding in neocortex of patient with mTLE.^{19–21} These evidences may partly explain the decrease in the receptors functional activity in our results.

It is important to mention that 5-HT_{1A} receptor-induced G-protein functional activation induces hyperpolarization *via* potassium channels and therefore inhibitory effects.^{34–36} Therefore, the decreased functional activity of 5-HT_{1A} receptors associated with reduced density of these receptors in neocortex; it could explain the excitability and facilitation of seizure activity in neocortex of patients with mTLE. On the contrary, study in epileptic hippocampal tissue show an increase of the 5-HT_{1A} receptor-induced G-protein functional activation³⁷ suggest as a compensatory mechanism in response to the epileptic discharges that could contribute to the modulation of neuronal hyperexcitability.

The results of the present study indicate that patients with mTLE do not observe correlation with clinical aspects. These results suggest that clinical aspects do not play an important role in changes in the attachment of G $\alpha_{i/o}$ proteins to 5-HT_{1A} receptors. However, we cannot rule out other factors such as antiepileptic drugs, cortical atrophy or other processes underlying epilepsy may be involved in the observed alterations in functional activity.

The development of drugs that target the activation of GPCR receptors specifically those coupled to G $\alpha_{i/o}$ proteins has recently gained new interest for its efficacy in preclinical models by their anticonvulsive, antiepileptic, and neuroprotective effects. However, it is important to consider that at present is insufficient evidence of the alterations in coupled to G $\alpha_{i/o}$ proteins have in pharmacoresistance in patients with epilepsy. These studies in surgically resected tissue of patients with pharmacoresistant epilepsy will allow us to investigate and provide the scientific basis for these alterations in epilepsy, thus leading to know whether they can be used as a therapeutic target to treat epilepsy in the future.

Conflict of interest statement

None of the authors has any conflict of interest to disclose.

References

- Gilman AG. G proteins: transducers of receptor-generated signals. *Ann Rev Biochem.* 1987;56:615–49.
- Sunahara RK, Dessauer CW, Gilman AG. Complexity and diversity of mammalian adenylyl cyclases. *Annu Rev Pharmacol Toxicol.* 1996;36:461–80.
- Fisher RS, Van Emde Boas W, Blume W, et al. Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia.* 2005;46:470–2.
- Holmes GL. Role of glutamate and GABA in the pathophysiology of epilepsy. *Ment Retard Dev Disabil Res Rev.* 1995;1:208–19.
- DeFelipe J, Marco P, Sola RG, et al. Local changes in GTP-binding protein immunoreactivities in human epileptogenic neocortex. *Exp Brain Res.* 1998;119:153–8.
- Iwasa H, Kikuchi S, Miyagishima H, et al. Altered expression levels of G protein subclass mRNAs in various seizure stages of the kindling model. *Brain Res.* 1999;818:570–4.
- Iwasa H, Kikuchi S, Mine S, et al. Functional significance of stimulatory GTP-binding protein in hippocampus is associated with kindling-elicited epileptogenesis. *Psychiatry Clin Neurosci.* 2000;54:191–7.
- Potier S, Sénécal J, Chabot JG, et al. A pentylenetetrazole-induced generalized seizure in early life enhances the efficacy of muscarinic receptor coupling to G-protein in hippocampus and neocortex of adult rat. *Eur J Neurosci.* 2005;21:1828–36.
- Watanabe Y, Mori N, Uemura S. Suppression of kindled seizure following intraamygdaloid injection of pertussis toxin rats. *Neurosci Lett.* 1991;130:199–202.
- Snead OC. Evidence for G protein modulation of experimental-generalized absence seizures in rat. *Neurosci Lett.* 1992;148:15–8.
- Bruno V, Battaglia G, Ksiazek I, et al. Selective activation of mGlu4 metabotropic glutamate receptors is protective against excitotoxic neuronal death. *J Neurosci.* 2002;20:6413–20.
- Caulder EH, Riegle MA, Godwin DW. Activation of Group 2 metabotropic glutamate receptors reduces behavioral and electrographic correlates of pilocarpine induced status epilepticus. *Epilepsy Res.* 2014;108:171–81.
- Rocha L, Orozco-Suarez S, Alonso-Vanegas M, et al. Temporal lobe epilepsy causes selective changes in mu opioid and nociceptin receptor binding and functional coupling to G-proteins in human temporal neocortex. *Neurobiol Dis.* 2009;35:466–73.
- Cuellar-Herrera M, Velasco AL, Velasco F, et al. Mu opioid receptor mRNA expression, binding, and functional coupling to G-proteins in human epileptic hippocampus. *Hippocampus.* 2012;22:122–7.
- Bañuelos-Cabrera I, Cuellar-Herrera M, Velasco AL, et al. Pharmacoresistant temporal lobe epilepsy modifies histamine turnover and H3 receptor function in the human hippocampus and temporal neocortex. *Epilepsia.* 2016;57:e76–80.
- Salgado-Commissariat D, Alkadhhi KA. Serotonin inhibits epileptiform discharge by activation of 5-HT_{1A} receptors in CA1 pyramidal neurons. *Neuropharmacology.* 1997;36:1705–12.
- Wada Y, Nakamura M, Hasegawa H, et al. Role of serotonin receptor subtype in seizures kindled from the feline hippocampus. *Neurosci Lett.* 1992;141:21–4.
- López-Meraz ML, Martínez A, Rocha L. Effect of 8-OH-DPAT on electrographic activity during the kainic acid-induced status epilepticus in rats. *Seizure.* 2007;16:365–70.
- Toczek MT, Carson RE, Lang L, et al. PET imaging of 5-HT_{1A} receptor binding in patients with temporal lobe epilepsy. *Neurology.* 2003;60:749–56.
- Savic I, Lindström P, Gulyás B, et al. Limbic reductions of 5-HT_{1A} receptor binding in human temporal lobe epilepsy. *Neurology.* 2004;62:1343–51.
- Giovacchini G, Toczek MT, Bonwetsch R, et al. 5-HT_{1A} receptors are reduced in temporal lobe epilepsy after partial-volume correction. *J Nucl Med.* 2005;46:1128–35.
- Rocha L, Lorigados-Pedre L, Orozco-Suárez S, et al. Autoradiography reveals selective changes in serotonin binding in neocortex of patients with temporal lobe epilepsy. *Prog Neuro-Psychopharmacol Biol Psychiatry.* 2007;31:1208–18.

23. Chagnac-Amitai Y, Connors BW. Horizontal spread of synchronized activity in neocortex and its control by GABA-mediated inhibition. *J Neurophysiol.* 1989;61:747–58.
24. Chervin RD, Pierce PA, Connors BW. Periodicity and directionality in the propagation of epileptiform discharges across neocortex. *J Neurophysiol.* 1988;60:1695–713.
25. Marco P, DeFelipe J. Altered synaptic circuitry in the human temporal neocortex removed from epileptic patients. *Exp Brain Res.* 1997;114:1–10.
26. Gonzalez-Maeso J, Rodriguez-Puertas R, Gabilondo AM, et al. Characterization of receptor-mediated (35S)GTPγS binding to cortical membranes from postmortem human brain. *Eur J Pharmacol.* 2000;390:25–36.
27. Benyhe S, Farkas J, Tóth G, et al. Met5-enkephalin-Arg6-Phe7, an endogenous neuropeptide, binds to multiple opioid and nonopioid sites in rat brain. *J Neurosci Res.* 1997;48:249–58.
28. Lowry OH, Rosebrough NJ, Farr L, et al. Protein measurement with the Folin phenol reagent. *J Biol Chem.* 1951;193:265–75.
29. Spetea M, Monory K, Tomboly C, et al. In vitro binding and signaling profile of the novel mu opioid receptor agonist endomorphin 2 in rat brain membranes. *Biochem Biophys Res Commun.* 1998;250:720–5.
30. Dillon KA, Gross-Isseroff R, Israeli M, et al. Autoradiographic analysis of serotonin 5-HT1A receptor binding in the human brain postmortem: effects of age and alcohol. *Brain Res.* 1991;554:56–64.
31. Burnet PWJ, Eastwood SL, Harrison PJ. Detection and quantitation of 5-HT1A and 5-HT2A receptor mRNAs in human hippocampus using a reverse transcriptase-polymerase chain reaction (RT-PCR) technique and their correlation with binding site densities and age. *Neurosci Lett.* 1994;178:85–9.
32. González-Maeso J, Torre I, Rodríguez-Puertas R, et al. Effects of age, postmortem delay and storage time on receptor-mediated activation of G-proteins in human brain. *Neuropsychopharmacology.* 2002;26:468–78.
33. Palchoudhuri M, Flügge G. 5-HT1A receptor expression in pyramidal neurons of cortical and limbic brain regions. *Cell Tissue Res.* 2005;321:159–72.
34. Bobula B, Zahorodna A, Bijak M. Different receptor subtypes are involved in the serotonin-induced modulation of epileptiform activity in rat frontal cortex in vitro. *J Physiol Pharmacol.* 2001;52:265–74.
35. Lanfumey L, Hamon M. 5-HT1 receptors. *Curr Drug Targets CNS Neurol Disord.* 2004;3:1–10.
36. Wada Y, Shiraishi J, Nakamura M, et al. Role of serotonin receptor subtypes in the development of amygdaloid kindling in rats. *Brain Res.* 1997;747:338–42.
37. Cuellar-Herrera M, Velasco AL, Velasco F, et al. Alterations of 5-HT1A receptor-induced G-protein functional activation and relationship to memory deficits in patients with pharmacoresistant temporal lobe epilepsy. *Epilepsy Res.* 2014;108:1853–63.