



REVIEW

Endoscopic ultrasound and endoscopic retrograde cholangiopancreatography: Can they be successfully combined?



Joan B. Gornals^{a,*}, José Miguel Esteban^b, Carlos Guarner-Argente^c,
Carlos Marra-Lopez^d, Alejandro Repiso^e, Oriol Sendino^f, Carme Loras^g

^a Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge-IDIBELL, Barcelona, Catalonia, Spain

^b Endoscopy Unit, Department of Digestive Diseases, Hospital Clinic San Carlos, Madrid, Spain

^c Endoscopy Unit, Department of Digestive Diseases, Hospital Santa Creu i Sant Pau, Barcelona, Catalonia, Spain

^d Endoscopy Unit, Department of Digestive Diseases, Hospital Universitario Araba sede Txagorritxu, Alava, Spain

^e Endoscopy Unit, Department of Digestive Diseases, Hospital Virgen de la Salud, Toledo, Spain

^f Endoscopy Unit, Department of Digestive Diseases, Hospital Clinic, Barcelona, Catalonia, Spain

^g Endoscopy Unit, Department of Digestive Diseases, Hospital Mútua Terrassa, CIBEREHD, Terrassa, Catalonia, Spain

Available online 23 February 2016

KEYWORDS

Endoscopic retrograde cholangiopancreatography; Single-session procedure; Endoscopic ultrasound; Biliary drainage; Endoscopy; Digestive system; Therapeutics

Abstract Endoscopic ultrasound (EUS) and endoscopic retrograde cholangiopancreatography (ERCP) have much in common, including their main indications (biliopancreatic disorders), powerful therapeutic capacities and a steep learning curve. Over the years they have evolved from novel diagnostic procedures to interventional therapeutic techniques, but along different paths (different scopes or devices and endoscopists specializing exclusively in one or the other technique). However, EUS has gradually developed into a therapeutic technique that requires skills in the use of ERCP devices and stents, leading some ERCP specialists to explore the therapeutic potential of EUS. The corresponding literature, which has grown exponentially, includes recent experiments on combining the two techniques, which have gradually come to be used in routine care in a number of centers, with positive technical, clinical and financial outcomes. We review EUS and ERCP as individual or combined procedures for managing biliopancreatic disorders. © 2016 Elsevier España, S.L.U., AEEH y AEG. All rights reserved.

PALABRAS CLAVE

Colangiopancreatografía retrógrada endoscópica; Procedimiento en una sola sesión;

Ecografía endoscópica y colangiopancreatografía retrógrada endoscópica: ¿Es posible combinarlas con éxito?

Resumen La ecografía endoscópica (EE) y la colangiopancreatografía retrógrada endoscópica (CPRE) comparten muchas cosas, tales como sus indicaciones principales (las enfermedades biliopancreáticas) y potentes capacidades terapéuticas, y una empinada curva de aprendizaje.

* Corresponding author.

E-mail address: jgornals@bellvitgehospital.cat (J.B. Gornals).

Ecografía
endoscópica;
Drenaje biliar;
Endoscopia;
Aparato digestivo;
Terapéutica

Con el paso del tiempo, estos procedimientos diagnósticos novedosos han evolucionado por diferentes vías (distintos ámbitos o dispositivos y endoscopistas especializados exclusivamente en una u otra técnica), hasta convertirse en técnicas terapéuticas intervencionistas. Sin embargo, de manera gradual, la EE ha llegado a ser una técnica terapéutica que requiere habilidades en el manejo de los instrumentos y stents que se emplean en la CPRE, lo que ha conducido a algunos especialistas en CPRE a explorar el potencial terapéutico de la EE. Las publicaciones relativas a este tema, que han crecido de forma exponencial, incluyen experimentos recientes de combinación de estas técnicas, que diversos centros han introducido progresivamente en sus protocolos de atención rutinaria, con resultados técnicos, clínicos y económicos positivos. Hemos revisado la EE y la CPRE como procedimientos individuales o combinados en el tratamiento de enfermedades biliopancreáticas.

© 2016 Elsevier España, S.L.U., AEEH y AEG. Todos los derechos reservados.

Introduction

Endoscopic Ultrasound (EUS) and Endoscopic Retrograde Cholangiopancreatography (ERCP) are advanced endoscopic procedures. The two techniques have similarities and differences. ERCP is older, indications are purely focused on biliopancreatic disorders, and it is mainly a therapeutic intervention procedure. On the other hand, EUS is a more modern technique, indications are wider and more varied (mediastinal and rectal lesions, staging of GI lesions, subepithelial lesions), and it is mainly a diagnostic procedure, including interventional diagnostic procedures such as EUS-guided puncture with fine needle aspiration. EUS and ERCP have many similarities: a demanding learning curve, a common main indication (biliopancreatic disorders), and powerful therapeutic capacity.¹

In recent years, the literature has included experiments that combine the two techniques, and in some centers they have been introduced naturally in routine care. The positive effects of this combination are technical, clinical, logistical, and financial.²

This review focuses on the relationship between the two techniques, firstly considering them individually, and secondly as competing procedures, with a final look at combined procedures.

Endoscopic ultrasound

The learning curve for EUS and EUS-guided fine needle aspiration (FNA)

As the applications for EUS have become increasingly recognized by other clinical practitioners, the demand for well-trained endosonographers has increased.³ The limited availability of EUS is largely the result of a lack of skilled endosonographers. A relative lack of training centers combined with the extensive commitment required by the trainee has limited the growth of EUS and its availability in community practice. For most trainees, the amount of EUS exposure and training is highly variable and often

program-dependent. Many fellowship programs do not provide the opportunity to learn EUS.

EUS is operator-dependent. The American Society for Gastrointestinal Endoscopy (ASGE) recommends a minimum of 150 total supervised procedures, 75 of which have a pancreatobiliary indication and 50 cases of FNA (25 of which are pancreatic FNA) before competency can be determined.⁴ Other studies by experts claim that the number of procedures must be twice the ASGE recommendations.⁵⁻⁷

The learning process of EUS-FNA has been studied for solid pancreatic lesions and has shown a learning curve with increasing sensitivity in the cytopathological diagnosis of cancer (reaching 80% after 20-30 EUS-FNA), with a decreasing number of passes needed to obtain adequate results (reaching a median of 3 after 150 EUS-FNA) but without variation in severe morbidity.^{8,9}

Trainees should demonstrate competence in linear EUS before undertaking EUS-FNA.^{10,11} The Technical Guidelines of the European Society of Gastrointestinal Endoscopy (ESGE) recommend the use of a combination of different simulators and, if available, live pigs during training in EUS-FNA. They recommend that a minimum of 20-30 supervised non-pancreatic and pancreatic lesions, respectively, be performed with rapid on-site cytopathological examination (ROSE).¹² The clinical effectiveness of EUS and EUS-FNA depends on the judicious use of these techniques and the skill of the endosonographer, but also on the indicators established by the ASGE to aid in the recognition of high quality EUS examinations.¹³

General review (overview) of EUS indications: diagnostic, interventional, and therapeutic

EUS continues to evolve as a diagnostic and therapeutic technique. EUS should be performed when it has the potential to affect patient management, such as when establishing a diagnosis, performing locoregional tumor staging, and providing therapeutic interventions. Since the introduction of EUS in 1980, its indications and role have continued to expand.

We must differentiate between diagnostic and therapeutic indications. The indications for EUS are determined by

Table 1 Diagnostic indications for EUS.

- *GI tract cancer staging*
Gastroesophageal cancer
Rectal cancer
Ampullary cancer
- *Mass imaging and delineation of*
Pancreas
Bile ducts
Mediastinum (including lung cancer)
- *Evaluating abnormalities of the GI tract wall or adjacent structures*
- *Tissue sampling of lesions within, or adjacent to, the wall of the GI tract*
- *Evaluation of abnormalities of the pancreas and biliary tree*
Masses
Dilations
Lithiasis
- *Evaluation of patients at increased risk of pancreatic cancer*

the anatomic conditions and the technical capabilities of the equipment. The primary role of EUS is to delineate gastrointestinal tract (GIT) and immediate surroundings to a depth of 4–5 cm.¹⁴ Table 1 summarizes diagnostic indications. These indications can be divided between GIT cancer staging, mass imaging and delineation, evaluation of abnormalities of the pancreas and biliary tree, and study and evaluation of the mediastinum of patients at increased risk of pancreatic cancer.

The ability to pass a hollow needle under ultrasound guidance has expanded the applications of EUS. The needle is essentially a conduit that allows for the passage or placement of materials with therapeutic intent. The EUS therapeutic interventions have continued evolving since the beginning of the EUS-guided celiac plexus neurolysis or block, followed by the EUS-guided pseudocyst drainage. Table 2 summarizes the therapeutic indications at present. But perhaps in the near future some other indications that are now on evaluation will be established. New indications are evolving and now only expert centers are using them, or they are on evaluation in animal labs. Some of these indications have been published with small groups of patients included.

Table 2 Therapeutic indications for EUS on evaluation.

- *EUS-guided FNI into tumors*
Chemotherapy
Antitumor agents
- *EUS-guided FNI vascular interventions*
Sclerosant agents
Coil application
Intrahepatic portosystemic shunts
- *Drainage of the gallbladder*
- *EUS-guided anastomosis*
- *EUS-guided photodynamic therapy*
- *EUS-guided brachytherapy*

Endoscopic retrograde cholangiopancreatography

The learning curve for therapeutic ERCP

Implementing a standardized protocol for an ERCP training program is very difficult, and might not be feasible in practice. Procedure volume, indications, and technical approaches vary widely among different countries, institutions, and even endoscopists in a same institution.

Competence has been classically evaluated according to a minimum number of procedures performed. Based on a first prospective study,¹⁵ the ASGE recommended a minimum of 180–200 ERCPs. The number of procedures recommended in most training programs ranges from 100 to 200. Current research supports establishing a standard of 80–90% technical success before trainees are deemed competent in a specific skill.^{16,17} Nevertheless, individual trainees may differ in the acquisition of technical skills. Therefore these threshold numbers might be inadequate.

Recently, a study reported the use of a standardized form for continuous self-assessment.¹⁸ The form included previously proposed quality indicators for ERCP such as procedural indication, degree of technical difficulty, previous ERCP failure, and success or failure options (such as cannulation of the CBD or pancreatic duct, stent placement, sphincterotomy, and stone extraction). It also comprised an improvement plan for self-evaluation every 10 procedures. This method allowed not only for the quantitative evaluation of trainees, but also determined the learning curve of each individual and of the average group progression. As such, current training should probably be based on learning curves, although more studies are needed to validate these resources.

General overview of ERCP indications

ERCP was introduced more than four decades ago as a diagnostic procedure, but only evolved into therapeutic endoscopy once sphincterotomy was introduced. Nowadays, due to the technological evolution of non-invasive tests (i.e. imaging tests or EUS), and taking into account potentials complications, this technique should be reserved almost exclusively for therapeutic indications.

ERCP should be performed for appropriate indications as defined in previously published guidelines for >90% of the procedures.¹⁶ When there is a nonstandard indication, the reasons for this should be made sufficiently clear. Therefore, the indication has to be carefully evaluated, and radiological studies reviewed to anticipate the therapeutic strategy.¹⁶ The main indications have been previously described in detail, and they are summarized in Table 3.^{19–22} We will briefly review the most common indications.

ERCP for biliary diseases

Choledocholithiasis

Choledocholithiasis is the most common cause of obstruction of the bile duct and ERCP has sensitivity greater than 95% for diagnosis. ERCP can be performed after cholecystectomy.

Table 3 ERCP indications.

Biliary diseases	Pancreatic diseases
<ul style="list-style-type: none"> • <i>Cholelithiasis</i> 	<ul style="list-style-type: none"> • <i>Recurrent idiopathic acute pancreatitis</i>
<ul style="list-style-type: none"> • <i>Benign biliary stricture</i> Inflammatory Congenital Postoperative 	<ul style="list-style-type: none"> • <i>Chronic pancreatitis</i> Stricture Lithiasis
<ul style="list-style-type: none"> • <i>Malignant biliary stricture</i> 	<ul style="list-style-type: none"> • <i>Pancreatic duct leak</i> • <i>Pancreatic collections (pseudocysts, disconnected pancreatic duct syndrome)</i>
Pancreatic cancer Cholangiocarcinoma Ampullary lesions Infiltration by other neoplasias, lymph nodes or metastatic disease	
<ul style="list-style-type: none"> • <i>Bile leak</i> • <i>Sphincter of Oddi dysfunction</i> • <i>Rare indications: hemobilia, infectious (hydatidosis or others)</i> 	

However, in patients with jaundice, elevated liver enzymes, or worsening or persistent pancreatitis or cholangitis, it should be considered preoperatively, especially if a magnetic resonance cholangiopancreatography (MRCP) or EUS demonstrates choledocholithiasis. In addition, it should be performed urgently in patients with severe acute cholangitis, and those with severe acute biliary pancreatitis with suspected persistence of impacted lithiasis.

Biliary strictures

For malignant strictures, drainage with plastic and metal stents is effective in up to 90% of cases of the stenosis in the middle or distal choledochus. Nevertheless, ERCP is not generally indicated to relieve a biliary obstruction in patients with potentially resectable malignant distal bile duct obstruction, in whom surgical resection will not be delayed by neoadjuvant therapy or other preoperative assessments or treatments. Despite this, in current clinical practice preoperative biliary decompression is widely performed. Drainage of more proximal lesions, such as Klatskin tumors, is usually less successful due to incomplete biliary drainage.

For benign biliary strictures different dilation or stenting can be effective, although treatment success varies widely depending on the indication.

A special indication is primary sclerosing cholangitis. There is usually a good response to dilatation. However, there is also an increased risk of developing acute cholangitis. For this reason, indications need to be individualized and considered carefully. Additionally, cytology of the dominant stricture should always be performed.

Bile leak

Sphincterotomy alone may be sufficient. Combination with a plastic biliary stent is recommended for large or highly productive leaks. The treatment is successful 80–100% of cases, depending on size and location.

Sphincter of Oddi dysfunction

Patients with sphincter of Oddi type I dysfunction respond in as many as 90% of cases to sphincterotomy. For type II dysfunction, clinical response is usually associated with abnormal manometry. However, many groups perform this treatment empirically. ERCP is not recommended for type III dysfunction.

ERCP for pancreatic diseases^{23,24}

Acute recurrent pancreatitis

EUS and MRCP allow non-invasive diagnosis of anatomical pancreatic abnormalities and other etiologies of pancreatitis such as microlithiasis. As such, ERCP is generally reserved for treatment procedures. Occasionally, ERCP may be required to obtain the pancreatic duct anatomy or to collect bile for the study of microlithiasis. For selected patients with pancreas divisum, minor papilla sphincterotomy might be effective.

Chronic pancreatitis

ERCP is useful for the diagnosis and treatment of pancreatic duct lithiasis, strictures, and pseudocysts. Fragmentation and extraction of pancreatic duct lithiasis is often difficult or even impossible, and it seems less effective than surgical treatment. New approaches with direct cholangioscopy and laser or electrohydraulic lithotripsy might improve the endoscopic management of this situation.

Pancreatic leaks

Transpapillary pancreatic stents can treat pancreatic leaks. However, it is advisable to place larger stents to bridge the leak when feasible, as this is associated with better outcomes.

EUS or ERCP

For patients with suspected choledocholithiasis (or with intermediate probability of bile duct stones)

Cholelithiasis is quite prevalent in most western countries. Prevalence ranges between 10% and 20% while incidence increases with age.²⁵ However cholelithiasis is asymptomatic in 80% of patients. Among these patients, passage of gallstones into the common bile duct (CBD) stones or choledocholithiasis occurs in 15–20%.²⁶

When CBD stones are demonstrated prior to cholecystectomy, patients should undergo stone extraction, either by ERCP or by intraoperative bile duct examination during cholecystectomy.²⁷

The diagnosis of choledocholithiasis is usually made by cholangiography, either preoperatively with ERCP, MRCP, or EUS, or intraoperatively (IOC) at the time of cholecystectomy.^{27–30}

A combination of clinical, biochemical, and morphological criteria is used to risk-stratify individuals with suspected choledocholithiasis and an intact gallbladder into low, intermediate and high risk of choledocholithiasis groups.²⁹ In patients with high risk of choledocholithiasis, ERCP remains the gold standard given its dual diagnostic and therapeutic role, even though it carries a risk of complications. At least two studies have analyzed the performance of ERCP as a diagnostic test and compared the performance of ERCP to that of EUS, along with ERCP vs intraductal ultrasound for CBD stone detection in patients at high risk of choledocholithiasis. ERCP showed 89–93% sensitivity, 100% specificity, and 94% accuracy for choledocholithiasis.^{31,32} Patients with intermediate risk of choledocholithiasis benefit the most from a non-invasive diagnostic evaluation of their CBD by EUS or MRCP, based on which a decision is then made to perform a therapeutic ERCP.^{28,29} EUS is an accurate test for diagnosis of choledocholithiasis in patients who are at intermediate risk of having a CBD stone based on clinical predictors.³³

The performance of EUS for the evaluation for choledocholithiasis has been extensively studied. Two extensive meta-analyses reported 89–94% sensitivity and 94–95% specificity of EUS in detecting choledocholithiasis, with ERCP, IOC, or surgical exploration used as criterion standards.^{30,34}

It has been proposed that EUS could be used in selecting patients for therapeutic ERCP, avoiding diagnostic ERCP, given the higher morbidity of ERCP compared with EUS. For this reason, it has been suggested that EUS should replace diagnostic ERCP to assess patients with intermediate probability of choledocholithiasis.³⁵

EUS appears comparable to ERCP as a diagnostic test for CBD stones, is superior to other techniques for detecting biliary stones, and can be used to select patients who will need therapeutic ERCP. This results in a significantly lower risk of complications in comparison with the use of ERCP for both diagnosis and treatment of choledocholithiasis.^{30,36,40,41}

By performing EUS first, ERCP may be safely avoided in patients with common bile duct stones. Application of EUS in the selection of patients for therapeutic ERCP significantly reduces the complication rate.³⁵

However, there are only a few randomized studies in this setting.^{35–39} These 4 trials randomized patients at intermediate to high risk for choledocholithiasis to an EUS-first strategy versus an ERCP-first strategy. Patients found to have CBD stones at EUS underwent subsequent therapeutic ERCP, which was performed in the same setting in 3 of the 4 trials. Taken together, these studies provide evidence that the EUS-first strategy with selective therapeutic ERCP can reduce the number of diagnostic ERCPs by 60–75% in patients who are at moderate risk for choledocholithiasis.

It was also found that EUS first is more successful in evaluating the bile duct for stones because of possibly unsuccessful biliary cannulation with diagnostic ERCP. Finally, the studies in question suggest that there was either less morbidity or a trend toward less morbidity and fewer complications using EUS first to screen for CBD stones, rather than starting with ERCP.

For tissue diagnosis of biliary strictures (or tissue sampling in suspected malignant biliary obstruction)

It is difficult to differentiate between benign and malignant biliary strictures. In this regard, endoscopic tissue acquisition may obviate the need for further invasive testing, thus allowing for optimal intervention without delay.⁴²

Biliary brush cytology has been a mainstay diagnostic method for suspected extrahepatic biliary tree malignancies because it is technically easy and generally is safe. However, its sensitivity for cancer is modest, ranging from 30 to 57% in most published studies.^{42,43} The addition of endobiliary forceps and endoscopic needle aspiration to brush cytology during ERCP increases the sensitivity of tissue sampling, but these procedures are time-consuming, increase the procedural costs, and may not be technically feasible for some biliary strictures.⁴⁴

Some studies have documented EUS-FNA as a useful technique for the diagnosis of bile duct strictures, with a reported sensitivity of 43–86%^{45–48} (Fig. 1). In addition, three studies have attempted to use EUS-FNA on biliary strictures following negative or unsuccessful ERCP sampling. These studies showed that EUS-FNA was technically feasible,



Figure 1 EUS-guided FNA of a malignant biliary stricture. (A) Radial EUS, bile duct wall thickening in relation to a distal cholangiocarcinoma; (B) EUS-FNA; (C) ERCP reveals a distal biliary stricture.

Dr. A Repiso, H. Virgen de la Salud, Toledo.

Table 4 Overall sensitivity of EUS-FNA and ERCP-based tissue sampling for patients with biliary strictures.

	No. of patients	Study	Sensitivity (%)		p-value
			EUS-FNA	ERCP	
Rösch et al. ⁵²	50	Prospective	43	54	NS
Oppong et al. ⁵³	37	Retrospective	53	29	<0.05
Weilert et al. ⁵⁴	51	Prospective	94	50	<0.05

without significant risks, and with sensitivities after negative ERCP sampling of 68–89%.^{49–51}

Three studies have directly compared the diagnostic yield of EUS-FNA and ERCP-based tissue sampling in patients with suspected malignant biliary obstruction.^{52–54} They found that EUS-FNA had greater sensitivity compared with ERCP (Table 4). In these studies, the sensitivity and accuracy of EUS-FNA were significantly superior to ERCP sampling among patients with pancreatic masses; EUS-FNA directly samples pancreatic mass, whereas ERCP samples are usually obtained from the area where a mass is causing compression on the bile duct. However, the sensitivity and accuracy of EUS-FNA seemed comparable to or lower than those of ERCP tissue sampling among patients with biliary masses and strictures.^{52–54} Therefore, it appears reasonable, when a tissue diagnosis is required, to start with EUS-FNA when a biliopancreatic mass is suspected, and with ERCP in biliary strictures in the absence of a biliopancreatic mass.

One potential additional risk in EUS-FNA with malignant biliary strictures is peritoneal tumor seeding.⁵⁵ Although a published study suggested there was no adverse effect on overall or progression-free survival from preoperative EUS-FNA, the theoretical risk of tumor seeding must be considered.⁵⁶ Bile leakage during EUS-FNA may raise the risk of cancer-cell seeding in EUS-FNA performed for biliary lesions.⁵¹ Therefore, care must be taken not to puncture the lesion through the bile duct lumen, if possible, in order to avoid bile leakage.

EUS and ERCP

Single or separate sessions

EUS with or without FNA and ERCP are very often required for the management of pancreaticobiliary disorders, and they are usually performed in separate sessions.⁵⁷ The experience of combining both procedures in a single session is limited and it is not common in routine care. EUS is a safe and useful technique in the evaluation of pancreaticobiliary diseases, especially in the evaluation of small choledocholithiasis and small pancreatic tumors, allowing a locoregional staging for neoplastic lesions with the possibility of performing FNA for cytological diagnosis.^{58,59} Currently, ERCP should only be performed as a therapeutic technique for biliary drainage with or without stent insertion.^{60,61} In cases of failed transpapillary ERCP cannulation, EUS-guided interventions could provide access for direct drainage, as the procedure was first described in 1996.⁶² Therefore, the

combination of the two techniques in a single session offers the advantage of adding the potential of both techniques, allowing accurate diagnosis with endoscopic drainage.^{1,63} Despite the numerous potential advantages, implementation in clinical practice has encountered important obstacles regarding safety, diagnostic accuracy, and cost.

Table 5 provides a summary of the various papers published in relation to single-session EUS-ERCP in pancreaticobiliary diseases.^{36,63–76}

Combined EUS and ERCP for patients with suspected choledocholithiasis

Performing EUS before ERCP can prevent two thirds of unnecessary ERCPs.^{36,59} Although MRCP is as powerful as EUS in choledocholithiasis detection, we have to take into account that EUS is more accurate than MRCP in the detection of small CBD stones, and it is preferred in cases where MRCP is not possible, such as with patients with claustrophobia and patients with metal devices.^{36,77} As previously reported, it seems that performing EUS-ERCP in a single session for choledocholithiasis is a safe, feasible and effective strategy, with no increase in procedure- or sedation-related complications (Table 5). Moreover, in one study an increased risk of complications was noted when ERCP was delayed, resulting in significant biliary complications in 14% of patients.⁷¹ It also seems that a single session reduces hospital stay, costs and repeat sedation compared to separate sessions.^{36,66,72} A negative point of this strategy is the long procedure duration that may be especially problematic in elderly patients. As reported by Iles-Shih, a single session in elderly patients was not related with more adverse events than in nonelderly patients.⁷⁰ An important element in establishing this approach is having technology that allows the procedure scope without the need to change the scope. Although there is a study that did the whole procedure with the linear echoendoscopy, in real practice the cannulation of the papilla is not possible in the majority of the cases; improvement in the endoscopic devices is necessary.⁶³ Another situation in which a single-session EUS and ERCP would be useful is the case of pregnant patients with suspected choledocholithiasis. As reported by Vohra et al., a single-session procedure was done without fluoroscopy in 10 pregnant women. The whole procedure was done with two different tubes and EUS helped the study of the number, size and location of common bile stones.⁷⁶

Table 5 Summary of single-session EUS and ERCP procedures reported in the literature.

Author year (reference)	Combined procedures (EUS FNA)	Study	Pathology	FNA Accuracy (%)	ERCP details	Duration (min.)	Complications
Liu ³⁶	70 (0)	Comparative prospective	Benign	NA	100% BD cannulation rate (successful in 25 of 25)	–	7% (pancreatitis, cholangitis, 2 deaths)
Rocca ⁶³	19 (0)	Observational	Benign	NA	94% BD cannulation rate (successful in 15 of 16)	27 (22–36)	None
Tarantino ⁶⁴	72 (25)	Prospective observational	Malignant and benign (76–24%)	92	'All ERCP completed' (1 precut)	58.6 ± 9 (30–91)	8% 2 mild bleeding
Ross ⁶⁵	114 (87)	Retrospective	Malignant and benign (70–30%)	87.8	84% B stent placement (in 96 patients) 51 sphincterotomies	73.6 ± 30 (25–148)	10.5% 6 pancreatitis
Fabbri ⁶⁶	40 (0)	Comparative Prospective	Benign	NA	–	59.3 ± 8	2.5% 1 mild pancreatitis
Ascunce ⁶⁷	35 (28)	Prospective observational	Malignant and benign (80/20%)	96	97.1% BD cannulation rate 5 precut 82.9% stents deployed (29 patients)	83.7	2% 1 bleeding
Vila ⁶⁸	39 (19)	Comparative retrospective	Malignant and benign (55–45%)	90.6	–	93 ± 32.78	5% (SSG) 1 pancreatitis, 1 bleeding
Aslanian ⁶⁹	29 (24)	Comparative retrospective	Malignant	92	72% B stent placement	75 (60–90)	3.4% (SSG) 1 bleeding
Iles-Shih ⁷⁰	206 (110)	Retrospective	Malignant and benign (70–25%)	–	–	56 ± 2.1 58 ± 1.8	1 immediate AE 20 long-term AE (9%)
Benjaminov ⁷¹	71 (0)	Comparative retrospective	Benign	NA	98.5% BD cannulation rate	–	15.7% (SSG) minor complications
Gornals ⁷²	53	Retrospective	Malignant and benign (68–32%)	90	67% BD cannulation rate, 89% BD drainage (wEUS) 60% B sent placement	65.5 ± 3	5.6% complications (1 perforation/2 cholangitis)
Chu ⁷³	60 (14)	Comparative prospective	Malignant and benign	81.8	98.3% BD cannulation rate 31.6% B stent placement	70.05 ± 15.35	8% complications (1 pancreatitis/4 hypoxemia)
Weilert ⁵⁴	51 (51)	Prospective	Malignant	94	86% BD cannulation rate 38% B stent placement	70 (40–126)	None
Kawakubo ⁷⁴	68 (30)	Retrospective	Malignant and benign	100	98.5% BD cannulation rate 71% B stent placement	75 (53–88)	8.8% pancreatitis, 1.5% bleeding
Noma ⁷⁵	100 (100)	Prospective	Malignant	91	97% BD cannulation rate 98% PD cannulation rate	61 (27–97)	10% pancreatitis
Vohra ⁷⁶	6 (0)	Retrospective (pregnant patients)	Benign	NA	100% BD cannulation rate	33 (20–42)	None

AE: adverse event; B: biliary; BD: biliary duct; PD: pancreatic duct; NA: not applicable; SSG: single session group; wEUS: with EUS helping.

Combined EUS-guided FNA and ERCP for obstructive jaundice from presumed biliopancreatic malignancy

EUS is highly sensitive for the detection of pancreatic tumors with a very high negative predictive value and with the ability to provide FNA.⁷⁸ Establishing a tissue diagnosis of malignancy is mandatory prior to surgical or oncologic treatment in patients with suspected malignant biliary obstruction. ERCP is a well-established procedure for evaluating and managing biliary obstructions. These two procedures are often required for the diagnosis and treatment of biliopancreatic malignancy. Therefore it is reasonable to perform these two techniques during a single session under the same anesthesia. Based on the published data, it seems that combined EUS-FNA and ERCP for obstructive jaundice is a safe, feasible procedure (Table 5).

Regarding which technique is better to perform first, some authors advocate beginning with ERCP and stent placement, to improve the EUS study of biliary system using the stent as a guide.⁵⁷ In contrast, other evidence supports performing EUS prior to ERCP, as it is more accurate for cancer staging^{72,79,80} and because EUS-FNA has greater sensitivity than ERCP for detecting malignancy, especially in primary pancreatic cancer.⁵⁴ This strategy could avoid unnecessary ERCP, brushings and stents, and decrease the possibility of inconclusive cytology in the case of stent placement. Moreover, as reported in several papers, the sensitivity and accuracy of FNA are not affected when the two procedures are done in the same session (Table 5).

Regarding safety, although initial experiences reported severe complications related to EUS-FNA prior to ERCP, with the consequence of contrast leakage and pneumoperitoneum,^{81,82} this has not been observed in later studies with series of patients ranging from 19 to 206 (Table 5).

Regarding procedure duration, as previously noted it does not seem to be the case that a longer procedure is associated with more cardiopulmonary adverse events.^{65,70,72} Moreover in those studies that have compared the procedure time of the two techniques in a separate session versus in a single sessions, it seems that there are no differences or even a timesaving in the single session strategy.^{66,68,69,73} In these previous comparative studies similar results were obtained, with the dose of propofol administered being better for the single session group.

Another important issue to be considered is the financial. As we mentioned above, for benign disease there have been few cost-analysis studies performed, and these show a clear benefit in single-session EUS-ERCP.^{36,66,72}

In conclusion, it seems that a single-session EUS-ERCP for pancreaticobiliary diseases provides benefits in clinical and financial terms, by simplifying patient management, shortening hospital stay, reducing cost, and avoiding repeat sedation. The procedure time and the logistical possibility of performing the two techniques at the same time are the principal limiting factors for applying this strategy in real practice.

Biliary and pancreatic drainage guided by EUS after failed ERCP

ERCP is the preferred procedure for biliary and pancreatic drainage. However failure occurs in 3–10% of cases due to anatomic variation, prior surgery, operator inexperience, tumor extension, or periampullary pathology. In these situations, the rescue options include precut, percutaneous transhepatic drainage (PTBD), surgical intervention, EUS-guided biliary drainage (BD), and pancreatic duct (PD) drainage.^{68,78,83–85}

Biliary drainage guided by EUS

EUS-guided biliary drainage (EUS-BD) has been described as an alternative method to achieve internal biliary drainage in those patients in whom ERCP has failed. This is a promising technique which offers some advantages over the other options (radiological or surgery). Multiple retrospective and some prospective studies have shown it to be safe and effective. But it is important to note that all these data come from highly skilled advanced endoscopists at tertiary centers with expertise in both EUS and therapeutic endoscopy.

EUS-guided BD has been performed in more than 1000 published cases, with mean technical and clinical success rates of 91% and 88%, respectively. However, the mean overall complication rate was 26% (16–35%) with mortality of 1–5%, especially in the early stage of the learning-curve (Table 6).^{68,72,85–110}

The nomenclature of the different variations of this technique can be confusing and repetitive. In short, there are two different routes: intrahepatic and extrahepatic. After bile duct access, drainage can be accomplished via (A) direct transgastric transluminal stenting (i.e., hepaticogastrotomy) or transduodenally (i.e., choledochoduodenostomy); (B) rendezvous technique passing a guidewire through to the papilla; and (C) antegrade stent placement, similar to the PTBD technique.

When these different methods are compared, EUS-guided BD shows similar technical and clinical success rates to both intrahepatic (usually transgastric) and extrahepatic (usually transduodenal) access. However, the extrahepatic route seems to be safer than the intrahepatic.^{84,109}

This technique has been compared with the percutaneous BD, and when both techniques are performed by skilled interventional operators, the technical and clinical success rates are similar, with no difference in the complication rate.⁸³

Fig. 2 shows a case of EUS-guided BD after a failed ERCP in a malignant biliary stricture with signs of tumor invasion in the papilla. In this case extrahepatic access (transduodenal route) with transmural drainage (choledochoduodenostomy) was accomplished with a novel device designed for use with an echoendoscope. It creates an internal ostomy and delivers a biliary lumen-apposing metal stent at the same time (6-mm diameter, 8-mm saddle length, HotAXIOS, Xlumena Inc., Mountain View, CA, USA).

Table 6 EUS-guided biliary drainage in the literature.

Study	Design	Cases (n)	Technical success (%)	Clinical success (%)	Adverse events (%)
Bories et al. ⁸⁶	R	11	91	80	72
Maranki et al. ⁸⁷	R	49	84	80	18
Brauer et al. ⁸⁸	P	12	92	72	16
Horaguchi et al. ⁸⁹	P	16	100	94	37
Kim et al. ⁹⁰	R	15	80	80	None
Fabbri et al. ⁹¹	P	16	75	75	8
Park et al. ⁹²	R	57	96	89	47
Hara et al. ⁹³	P	18	94	94	77
Komaki et al. ⁹⁴	R	15	100	100	46
Ramírez-Luna et al. ⁹⁵	P	11	91	82	18
Shah et al. ⁹⁶	R	68	85	85	9
Iwashita et al. ⁹⁷	R	40	73	73	12
Dhir et al. ⁹⁸	R	58	98	98	3
Artifon et al. ⁸⁵	RCT	13	100	100	15
Song et al. ⁹⁹	P	15	87	87	47
Kim et al. ¹⁰⁰	P	13	92	84	38
Vila et al. ¹⁰¹	R	106	70	70	23
Horaguchi et al. ¹⁰²	R	21	100	100	10
Hara et al. ¹⁰³	P	18	94	89	27
Park et al. ¹⁰⁴	P	45	91	87	11
Kawakubo et al. ¹⁰⁵	R	14	100	100	14
Dhir et al. ¹⁰⁶	R	35	97	97	23
Khashab et al. ¹⁰⁷	R	35	94	91	14
Gornals et al. ⁷²	R	15	87	73	40
Gupta et al. ¹⁰⁸	R	240	99	87	35
Dhir et al. ¹⁰⁹	R	68	97	97	21
Kawakubo et al. ¹¹⁰	R	64	95	95	42
Total 27 studies	27	1088	91 (70–100)	87 (70–100)	29 (3–77)

Table modified from Fabbri et al.⁸⁴

RCT: randomized controlled trial; P: prospective study; R: retrospective study.

Pancreatic duct drainage guided by EUS

EUS-guided PD is indicated after failed ERCP, usually in patients with benign disorders such as stenosis, lithiasis, or post-interventional strictures. At present, there are more than 300 descriptions of this technique in the literature.

This technique is more challenging technically than EUS-guided BD. For this reason, general clinical and technical outcomes (78%) are poorer.^{83,84}

As with the previous technique, there is a nomenclature of the different variations. Briefly, there are 2 different approaches: transmural, usually transgastric (i.e., pancreaticogastrostomy), and transpapillary. Transpapillary guidewire placement allows retrograde access via rendezvous ERCP and antegrade stent placement for PD drainage.

The mean overall complication rate is 20% (7–55%). The transmural route is associated with a higher complication rate than the rendezvous route. This is clearly due to the fact that EUS-guided transmural PD stenting requires more aggressive dilation of the tract than does the rendezvous technique.⁸⁴

Fig. 3 illustrates EUS-guided PD drainage in a case of benign stenosis in chronic pancreatitis. The internal ostomy

is performed using a 6F cystotome, and a 7F transgastric plastic stent is delivered.

Miscellaneous

Endoscopic papillectomy for ampullary lesions

EUS is useful for the differential diagnosis of a prominent ampulla, with the majority of these lesions (>95%) being either adenomas or adenocarcinomas.

The universal incorporation of the EUS into the diagnostic evaluation of an ampullary adenoma is still controversial. If clinical suspicion of invasive carcinoma is low, and the lesion appears amenable to endoscopic resection, then EUS may not impact the decision to stage the lesion via papillectomy.¹¹¹

However, EUS is recommended to assess the endoscopic resectability of ampullary adenomas because it is the best technique for T staging of these lesions and it is highly accurate in evaluating intraductal extension and infiltration of the periampullary wall layers.^{112,113}

Although the accuracy of ERCP compared with EUS for delineating ductal extension of tumor requires further study,




Figure 2 EUS-guided biliary drainage. (A) Failed ERCP in a duodenal papilla with tumor invasion signs; (B) common bile duct (CBD) access by EUS-guided puncture; (C) cholangiography shows a dilated CBD with a malignant distal stenosis; (D) EUS-guided choledochoduodenostomy using a cautery-tipped stent delivery system (HotAXIOS, Xlumena Inc.); (E) EUS image of a lumen-apposing metal stent delivered inside the CBD; (F) biliary AXIOS stent (6 × 8 mm) delivered and well positioned in the duodenum. Dr. JB Gornals, H. Bellvitge, Barcelona.

the performance of a cholangiopancreatogram at the time of resection is helpful unless EUS has previously confirmed ductal involvement or irresectability.

In addition, ERCP has a crucial role in palliation of obstructive jaundice in the setting of ampullary adenocarcinoma and in minimizing the risk of post-papillectomy pancreatitis by placement of a pancreatic stent.¹¹⁴

Endoscopic drainage for pancreatic collections with disrupted pancreatic duct

Endoscopic transmural drainage of pancreatic collections (pseudocysts or walled-off pancreatic necrosis), with or without EUS-guidance, is safe, effective, has lower morbidity and mortality than surgery, and has been adopted as the first therapeutic option for drainage of pancreatic collections, when indicated, in many centers.¹¹⁵

ERCP allows the endoscopic transpapillary drainage by placement of an endoprosthesis through the papilla into the PD and is highly effective in the presence of a partial PD disruption that can be successfully bridged.

Data on the role of combined transmural and transpapillary drainage of pancreatic collections are limited and inconsistent. Based on available evidence, it seems reasonable to combine the two therapies in patients with pseudocysts and underlying chronic pancreatitis or known ductal abnormalities.^{116,117}

In the event of complete PD disruption, transpapillary drainage is hardly effective, and the placement of a permanent transmural stent is usually recommended.

The future of the relationship: discussion

With all the above in mind, it is easy to think that the relationship between EUS and ERCP will be close and strong, in the management of biliopancreatic disorders. The two techniques have experienced similar growth over the years, as new diagnostic procedures that have become interventional therapeutic techniques. Until recently, the two techniques were treated as different scopes or devices, and two different groups of endoscopists applied themselves to either ERCP or EUS.

However, in the last 5 or 10 years, EUS has undergone a transformation into a therapeutic technique, demanding skills in devices and stents from the ERCP. For this reason, some endoscopists dedicated to ERCP have taken up EUS, exploiting its therapeutic potential, which, to judge by the literature, has grown exponentially.

Following on from the era in which EUS and ERCP were in different worlds, some young endoscopists have been able to bridge this gap between the two advanced endoscopy techniques, and in so doing have created a single specialization.

The belief that the upcoming endoscopist should have to choose between EUS and ERCP is negative and detrimental to the progress of advanced endoscopy.¹¹⁸

The learning-curve of the two techniques will be demanding and challenging because a high level of skill is required for these pancreaticobiliary endoscopy procedures. Modern training must view the two techniques as one in order to facilitate training and motivate the learner.




Figure 3 EUS-guided pancreatic duct drainage after failed ERCP. (A) Pancreatic duct (PD) access by EUS-guided puncture using a 19G needle; (B) pancreatography and creation of a pancreaticogastrostomy using a cystotome over a guidewire; (C) plastic stent delivered inside the PD. (D) A 7Fr transgastric plastic stent, well positioned in the gastric cavity. Dr. JB Gornals, H. Bellvitge, Barcelona.

Conclusions

EUS and ERCP in good hands are advanced endoscopic procedures, and if they are used in combination, they offer a powerful positive effect on the management of biliopancreatic disorders. Nowadays, the relationship between them depends on the historic context of each center where they are used. But have no doubt that both techniques share a promising future together. More than friends, they are brothers.

Author contributions

Joan B. Gornals was involved in all the stages and in the conception and design of the article, and distributed each author with each section. J. M. Esteban, C. Guarnar-Argente, C. Marra-López, A. Repiso, O. Sendino, and C. Loras were involved with all stages of the manuscript development: analysis and interpretation of the data, drafting of the article, critical revision of the article, and final approval of the article.

Conflict of interest

Authors declare no conflict of interests for this article.

References

- Hollerbach S. EUS and ERCP: brothers in arms. *Gastrointest Endosc.* 2008;68:467–9, <http://dx.doi.org/10.1016/j.gie.2008.01.022>. PMID: 18760174.
- Gornals JB [PhD dissertation] Introduction of the interventional and therapeutic endoscopic ultrasound in an university hospital. Its combination with endoscopic retrograde cholangiopancreatography in biliopancreatic disorders: clinical and economical aspects. Barcelona: University of Barcelona, School of Medicine; 2013. <http://hdl.handle.net/10803/121465>
- Parada KS, Peng R, Erickson RA, Hawes R, Sahai AV, Ziogas A, et al. A resource utilization projection study of EUS. *Gastrointest Endosc.* 2002;55:328–34, <http://dx.doi.org/10.1067/mge.2002.118948>. PMID: 11868004.
- Èlsen GM, Dominitz JA, Faigel DO, Goldtein JA, Petersen GT, Radawi HM, et al. Guidelines for credentialing and grating privileges for endoscopic ultrasound. *Gastrointest Endosc.* 2001;54:811–4, [http://dx.doi.org/10.1016/s0016-5107\(01\)70082-X](http://dx.doi.org/10.1016/s0016-5107(01)70082-X). PMID: 11726873.
- Faigel DO. Economic realities of EUS in an academic practice. *Gastrointest Endosc.* 2007;65:287–9, <http://dx.doi.org/10.1016/j.gie.2006.06.045>. PMID: 17258989.
- Wani S, Coté GA, Keswani R, Mallady D, Azar R, Murad F, et al. Learning curves for EUS by using cumulative sum analysis: implications for American Society for Gastrointestinal Endoscopy recommendations for training. *Gastrointest Endosc.*

- 2013;77:558–65, <http://dx.doi.org/10.1016/j.gie.2012.10.012>. PMID: 23260317.
7. Early DS, Hall M, Aslanian H, Casey B, Burbridge R, Chak A, et al. A Prospective, Multicenter Study Research the Aptitude of Trainees in Endoscopic Ultrasonography (Rate US Study) Using Cumulative sum Analysis (CUSUM). *Gastrointest Endosc.* 2014;79:AB137–8, <http://dx.doi.org/10.1016/j.gie.2014.02.093>
 8. Mertz H, Gautam S. The learning curve for EUS-guided FNA of pancreatic cancer. *Gastrointest Endosc.* 2004;59:33–7, [http://dx.doi.org/10.1016/S0016-5107\(03\)02028-5](http://dx.doi.org/10.1016/S0016-5107(03)02028-5). PMID: 14722544.
 9. Eloubeidi MA, Tamhane A. EUS-guided FNA of solid pancreatic masses: a learning curve with 300 consecutive procedures. *Gastrointest Endosc.* 2005;61:700–8, [http://dx.doi.org/10.1016/S0016-5107\(05\)00363-9](http://dx.doi.org/10.1016/S0016-5107(05)00363-9). PMID: 15855975.
 10. Paquin SC. Training in endoscopic ultrasound-guided fine needle aspiration. *Endosc Ultrasound.* 2014;3:12–6, <http://dx.doi.org/10.4103/2303-9027.127123>. PMID: 24949405.
 11. Iglesias-Garcia J, Lariño-Noia J, Dominguez-Muñoz JE. When to puncture, when not to puncture: pancreatic masses. *Endosc Ultrasound.* 2014;3:91–7, <http://dx.doi.org/10.4103/2303-9027.123007>. PMID: 2495338.
 12. Polkowski M, Larghi A, Weynand B, Boustière C, Giovannini M, Pujol B, et al. Endoscopic ultrasound (EUS)-guided sampling in gastroenterology: European Society of Gastrointestinal Endoscopy (ESGE) Technical Guideline. *Endoscopy.* 2012;44:190–205, <http://dx.doi.org/10.1055/s-0031-1291543>. PMID: 22180307.
 13. Jacobson BC, Chak A, Hoffman B, Baron TH, Cohen J, Deal SE, et al. Quality indicators for endoscopic ultrasonography. *Gastrointest Endosc.* 2006;63:535–8, <http://dx.doi.org/10.1016/j.gie.2006.02.020>. PMID: 16564910.
 14. Bhutani MS. Interventional endoscopic ultrasonography: state of the art at the new millenium. *Endoscopy.* 2000;32:62–71, <http://dx.doi.org/10.1055/s-2000-139>. PMID: 10691275.
 15. Jowell PS, Baillie J, Branch MS, Affronti J, Browning CL, Bute BP. Quantitative assessment of procedural competence: a prospective study of training in endoscopic retrograde cholangiopancreatography. *Ann Intern Med.* 1996;125:983–9, <http://dx.doi.org/10.7326/0003-4819-125-12-199612150-00009>. PMID: 8967710.
 16. Adler DG, Lieb JG, Cohen J, Pike IM, Park WG, Rizk MK, et al. Quality indicators for ERCP. *Gastrointest Endosc.* 2015;81:54–66, <http://dx.doi.org/10.1016/j.gie.2014.07.056>. PMID: 25480099.
 17. Chutkan RK, Ahmad AS, Cohen J, Cruz-Correa MR, Desilets DJ, Dominitz JA, et al. ERCP core curriculum. *Gastrointest Endosc.* 2006;63:361–76. PMID: 16500380.
 18. Ekkelenkamp VE, Koch AD, Rauws EA, Borsboom GJ, de Man RA, Kuipers EJ. Competence development in ERCP: the learning curve of novice trainees. *Endoscopy.* 2014;46:949–55, <http://dx.doi.org/10.1055/s-0034-1377930>. PMID: 25208031.
 19. Adler DG, Baron TH, Davila RE, Egan J, Hirota WK, Leighton JA, et al. Standards of Practice Committee of American Society for Gastrointestinal Endoscopy. ASGE guideline: the role of ERCP in diseases of the biliary tract and the pancreas. *Gastrointest Endosc.* 2005;62:1–8. PMID: 15990812.
 20. Early DS, Ben-Menachem T, Decker GA, Evans JA, Fanelli RD, Fisher DA, et al. Appropriate use of GI endoscopy. *Gastrointest Endosc.* 2012;75:1127–31, <http://dx.doi.org/10.1016/j.gie.2012.01.011>. PMID: 22624807.
 21. ASGE Standards of Practice Committee, Chathadi KV, Chandrasekhara V, Acosta RD, Decker GA, Early DS, et al. The role of ERCP in benign diseases of the biliary tract. *Gastrointest Endosc.* 2015, <http://dx.doi.org/10.1016/j.gie.2014.11.019>. pii:S0016-5107(14)02433-X, PMID: 2566593.
 22. Anderson MA, Appalaneni V, Ben-Menachem T, Decker GA, Early DS, Evans JA, et al. The role of endoscopy in the evaluation and treatment of patients with biliary neoplasia. *Gastrointest Endosc.* 2013;77:167–74, <http://dx.doi.org/10.1016/j.gie.2012.09.029>. PMID: 23219047.
 23. Christodoulou DK, Tsianos EV. Role of endoscopic retrograde cholangiopancreatography in pancreatic diseases. *World J Gastroenterol.* 2010;16:4755–61. PMID: 20939103.
 24. Buxbaum J. The role of endoscopic retrograde cholangiopancreatography in patients with pancreatic disease. *Gastroenterol Clin N Am.* 2012;41:23–45, <http://dx.doi.org/10.1016/j.gtc.2011.12.010>. PMID: 22341248.
 25. Shaffer EA. Epidemiology of gallbladder stone disease. *Best Pract Res Clin Gastroenterol.* 2006;20:98, <http://dx.doi.org/10.1016/j.bpg.2006.05.004>. PMID: 17127183.
 26. Collins C, Maguire D, Ireland A, Fitzgerald E, O'Sullivan GC. A prospective study of common bile duct calculi in patients undergoing laparoscopic cholecystectomy: natural history of choledocholithiasis revisited. *Ann Surg.* 2004;239:28–33, <http://dx.doi.org/10.1097/01.sla.0000103069.00170.9c>. PMID: 14685097.
 27. National institutes of health consensus development conference statement on gallstones and laparoscopic cholecystectomy. *Am J Surg.* 1993;165:390–8, [http://dx.doi.org/10.1016/S0002-9610\(05\)80929-8](http://dx.doi.org/10.1016/S0002-9610(05)80929-8). PMID: 8480870.
 28. Williams EJ, Green J, Beckingham I, Parks R, Martin D, Lombard M. Guidelines on the management of common bile duct stones (CBDs). *Gut.* 2008;57:1004–21, <http://dx.doi.org/10.1136/gut.2007.121657>. PMID: 18321943.
 29. Maple JT, Ikenberry SO, Anderson MA, Appalaneni V, Decker GA, Early D, et al. The role of endoscopy in the management of choledocholithiasis. *Gastrointest Endosc.* 2011;74:731–43, <http://dx.doi.org/10.1016/j.gie.2011.04.012>. PMID: 21951472.
 30. Tse F, Liu L, Barkun AN, Armstrong D, Moayyedi P. Nonoperative imaging techniques in suspected biliary tract obstruction. *Gastrointest Endosc.* 2008;67:235–44, <http://dx.doi.org/10.1016/j.gie.2007.09.047>. PMID: 18226685.
 31. Prat F, Amouyal G, Amouyal P, Pelletier G, Fritsch J, Choury AD, et al. Prospective controlled study of endoscopic ultrasonography and endoscopic retrograde cholangiography in patients with suspected common bile duct lithiasis. *Lancet.* 1996;347:75–9, [http://dx.doi.org/10.1016/S0140-6736\(96\)90208-1](http://dx.doi.org/10.1016/S0140-6736(96)90208-1). PMID: 8538344.
 32. Tseng LJ, Jao YT, Mo LR, Lin RC. Over-the-wire US catheter probe asan adjunct to ERCP in the detection of choledocholithiasis. *Gastrointest Endosc.* 2001;54:720–3, <http://dx.doi.org/10.1067/mge.2001.119255>. PMID: 11726847.
 33. Larghi A, Petrone MC, Galasso D, Arcidiacono PG. Endoscopic ultrasound in the evaluation of pancreaticobiliary disorders. *Dig Liver Dis.* 2009;42:6–15, <http://dx.doi.org/10.1016/j.dld.2009.06.021>. PMID: 19665951.
 34. Garrow D, Miller S, Sinha D, Conway J, Hoffman BJ, Hawes RH, et al. Endoscopic ultrasound: a metaanalysis of test performance in suspected biliary obstruction. *Clin Gastroenterol Hepatol.* 2007;5:616–23, <http://dx.doi.org/10.1016/j.cgh.2007.02.027>. PMID: 17478348.
 35. Petrov MS, Savides TJ. Systematic review of endoscopic ultrasonography versus endoscopic retrograde cholangiopancreatography for suspected choledocholithiasis. *Br J Surg.* 2009;96:967–74, <http://dx.doi.org/10.1002/bjs.6667>. PMID: 19644975.
 36. Liu CL, Fan ST, Lo CM, Tso WK, Wong Y, Poon RT, et al. Comparison of early endoscopic ultrasonography and endoscopic retrograde cholangiopancreatography in the

- management of acute biliary pancreatitis: a prospective randomized study. *Clin Gastroenterol Hepatol.* 2005;3:1238–44, [http://dx.doi.org/10.1016/s1542-3565\(05\)00619-1](http://dx.doi.org/10.1016/s1542-3565(05)00619-1). PMID: 16361050.
37. Lee YT, Chan FK, Leung WK, Chan HL, Wu JC, Yung MY, et al. Comparison of EUS and ERCP in the investigation with suspected biliary obstruction caused by choledocholithiasis: a randomized study. *Gastrointest Endosc.* 2008;67:660–8, <http://dx.doi.org/10.1016/j.gie.2007.07.025>. PMID: 18155205.
 38. Polkowski M, Regula J, Tilszer A, Butruk E. Endoscopic ultrasound versus endoscopic retrograde cholangiography for patients with intermediate probability of bile duct stones: a randomized trial comparing two management strategies. *Endoscopy.* 2007;39:296–303, <http://dx.doi.org/10.1055/s-2007-966264>. PMID: 17427065.
 39. Karakan T, Cindoruk M, Alagozlu H, Ergun M, Dumlu S, Unal S. EUS versus endoscopic retrograde cholangiography for patients with intermediate probability of bile duct stones: a prospective randomized trial. *Gastrointest Endosc.* 2009;69:244–52, <http://dx.doi.org/10.1016/j.gie.2008.05.023>. PMID: 19019364.
 40. Amouyal P, Amouyal G, Lévy P, Tuzet S, Palazzo L, Vilgrain V, et al. Diagnosis of choledocholithiasis by endoscopic ultrasonography. *Gastroenterology.* 1994;106:1062–7, <http://dx.doi.org/10.1002/hep.1840200445>. PMID: 8143973.
 41. Sugiyama M, Atomi Y. Endoscopic ultrasonography for diagnosing choledocholithiasis: a prospective comparative study with ultrasonography and computed tomography. *Gastrointest Endosc.* 1997;45:143–6, [http://dx.doi.org/10.1016/s0016-5107\(97\)70237-2](http://dx.doi.org/10.1016/s0016-5107(97)70237-2). PMID: 9040999.
 42. Yoo WJ, Brugge ER. Endoscopic evaluation of bile duct strictures. *Gastrointest Endosc Clin N Am.* 2013;23:277–93, <http://dx.doi.org/10.1016/j.giec.2012.12.002>. PMID: 23540961.
 43. Foutch PG. Diagnosis of cancer by cytologic methods performed during ERCP. *Gastrointest Endosc.* 1994;40:249–52. PMID: 8013838.
 44. Weber A, von Weyhern C, Fend F, Schneider J, Neu B, Meining A, et al. Endoscopic transpapillary brush cytology and forceps biopsy in patients with hilar cholangiocarcinoma. *World J Gastroenterol.* 2008;14:1097–101. PMID: 18286693.
 45. Lee JH, Salem R, Aslanian H, Chacho M, Topazian M. Endoscopic ultrasound-guided fine-needle aspiration of unexplained bile duct strictures. *Am J Gastroenterol.* 2004;99:1069–73. PMID: 15180727.
 46. Eloubeidi MA, Chen VK, Jhala NC, Eltoun IE, Jhala D, Chhieng DC, et al. Endoscopic ultrasound-guided fine-needle aspiration biopsy of suspected cholangiocarcinoma. *Clin Gastroenterol Hepatol.* 2004;2:209–13. PMID: 15017604.
 47. Fritscher-Ravens A, Broering DC, Sriram PV, Topalidis T, Jaekle S, Thonke F, et al. EUS-guided fine-needle aspiration cytodiagnosis of hilar cholangiocarcinoma: a case series. *Gastrointest Endosc.* 2000;52:534–40. PMID: 11023576.
 48. Byrne MF, Gerke H, Mitchell RM, Stiffler HL, McGrath K, Branch MS, et al. Yield of endoscopic ultrasound-guided fine-needle aspiration of bile duct lesions. *Endoscopy.* 2004;36:715–9. PMID: 15280978.
 49. DeWitt J, Misra VL, Leblanc JK, McHenry L, Sherman S. EUS-guided FNA of proximal biliary strictures after negative ERCP brush cytology results. *Gastrointest Endosc.* 2006;64:325–33. PMID: 16923477.
 50. Fritscher-Ravens A, Broering DC, Knoefel WT, Rogiers X, Swain P, Thonke F, et al. EUS-guided fine-needle aspiration of suspected hilar cholangiocarcinoma in potentially operable patients with negative brush cytology. *Am J Gastroenterol.* 2004;99:45–51. PMID: 14687140.
 51. Ohshima Y, Yasuda I, Kawakami H, Kuwatani M, Mukai T, Iwashita T, et al. EUS-FNA for suspected malignant biliary strictures after negative endoscopic transpapillary brush cytology and forceps biopsy. *J Gastroenterol.* 2011;46:921–8, <http://dx.doi.org/10.1007/s00535-011-0404-z>. PMID: 21526370.
 52. Rösch T, Hofrichter K, Frimberger E, Meining A, Born P, Weigert N, et al. ERCP or EUS for tissue diagnosis of biliary strictures? A prospective comparative study. *Gastrointest Endosc.* 2004;60:390–6. PMID: 15332029.
 53. Oppong K, Raine D, Nayar M, Nayar M, Wadehra V, Ramakrishnan S, et al. EUS-FNA versus biliary brushings and assessment of simultaneous performance in jaundiced patients with suspected malignant obstruction. *JOP.* 2010;11:560–7. PMID: 21068487.
 54. Weilert F, Bhat YM, Binmoeller KF, Kane S, Jaffee IM, Shaw RE, et al. EUS-FNA is superior to ERCP-based tissue sampling in suspected malignant biliary obstruction: results of a prospective, single-blind, comparative study. *Gastrointest Endosc.* 2014;80:97–104, <http://dx.doi.org/10.1016/j.gie.2013.12.031>. PMID: 24559784.
 55. Khashab M, Fockens P, Al-Haddad MA. Utility of EUS in patients with indeterminate biliary stricture and suspected extrahepatic cholangiocarcinoma. *Gastrointest Endosc.* 2012;76:1024–33.
 56. Chafic AH, DeWitt J, Leblanc JK, et al. Impact of preoperative endoscopic ultrasound-guided fine needle aspiration on preoperative recurrence and survival in cholangiocarcinoma patients. *Endoscopy.* 2013;45:883–9.
 57. Peter S, Eloubeidi MA. Feasibility of combined EUS-FNA and ERCP for obstructive jaundice from presumed pancreatic malignancy. *Nat Clin Pract Gastroenterol Hepatol.* 2009;6:132–3, <http://dx.doi.org/10.1038/ncpgasthep1365>. PMID: 19204740.
 58. Hewitt MJ, McPhail MJ, Possamai L, Dhar A, Vlavianos P, Monahan KJ. EUS-guided FNA for diagnosis of solid pancreatic neoplasms: a meta-analysis. *Gastrointest Endosc.* 2012;75:319–31, <http://dx.doi.org/10.1016/j.gie.2011.08.049>. PMID: 22248600.
 59. Fusaroli P, Kypraios D, Caletti G, Eloubeidi MA. Pancreaticobiliary endoscopic ultrasound: a systematic review of the levels of evidence, performance and outcomes. *World J Gastroenterol.* 2012;18:4243–56, <http://dx.doi.org/10.3748/wjg.v18.i32.4243>. PMID: 22969187.
 60. Moss AC, Morris E, Mac Mathuna P. Palliative biliary stents for obstructing pancreatic carcinoma. *Cochrane Database Syst Rev.* 2006;19:CD004200. PMID: 16625598.
 61. Standards of Practice Committee, Maple JT, Ikenberry SO, Anderson MA, Appalaneni V, Decker GA, et al. The role of endoscopy in the management of choledocholithiasis. *Gastrointest Endosc.* 2011;74:731–44, <http://dx.doi.org/10.1016/j.gie.2011.04.012>. PMID: 21951472 [Erratum in: *Gastrointest Endosc.* 2012; 75:230–230.e14].
 62. Wiersema MJ, Sandusky D, Carr R, Wiersema LM, Erdel WC, Frederick PK. Endoscopy-guided cholangiopancreatography. *Gastrointest Endosc.* 1996;43:102–6. PMID: 8635700.
 63. Rocca R, De Angelis C, Castellino F, Masoero G, Daperno M, Sostegni R, et al. EUS diagnosis and simultaneous endoscopic retrograde cholangiography treatment of common bile duct stones by using an oblique-viewing echoendoscope. *Gastrointest Endosc.* 2006;63:479–84. PMID: 16500400.
 64. Tarantino I, Barresi L, Di Pisa M, Traina M. Simultaneous endoscopic ultrasound fine needle aspiration and endoscopic retrograde cholangio-pancreatography: evaluation of safety. *World J Gastroenterol.* 2007;13:3861–3. PMID: 17657842.
 65. Ross WA, Wasan SM, Evans DB, Wolff RA, Trapani LV, Staerckel GA, et al. Combined EUS with FNA and ERCP for the evaluation of patients with obstructive jaundice

- from presumed pancreatic malignancy. *Gastrointest Endosc.* 2008;68:461–6, <http://dx.doi.org/10.1016/j.gie.2007.11.033>. PMID: 18384788.
66. Fabbri C, Polifemo AM, Luigiano C, Cennamo V, Fuccio L, Billi P, et al. Single session versus separate session endoscopic ultrasonography plus endoscopic retrograde cholangiography in patients with low to moderate risk for choledocholithiasis. *J Gastroenterol Hepatol.* 2009;24:1107–12, <http://dx.doi.org/10.1111/j.1440-1746.2009.05828.x>. PMID: 19638088.
 67. Ascunze G, Ribeiro A, Rocha-Lima C, Larsen M, Sleeman D, Merchan J, et al. Single-session endoscopic ultrasonography and endoscopic retrograde cholangiopancreatography for evaluation of pancreaticobiliary disorders. *Surg Endosc.* 2010;24:1447–50, <http://dx.doi.org/10.1007/s00464-009-0798-3>. PMID: 20054580.
 68. Vila JJ, Kutz M, Goñi S, Ostiz M, Amorena E, Prieto C, et al. Endoscopic and anesthetic feasibility of EUS and ERCP combined in a single session versus two different sessions. *World J Gastrointest Endosc.* 2011;3:57–61, <http://dx.doi.org/10.4253/wjge.v3.i3.57>. PMID: 21455343.
 69. Aslanian HR, Estrada JD, Rossi F, Dziura J, Jamidar PA, Siddiqui UD. Endoscopic ultrasound and endoscopic retrograde cholangiopancreatography for obstructing pancreas head masses: combined or separate procedures? *J Clin Gastroenterol.* 2011;45:711–3, <http://dx.doi.org/10.1097/MCG.0b013e3182045923>. PMID: 21301359.
 70. İles-Shih L, Hilden K, Adler DG. Combined ERCP and EUS in one session is safe in elderly patients when compared to non-elderly patients: outcomes in 206 combined procedures. *Dig Dis Sci.* 2012;57:1949–53, <http://dx.doi.org/10.1007/s10620-012-2135-2>. PMID: 22453997.
 71. Benjaminov F, Stein A, Lichtman G, Pomeranz I, Konikoff FM. Consecutive versus separate sessions of endoscopic ultrasound (EUS) and endoscopic retrograde cholangiopancreatography (ERCP) for symptomatic choledocholithiasis. *Surg Endosc.* 2013;27:2117, <http://dx.doi.org/10.1007/s00464-012-2720-7>. PMID: 23389062.
 72. Gornals JB, Moreno R, Castellote J, Loras C, Barranco R, Catala I, et al. Single-session endosonography and endoscopic retrograde cholangiopancreatography for biliopancreatic diseases is feasible, effective and cost beneficial. *Dig Liver Dis.* 2013;45:578–83, <http://dx.doi.org/10.1016/j.dld.2013.01.023>. PMID: 23465682.
 73. Chu YL, Wang XF, Gao XZ, Qiao XL, Liu F, Yu SY, et al. Endoscopic ultrasonography in tandem with endoscopic retrograde cholangiopancreatography in the management of suspected distal obstructive jaundice. *Eur J Gastroenterol Hepatol.* 2013;25:455–9, <http://dx.doi.org/10.1097/MEG.0b013e32835ca1d7>. PMID: 23249605.
 74. Kawakubo K, Kawakami H, Kuwatani M, Haba S, Kudo T, Abe Y, et al. Safety and utility of single-session endoscopic ultrasonography and endoscopic retrograde cholangiopancreatography for the evaluation of pancreatobiliary diseases. *Gut Liver.* 2014;8:329–32, <http://dx.doi.org/10.5009/gnl.2014.8.3.329>. PMID: 248276632.
 75. Noma Y, Kawamoto H, Kato H, Iwamura M, Hirao K, Fujii M, et al. The efficacy and safety of single-session endoscopic ultrasound-guided fine needle aspiration and endoscopic retrograde cholangiopancreatography for evaluation of pancreatic masses. *Hepatogastroenterology.* 2014;61:1775–9, <http://dx.doi.org/10.1097/MCG.0b013e3182045923>. PMID: 21301359.
 76. Vohra S, Holt EW, Bhat YM, Kane S, Shah JN, Binmoeller KF. Successful single-session endosonography-based endoscopic retrograde cholangiopancreatography without fluoroscopy in pregnant patients with suspected choledocholithiasis: a case series. *J Hepatobiliary Pancreat Surg.* 2014;21:93–7, <http://dx.doi.org/10.1002/jhbp.7>. PMID: 23798477.
 77. Verma D, Kapadia A, Eisen GM, Adler DG. EUS vs MRCP for detection of choledocholithiasis. *Gastrointest Endosc.* 2006;64:248–54. PMID: 16860077.
 78. Gonzalo-Marin J, Vila JJ, Perez-Miranda M. Role of endoscopic ultrasound in the diagnosis of pancreatic cancer. *World J Gastrointest Oncol.* 2014;6:360–8, <http://dx.doi.org/10.4251/wjgo.v6.i9.360>. PMID: 25232461.
 79. Fusaroli P, Manta R, Fedeli P, Maltoni S, Grillo A, Giovannini E, et al. The influence of endoscopic biliary stents on the accuracy of endoscopic ultrasound for pancreatic head cancer staging. *Endoscopy.* 2007;39:813–7. PMID: 17703391.
 80. Shami VM, Mahajan A, Sundaram V, Davis EM, Loch MM, Kahaleh M. Endoscopic ultrasound staging is adversely affected by placement of a self-expandable metal stent: fact or fiction? *Pancreas.* 2008;37:396–8, <http://dx.doi.org/10.1097/MPA.0b013e3181800d2e>. PMID: 18953251.
 81. Mergener K, Jowell PS, Branch MS, Baillie J. Pneumoperitoneum complicating ERCP performed immediately after EUS-guided fine needle aspiration. *Gastrointest Endosc.* 1998;47:541–2. PMID: 9647385.
 82. Di Matteo F, Shimpi L, Gabbriellini A, Martino M, Caricato M, Esposito A, et al. Same-day endoscopic retrograde cholangiopancreatography after transduodenal endoscopic ultrasound-guided needle aspiration: do we need to be cautious? *Endoscopy.* 2006;38:1149–51. PMID: 17111340.
 83. Kahaleh M, Artifon ELA, Perez-Miranda M, Gaidhane M, Rondon C, Itoi T, et al. Endoscopic ultrasonography guided drainage: summary of consortium meeting, May 21, 2012, San Diego, California. *World J Gastroenterol.* 2015;21:726–41, <http://dx.doi.org/10.3748/wjg.v21.i3.726>. PMID: 25624708.
 84. Fabbri C, Luigiano C, Lisotti A, Cennamo V, Virgilio C, Caletti G, et al. Endoscopic ultrasound-guided treatments: are we getting evidence based – a systematic review. *World J Gastroenterol.* 2014;20:8424–48, <http://dx.doi.org/10.3748/wjg.v20.i26.8424>. PMID: 25024600.
 85. Artifon EL, Aparicio D, Paione JB, Lo SK, Bordini A, Rabello C, et al. Biliary drainage in patients with unresectable, malignant obstruction where ERCP fails: endoscopic ultrasonography-guided choledochoduodenostomy versus percutaneous drainage. *J Clin Gastroenterol.* 2012;46:768–74, <http://dx.doi.org/10.1097/MCG.0b013e31825f264c>. PMID: 22810111.
 86. Bories E, Pesenti C, Caillol F, Lopes C, Giovannini M. Transgastric endoscopic ultrasonography-guided biliary drainage: results of a pilot study. *Endoscopy.* 2007;39:287–91, <http://dx.doi.org/10.1055/s-2007-966212>. PMID: 17357952.
 87. Maranki J, Hernandez AJ, Arslan B, Jaffan AA, Angle JF, Shami VM, et al. Interventional endoscopic ultrasound-guided cholangiography: long-term experience of an emerging alternative to percutaneous transhepatic cholangiography. *Endoscopy.* 2009;41:532–8, <http://dx.doi.org/10.1055/s-0029-1214712>. PMID: 19533558.
 88. Brauer BC, Chen YK, Fukami N, Shah RJ. Single-operator EUS-guided cholangiopancreatography for difficult pancreaticobiliary access (with video). *Gastrointest Endosc.* 2009;70:471–9, <http://dx.doi.org/10.1016/j.gie.2008.12.233>. PMID: 19560768.
 89. Horaguchi J, Fujita N, Noda Y, Kobayashi G, Ito K, Obana T, et al. Endosonography-guided biliary drainage in cases with difficult transpapillary endoscopic biliary drainage. *Dig Endosc.* 2009;21:239–44, <http://dx.doi.org/10.1111/j.1443-1661.2009.00899.x>. PMID: 19961522.
 90. Kim YS, Gupta K, Mallery S, Li R, Kinney T, Freeman ML. Endoscopic ultrasound rendezvous for bile duct access using a transduodenal approach: cumulative experience at

- a single center. A case series. *Endoscopy*. 2010;42:496–502, <http://dx.doi.org/10.1055/s-0029-1244082>. PMID: 20419625.
91. Fabbri C, Luigiano C, Fuccio L, Polifemo AM, Ferrara F, Ghersi S, et al. EUS-guided biliary drainage with placement of a new partially covered biliary stent for palliation of malignant biliary obstruction: a case series. *Endoscopy*. 2011;43:438–41, <http://dx.doi.org/10.1055/s-0030-1256097>. PMID: 21271507.
 92. Park do H, Jang JW, Lee SS, Seo DW, Lee SK, Kim MH. EUS guided biliary drainage with transluminal stenting after failed ERCP: predictors of adverse events and long-term results. *Gastrointest Endosc*. 2011;74:1276–84, <http://dx.doi.org/10.1016/j.gie.2011.07.054>. PMID: 21963067.
 93. Hara K, Yamao K, Niwa Y, Sawaki A, Mizuno N, Hijioka S, et al. Prospective clinical study of EUS-guided choledochoduodenostomy for malignant lower biliary tract obstruction. *Am J Gastroenterol*. 2011;106:1239–45, <http://dx.doi.org/10.1038/ajg.2011.84>. PMID: 21448148.
 94. Komaki T, Kitano M, Sakamoto H, Kudo M. Endoscopic ultrasonography-guided biliary drainage: evaluation of a choledochoduodenostomy technique. *Pancreatol*. 2011;11 Suppl. 2:47–51, <http://dx.doi.org/10.1159/000323508>. PMID: 21464587.
 95. Ramírez-Luna MA, Téllez-Ávila FI, Giovannini M, Valdovinos-Andraca F, Guerrero-Hernández I, Herrera-Esquivel J. Endoscopic ultrasound-guided biliodigestive drainage is a good alternative in patients with unresectable cancer. *Endoscopy*. 2011;43:826–30, <http://dx.doi.org/10.1055/s-0030-1256406>. PMID: 21833899.
 96. Shah JN, Marson F, Weilert F, Bhat YM, Nguyen-Tang T, Shaw RE, et al. Single-operator, single-session EUS-guided anterograde cholangiopancreatography in failed ERCP or inaccessible papilla. *Gastrointest Endosc*. 2012;75:56–64, <http://dx.doi.org/10.1016/j.gie.2011.08.032>. PMID: 22018554.
 97. Iwashita T, Lee JG, Shinoura S, Nakai Y, Park DH, Muthusamy VR, et al. Endoscopic ultrasound-guided rendezvous for biliary access after failed cannulation. *Endoscopy*. 2012;44:60–5, <http://dx.doi.org/10.1055/s-0030-1256871>. PMID: 22127960.
 98. Dhir V, Bhandari S, Bapat M, Maydeo A. Comparison of EUS-guided rendezvous and precut papillotomy techniques for biliary access (with videos). *Gastrointest Endosc*. 2012;75:354–9, <http://dx.doi.org/10.1016/j.gie.2011.07.075>. PMID: 22248603.
 99. Song TJ, Hyun YS, Lee SS, Park do H, Seo DW, Lee SK, et al. Endoscopic ultrasound-guided choledochoduodenostomies with fully covered self-expandable metallic stents. *World J Gastroenterol*. 2012;18:4435–40, <http://dx.doi.org/10.3748/wjg.v18.i32.4435>. PMID: 22969210.
 100. Kim TH, Kim SH, Oh HJ, Sohn YW, Lee SO. Endoscopic ultrasound-guided biliary drainage with placement of a fully covered metal stent for malignant biliary obstruction. *World J Gastroenterol*. 2012;18:2526–32, <http://dx.doi.org/10.3748/wjg.v18.i20.2526>. PMID: 22654450.
 101. Vila JJ, Pérez-Miranda M, Vazquez-Sequeiros E, Abadia MA, Pérez-Millán A, González-Huix F, et al. Initial experience with EUS-guided cholangiopancreatography for biliary and pancreatic duct drainage: a Spanish national survey. *Gastrointest Endosc*. 2012;76:1133–41, <http://dx.doi.org/10.1016/j.gie.2012.08.001>. PMID: 23021167.
 102. Horaguchi J, Fujita N, Noda Y, Kobayashi G, Ito K, Koshita S, et al. Metallic stent deployment in endosonography-guided biliary drainage: long-term follow-up results in patients with bilio-enteric anastomosis. *Dig Endosc*. 2012;24:457–61, <http://dx.doi.org/10.1111/j.1443-1661.2012.01316.x>. PMID: 23078440.
 103. Hara K, Yamao K, Hijioka S, Mizuno N, Imaoka H, Tajika M, et al. Prospective clinical study of endoscopic ultrasound-guided choledochoduodenostomy with direct metallic stent placement using a forward-viewing echoendoscope. *Endoscopy*. 2013;45:392–6, <http://dx.doi.org/10.1055/s-0032-1326076>. PMID: 23338620.
 104. Park do H, Jeong SU, Lee BU, Lee SS, Seo DW, Lee SK, et al. Prospective evaluation of a treatment algorithm with enhanced guidewire manipulation protocol for EUS-guided biliary drainage after failed ERCP (with video). *Gastrointest Endosc*. 2013;78:91–101, <http://dx.doi.org/10.1016/j.gie.2013.01.042>. PMID: 23523301.
 105. Kawakubo K, Isayama H, Sasahira N, Nakai Y, Kogure H, Hamada T, et al. Clinical utility of an endoscopic ultrasound-guided rendezvous technique via various approach routes. *Surg Endosc*. 2013;27:3437–43, <http://dx.doi.org/10.1007/s00464-013-2896-5>. PMID: 23508814.
 106. Dhir V, Bhandari S, Bapat M, Joshi N, Vivekanandarajah S, Maydeo A. Comparison of transhepatic and extrahepatic routes for EUS-guided rendezvous procedure for distal CBD obstruction. *United Eur Gastroenterol J*. 2013;1:103–8, <http://dx.doi.org/10.1177/2050640613480145>. PMID: 24917947.
 107. Khashab MA, Valeshabad AK, Modayil R, Widmer J, Saxena P, Idrees M, et al. EUS-guided biliary drainage by using a standardized approach for malignant biliary obstruction: rendezvous versus direct transluminal techniques (with videos). *Gastrointest Endosc*. 2013;78:734–41, <http://dx.doi.org/10.1016/j.gie.2013.05.013>. PMID: 23886353.
 108. Gupta K, Perez-Miranda M, Kahaleh M, Artifon EL, Itoi T, Freeman ML, et al. Endoscopic ultrasound-assisted bile duct access and drainage: multicenter, long-term analysis of approach, outcomes, and complications of a technique in evolution. *J Clin Gastroenterol*. 2014;48:80–7, <http://dx.doi.org/10.1097/MCG.0b013e31828c6822>. PMID: 23632351.
 109. Dhir V, Artifon EL, Gupta K, Vila JJ, Maselli R, Frazao M, et al. Multicenter study on endoscopic ultrasound-guided expandable biliary metal stent placement: choice of access route, direction of stent insertion, and drainage route. *Dig Endosc*. 2014;26:430–5, <http://dx.doi.org/10.1111/den.12153>. PMID: 23941261.
 110. Kawakubo K, Isayama H, Kato H, Itoi T, Kawakami H, Hanada K, et al. Multicenter retrospective study of endoscopic ultrasound-guided biliary drainage for malignant biliary obstruction in Japan. *J Hepatobiliary Pancreat Surg*. 2014;21:328–34, <http://dx.doi.org/10.1002/jhbp.27>. PMID: 24026963.
 111. Patel R, Varadarajulu S, Wilcox CM. Endoscopic ampullectomy: techniques and outcomes. *J Clin Gastroenterol*. 2012;46:8–15, <http://dx.doi.org/10.1097/MCG.0b013e318233a844>. PMID: 22064552.
 112. Manta R, Conigliaro R, Castellani D, Messerotti A, Bertani H, Sabatino G, et al. Linear endoscopic ultrasonography vs magnetic resonance imaging in ampullary tumors. *World J Gastroenterol*. 2010;16:5592–7. PMID: 21105192.
 113. Chen CH, Yang CC, Yeh YH, Chou DA, Nien CK. Reappraisal of endosonography of ampullary tumors: correlation with transabdominal sonography, CT, and MRI. *J Clin Ultrasound*. 2009;37:18–25, <http://dx.doi.org/10.1002/jcu.20523>. PMID: 18726967.
 114. Dumonceau JM, Rigaux J, Kahaleh M, Gomez CM, Vandermeeren A, Devière J. Prophylaxis of post-ERCP pancreatitis: a practice survey. *Gastrointest Endosc*. 2010;71:934–9, <http://dx.doi.org/10.1016/j.gie.2009.10.055>. PMID: 20226455.
 115. Singhal S, Rotman SR, Gaidhane M, Kahaleh M. Pancreatic fluid collection drainage by endoscopic ultrasound: an update. *Clin Endosc*. 2013;46:506–14, <http://dx.doi.org/10.5946/ce.2013.46.5.506>. PMID: 24143313.
 116. Shrode CW, Macdonough P, Gaidhane M, Northrup PG, Sauer B, Ku J, et al. Multimodality endoscopic treatment of pancreatic duct disruption with stenting and pseudocyst drainage: how

- efficacious is it? *Dig Liver Dis.* 2013;45:129–33, <http://dx.doi.org/10.1016/j.dld.2012.08.026>. PMID: 23036185.
117. Trevino JM, Tamhane A, Varadarajulu S. Successful stenting in ductal disruption favorably impacts treatment outcomes in patients undergoing transmural drainage of peripancreatic fluid collections. *J Gastroenterol Hepatol.* 2010;25:526–31, <http://dx.doi.org/10.1111/j.1440-1746.2009.06109.x>. PMID: 20074158.
118. Mangiavillano B, Baron TH. Changing our perspective in endoscopic ultrasound (EUS) and ERCP. *Endoscopy.* 2015; 47:176, <http://dx.doi.org/10.1055/s-0034-1391076>. PMID: 25635731.