



ORIGINAL ARTICLE

Immunomodulatory effects of black seeds and garlic on alloxan-induced *Diabetes* in albino rat

Bahaa K.A. Abel-Salam

Department of Biology, Deanship of Preparatory Year, Shaqra University, Shaqra, Saudi Arabia

Received 21 February 2011; accepted 25 July 2011

Available online 5 October 2011

KEYWORDS

Alloxan;
Leukocytes
proliferation;
IL-4;
IL-8;
TNF- α ;
Garlic;
Black seeds;
Albino rat

Abstract

Background: Alteration in the proliferation capacity of leukocytes and in the level of some cytokines, such as TNF- α , IL-4 and IL-8 have been suggested to associate with *Diabetes mellitus* in alloxan-induced diabetic rats given the potential immunomodulatory effects of black seeds and garlic.

Aim of the work: The aim of this study was to test the effects of these agents on the immune cells in alloxan-induced diabetic rats.

Methods: To this end, *Diabetes* was induced in albino rats by a single intraperitoneal injection of alloxan monohydrate (120 mg/kg of body weight). Diabetic rats were then fed normal diet or diet with black seeds or garlic for 28 days.

Results: The results showed significant increase in the numbers of monocytes and granulocytes, but with significant decreases in lymphocyte proliferation and the TNF- α , interleukin (IL)-4 and IL-8 levels in the diabetic group. Treatment of diabetic rats with black seeds or garlic induced significant amelioration in the numbers of monocytes and granulocytes, with significant increase in lymphocytes numbers and the TNF- α , IL-4 and IL-8 levels.

Conclusions: These results indicate the potential beneficial effects of black seeds and garlic as adjuvant treatment during treatment of *Diabetes*.

© 2011 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

Medicinal plants have been used to cure human illness since ancient times. Certain types of these plants are believed to promote positive health and maintain organism resistance against infection by re-establishing body equilibrium and

conditioning the body tissues. Among these plants, *Nigella sativa*, an annual herbaceous plant that belongs to family *Ranunculaceae*, has been used for thousands of years in traditional medicine. Its seeds are claimed to have anticancer, antileukaemic and antimicrobial effects.¹ The seeds have been found by some authors to possess an immunopotential activity.² Garlic (*Allium sativum*) also has traditional dietary and medicinal applications as an antimicrobial agent.³ Garlic is a common food spice widely distributed and used in all

E-mail address: manar_muhamad@yahoo.com

parts of the world as a spice and herbal medicine for the prevention and treatment of a variety of diseases, ranging from infections to heart diseases.⁴ Garlic is thought to have various pharmacological properties and medical applications. It is mainly consumed as a condiment in various prepared foods.⁵

Diabetes mellitus is a syndrome of impaired carbohydrate, fat, and protein metabolism caused by limited insulin secretion or lower tissue-sensitivity to insulin. This metabolic disorder is frequently diagnosed by hyperglycaemia, lipid abnormalities and vascular complications.⁶ In animals with experimental *Diabetes*, there is suppression of cellular immunity and reduced mitogenic response to several antigens.⁷ It has long been recognised that the inflammatory response in diabetic patients is impaired.^{8–13} Alloxan has been used to induce *Diabetes* into experimental animals. It acts through selective uptake by low affinity GLUT2 glucose transporter into the beta-cell leading to the destruction of the transporter protein by oxygen free radicals.¹⁴ Significant reduction in the body weight gain and hyperglycaemia are present in rats after alloxan injection.¹⁵

Alterations in lymphocytes are a common finding in both type I and type II *Diabetes*.¹⁶ Abnormalities of leukocyte function have been shown to occur during inflammation in alloxan-induced diabetic rats,⁸ including reduced number of leukocytes in inflammatory lesions¹⁷ and reduced production and transcription of pro-inflammatory (TNF- α) cytokines.^{8,9} Lymphocyte dysfunction might be the main cause of higher incidence of infections in diabetics, since an increased number of apoptotic lymphocytes were found in alloxan-induced diabetic animals and diabetic patients.¹⁸ Peripheral blood mononuclear cells from diabetic patients are reduced in their ability to produce cytokines¹⁹ and the proliferate responses of CD4+ T-cells to primary protein antigens are significantly reduced.²⁰ Some authors suggest that metabolic disturbances and therapeutic efforts to restore glucose metabolism may underlie part of the observed changes in lymphocyte subsets in either diabetic type.²¹ Elevated serum levels of IL-8, a potent chemoattractant for neutrophils and T-lymphocytes, were found in type 1 and 2 diabetic subjects, suggesting that this cytokine might contribute to the development of diabetic macroangiopathy.²²

The aim of this work was to evaluate the beneficial effects of black seeds and garlic on the altered leukocyte number and function as well as the levels of TNF- α , IL-4 and IL-8 in alloxan-induced diabetic rats with non-diabetics.

Materials and methods

Crude extract

Black seeds and garlic cloves purchased from the local market in El-Minia, Egypt were used. The animals were allowed free access to food 50 g of seeds and cloves daily during the course of the treatment, after fasting for about 12 h.

Animals

Male albino rats (*Rattus norvegicus*), weighing 120–150 g and of average age four months, were obtained from the Bio-

logical Supply Center, Theodore Bilharz Research Institute, TBRI, Cairo, Egypt and housed under specific pathogen-free conditions and maintained on a 12-h light-dark cycle, with food and water *ad libitum*. Animals were divided into four groups (10 animals each). The first group (control) was untreated. The second, third and fourth groups were artificially *Diabetes* induced by intraperitoneal injection with alloxan (Sigma Chemical Co., USA) 120 mg/kg of body weight, freshly dissolved in 5 mmol sterile normal saline.²³ The third group was intraperitoneal injected with alloxan once during the first week and allowed free access to food 50 g of black seeds daily during the course of the treatment, after fasting for about 12 h. The fourth group was treated similarly to the third group but with garlic instead of black seeds. Rats with blood glucose levels ≥ 250 mg/dl were considered to be diabetic. Plasma glucose levels in the control animals remained normal throughout the study.

Quantification of serum cytokines

The concentrations of TNF- α , IL-4 and IL-8 in collected serum samples were determined by enzyme-linked immunosorbent assay (ELISA) using commercially available kits according to the manufacturer's instructions (R&D Systems Inc., Minneapolis, MN, USA). The sensitivity of the assays was 15 pg/ml.

Leukocyte differential count

Freshly collected blood samples of about 20 μ l were spread on clean slides as a thin film using another smooth-edged glass slide. Each blood smear was left to air dry before being fixed with methanol for 2–3 min and then labelled by the number of the animal. Blood smears were stained with 10% Giemsa's stain (Aldrich) in buffered distilled water containing 0.021 M Na₂HPO₄/0.015 M KH₂PO₄. pH 7–7.2 for 30 min and kept away from sunlight. After that, the stain was removed by gentle washing with distilled water and the slides were air-dried at room temperature.²⁴ Using light microscopy at 400 \times magnification, different types of blood leukocytes were recorded. At least double smears for each blood sample were counted.

Statistical analysis

Data were analysed using SPSS program version 13.0. Statistical analysis of the obtained data was performed using one way analysis of variance (ANOVA) test followed by least square differences (LSD) analysis for comparison between means. Results were expressed as mean \pm standard error (SE). Values of $P < 0.05$ were considered statistically significant, while values of $P > 0.05$ were considered statistically non-significant.

Results

Quantification of TNF- α in sera

The TNF- α ELISA values demonstrated that the control group (1780.39 \pm 52.621 pg/ml) is higher than the diabetic group




Figure 1 Changes in TNF- α levels in alloxan groups versus control group and both black seeds and garlic treated groups.

(352.59 ± 4.204 pg/ml) which is lower than the diabetic group treated with either black seeds or garlic, counting 882.69 ± 3.468 pg/ml and 1100.28 ± 10.858 pg/ml, respectively (Fig. 1). This study showed a significant difference in TNF- α level between all groups ($P < 0.05$).

Quantification of serum IL-4

The results of ELISA in Fig. 2 showed that IL-4 recorded 850.83 ± 3.581 pg/ml in the control group. The diabetic group showed a lower level of IL-4 (144.47 ± 5.319 pg/ml) than the diabetic groups treated with either black seeds or garlic (379.20 ± 3.060 pg/ml and 452.16 ± 6.041 pg/ml, respectively). Statistical analysis showed a significant difference in IL-4 between the diabetic group and either the control group or the diabetic group treated with black seeds and garlic ($P < 0.05$).

Quantification of serum IL-8

In parallel to TNF- α and IL-4, ELISA tests of IL-8 showed that the diabetic group treated with black seeds (673.59 ± 3.243 pg/ml) or with garlic (731.02 ± 3.117 pg/ml) is higher than the diabetic group (232.10 ± 4.304 pg/ml), but lower than the control group (1090.21 ± 23.077 pg/ml). These differences can be seen in Fig. 3. Significant differences in IL-8 levels were monitored between the diabetic group and either the control group or the diabetic group treated with both black seeds and garlic ($P < 0.05$).




Figure 2 Black seeds and garlic modulation effects in IL-4 levels in diabetic male albino rats.




Figure 3 Changes in IL-8 levels in alloxan groups versus control group and both black seeds and garlic treated groups.




Figure 4 Black seeds and garlic modulation effects in granulocytes proliferation capacities in diabetic male albino rats.

Leukocyte proliferation

Fig. 4 shows that granulocytes percentage in the diabetic group (59.80 ± 0.200) is higher than both the control group (57.20 ± 0.374) and the diabetic group treated with black seeds (56.20 ± 1.800) and garlic (57.60 ± 0.245). On the other hand, lymphocytes percent in the diabetic group (28.40 ± 0.245) is slightly lower than both the control group (35.40 ± 0.245) and the diabetic group treated with black seeds and garlic recording 33.20 ± 0.200 and 33.80 ± 0.490 , respectively (Fig. 5). In parallel to granulocytes, Fig. 6 shows that the monocytes percentage in the diabetic group (11.80 ± 0.200) is higher than both the control group (7.40 ± 0.245) and the diabetic group treated with black




Figure 5 Changes in lymphocytes proliferation capacities in alloxan group versus control groups and both black seeds and garlic treated groups.




Figure 6 Black seeds and garlic modulation effects in monocytes proliferation capacities in diabetic male albino rats.

seeds and garlic (8.80 ± 0.200 and 8.60 ± 0.245 , respectively).

Statistical analysis showed that there is a significant difference in leukocyte count between the diabetic groups and either the control group or the diabetic group treated with both black seeds and garlic ($P < 0.05$).

Discussion

Alloxan has been known to be diabetogenic and induces DNA strand breaks in isolated rat pancreatic islets *in vitro* to cause *Diabetes mellitus*.²⁵ This is in accordance with our data, where alloxan injection showed statistically significant increase in glucose concentration versus control groups ($P < 0.05$). This rise in the glucose was significantly reduced after black seeds and garlic treatment compared to the glucose level in alloxan-treated groups. The mechanism of action of black seeds or garlic to decrease the glucose concentration in *Diabetes mellitus* is still unknown. Oral supplementation of black seeds after alloxan treatment resulted in lower plasma glucose levels and higher serum insulin values as compared with alloxan alone.²⁶ So, the decrease of glucose level in the black seeds and garlic treated groups revealed that these medicinal plants might be involved in reducing or inhibiting alloxan DNA strand breaks, increasing insulin secretion and consequently decreasing the glucose level.

Reinhold et al. showed that elevated glucose concentration significantly affected cytokine production.²⁷ In accordance, the results presented here showed changes in the level of the pro-inflammatory cytokine, TNF- α , the anti-inflammatory cytokine, IL-4, and the chemo-attractant cytokine, IL-8, between the diabetic group and the other groups. The changes in the TNF- α level could be due to the changes in insulin secretion, where insulin modulates the development of the inflammatory reaction to allergen challenge by its ability to modulate the production/release of TNF- α .¹⁰ In contrast to our data, a significant increase in IL-8 concentration in alloxan groups versus control groups (P value < 0.05) and a significant decrease after treatment compared with alloxan groups were detected.²⁸ In addition, circulating levels of IL-8 have been reported to be increased in patients with type 1 and type 2 *Diabetes* linking these chemokines with insulin resistance.²²

Leukocytes proliferation results suggested that alloxan-diabetic rates had an increase in granulocytes and

monocytes, and a decrease in lymphocytes proliferations. On the other hand, the diabetic rats treated with black seeds and garlic showed modulations in these changes. These are supported by a number of previous observations: (1) high glucose concentrations modified lymphocyte proliferate capacity²⁹; (2) Reinhold et al. showed that elevated glucose concentration significantly inhibits peripheral mononuclear cell proliferation by inhibiting the endogenous production of IL-2, IL-6 and IL-10²⁷; (3) hyperglycaemia in alloxan diabetic mice non-specifically impedes, but does not completely inhibit, the proliferation and growth of lymphocyte populations.³⁰

In contrast to our results, Martins et al. showed a significant reduction in the number of neutrophils and mononuclear leukocytes.¹⁵ They also mentioned that a complete recovery of the impaired responses was observed under the influence of insulin.

In conclusion, the data presented here showed that both black seeds and garlic could be involved in modulation of some immunological disorders induced by alloxan, when they were injected in combination.

Acknowledgement

I thank Dr. Mohamed Abuel-Maged (Zoology Department, Faculty of Science, Minia University, El-Minia, Egypt) for revising the manuscript.

References

- Hanafy MS, Hatem ME. Studies of the anti-microbial activity of *Nigella sativa* seeds (black comin). *J Ethnopharmacol.* 1991;34:275–8.
- El-Feki MA, Radwan ZA, Abdel-Moneim AA, Salah F. Effect of *Nigella sativa* and fish oil in alloxan diabetic rats. *Immunological studies. J Egypt Ger Soc Zool.* 1997;23:161–86.
- Ross ZM, O’Gara EA, Hill DJ, Sleightholme HV, Maslin DJ. Antimicrobial properties of garlic oil against human enteric bacteria: evaluation of methodologies and comparisons with garlic oil sulfides and garlic powder. *Appl Environ Microbiol.* 2001;67:475–80.
- Rivlin RS. Historical perspective on the use of garlic. *J Nutr.* 2001;131:951–4.
- Amagase H, Petesch BL, Matsuura H, Kasuga S, Itakura Y. Intake of garlic and its bioactive components. *J Nutr.* 2001;131:955–62.
- Maccuish AC, Urbaniak SJ, Campbell CJ, Duncan LJP, Irvine WJ. Phytohemagglutinin transformation and circulating lymphocyte subpopulations in insulin-dependent diabetic patients. *Diabetes.* 1974;23:708–12.
- Rayfield EJ, Ault MJ, Keusch GT, Brothers MJ, Nechemias C, Smith H. Infection and *Diabetes*: the case for glucose control. *Am J Med.* 1982;72:439–50.
- De Oliveira Martins J, Meyer-Pflug AR, Alba-Loureiro TC, Melbostad H, Cruz JWMC, Coimbra R, et al. Modulation of lipopolysaccharide induced acute lung inflammation: role of insulin. *Shock.* 2006;25:260–6.
- Martins JO, Zaroni FL, Martins DO, Coimbra R, Krieger JE, Jancar S, et al. Insulin regulates cytokines and intercellular adhesion molecule-1 gene expression through NF-kappaB activation in lipopolysaccharide-induced acute lung injury in rats. *Shock.* 2009;31:404–9.
- Martins JO, Ferracini M, Anger DB, Martins DO, Ribeiro Jr LF, Sannomiya P, et al. Signaling pathways and mediators in

- LPS-induced lung inflammation in diabetic rats: role of insulin. *Shock*. 2010;33:76–82.
11. Anjos-Valota EA, Martins JO, Oliveira MA, Casolari DA, Tostes RC, Fortes ZB, et al. Inhibition of tumor necrosis factor- α -induced intercellular adhesion molecule-1 expression in diabetic rats: role of insulin. *Inflamm Res*. 2006;55:16–22.
 12. Cavalher-Machado SC, Cuman RK, Sartoretto JL, Martins JO, de Lima WT, Martins MA, et al. The allergic inflammatory reaction in neonatal streptozotocin induced diabetic rats: evidence of insulin resistance and microvascular dysfunction. *Inflamm Res*. 2008;57:535–41.
 13. Ferracini M, Martins JO, Campos MRM, Anger DBC, Jancar S. Impaired phagocytosis by alveolar macrophages from diabetic rats is related to deficient coupling of LTs to the Fc γ R signaling cascade. *Mol Immunol*. 2010;47:1974–80.
 14. Elsner M, Tiedge M, Guldbakke B, Munday R, Lenzen S. Importance of the GLUT2 glucose transporter for pancreatic beta cell toxicity of alloxan. *Diabetologia*. 2002;45:1542–9.
 15. Martins JO, Campos CA, Cruz JW, Manzolli S, Alves VA, Vianna EO, et al. Insulin modulates cytokine release and selectin expression in the early phase of allergic airway inflammation in diabetic rats. *BMC Pulm Med*. 2010;10:39.
 16. von Kanel R, Mills PJ, Dimsdale JE. Short-term hyperglycemia induces lymphopenia and lymphocytes subset redistribution. *Life Sci*. 2001;69:255–62.
 17. Sannomiya P, Pereira MAA, Garcia-Leme J. Inhibition of leukocyte chemotaxis by serum factor in *Diabetes* mellitus: selective depression of cell response mediated by complement-derived chemoattractants. *Agents Actions*. 1990;30:369–76.
 18. Otton R, Soriano FG, Verlengia R, Curi R. *Diabetes* induces apoptosis in lymphocytes. *J Endocrinol*. 2004;145–56.
 19. Pickup JC, Chusney GD, Thomas SM, Burt D. Plasma interleukin-6, tumor necrosis factor alpha and blood cytokine production in type 2 *Diabetes*. *Life Sci*. 2000;67:291–300.
 20. Eibl N, Spatz M, Fischer GF, Mayr WR, Samstag A, Wolf HM, et al. Impaired primary immune response in type-1 *Diabetes*: results from a controlled vaccination study. *Clin Immunol*. 2002;103:249–59.
 21. Bouter KP, Meyling FH, Hoekstra JB, Masurel N, Erkelens DW, Diepersloot RJ. Influence of blood glucose levels on peripheral lymphocytes in patients with *Diabetes* mellitus. *Diabetes Res*. 1992;19:77–80.
 22. Zozulin D, Majchrzak A, Sobieska M, Wiktorowicz K, Wierusz W. Serum interleukin-8 level is increased in diabetic patients. *Diabetologia*. 1999;42:117–8.
 23. Yoruk M, Kanter M, Meral I, Agaoglu Z. Localization of glyco-gen in the placenta, fetal and maternal livers of cadmium exposed diabetic pregnant rats. *Biol Trace Elem Res*. 2004;96:217–26.
 24. Benett M, Daigle K. Temperature. Stress and the distribution of leukocytes in red-spotted newts, *Notophthalmus viridescens*. *J Comp Physiol*. 1983;153:81–3.
 25. Okamoto H. Ultrastructural aspects of streptozotocin cytotoxicity on rat pancreatic islets in vitro. *BioEssays*. 1985;2:15–21.
 26. Akhtar MS, Ali MR. The study of hypoglycemic activity of *Cuminum nigrum* seeds in normal and alloxan diabetic rabbits. *Planta Med*. 1984;2:81–5.
 27. Reinhold D, Bank U, Buhling F, Lendeckel U, Ulmer AJ, Flad HD, et al. Transforming growth factor h1 (TGFh1) inhibits DNA synthesis of PWM-stimulated PBMC via suppression of IL-2, IL-6 production. *Cytokine*. 1994;6:382–8.
 28. Ahmed MM, Bahaa KAA, Wafaey G. Protective effect of *Nigella Sativa* in alloxan induced diabetic rats. *Bull Eryp Soc Physiol Sci*. 2010;30:221–32.
 29. Sakowicz-Burkiewicz M, Kocbuch K, Grden M, Szutowicz A, Pawelczyk T. *Diabetes*-induced decrease of adenosine kinase expression impairs the proliferation potential of diabetic rat T lymphocytes. *Immunology*. 2006;118:402–12.
 30. Kozlov IA, Timofeeva EE, Zinger MG. Lymphopoietic function of mice with alloxan *Diabetes*. *Biull Eksp Biol Med*. 1986;101:407–9.