



Allergologia et immunopathologia

Sociedad Española de Inmunología Clínica,
Alergología y Asma Pediátrica

www.elsevier.es/ai



ORIGINAL ARTICLE

Food allergy in Catalonia: Clinical manifestations and its association with airborne allergens



J. Sánchez-López^a, V. Gázquez^b, N. Rubira^c, L. Valdesoiro^d, M. Guilarte^e,
A. Garcia-Moral^a, N. Depreux^f, L. Soto-Retes^g, M. De Molina^h, O. Luengoⁱ, R. Lleonartⁱ,
M. Basagaña^{f,*}

^a Unitat d'Al·lèrgia, Institut Clínic del Tòrax (ICT), Hospital Clínic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Universitat de Barcelona, Barcelona, Spain

^b Allergy Department, Hospital Universitari Joan XXIII, IISPV, Universitat Rovira i Virgili, Tarragona, Spain

^c Allergy Department, Hospital Platón, Barcelona, Spain

^d Pediatric Allergy Department, Corporació Sanitària Parc Taulí, Sabadell, Spain

^e Allergy Section, Medicine Department, Hospital Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain

^f Allergy Unit, Hospital Universitari Germans Trias i Pujol, Universitat Autònoma de Barcelona, Badalona, Spain

^g Servei de Pneumologia i Al·lèrgia, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain

^h Allergy Unit, Hospital General de l'Hospitalet, Consorci Sanitari Integral, L'Hospitalet de Llobregat, Spain

ⁱ Allergy Unit, Medicine Department, IDIBELL, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Spain

Received 28 January 2016; accepted 13 June 2016

Available online 31 August 2016

KEYWORDS

Allergy diagnosis;
Cross-sectional study;
Food allergy;
Food epidemiology;
Online database

Abstract

Background: Food allergy affects around 6% of the European population and its prevalence worldwide has been increasing in the last decades, but studies focused on investigating food allergy epidemiology in Europe are lacking.

Objective: The Cibus project was created to register the main culprit foods and their clinical manifestations in food allergic patients in Catalonia.

Methods: A specific online database was designed. Allergists from eight different Catalan hospitals registered the new diagnoses of food allergy.

Results: 618 food allergic patients were included. Egg and milk were the main elicitors in the early ages, while fruits and nuts were the most frequent in patients >14 years old. Fish was more frequent in children, while seafood and *Anisakis* allergy were more frequent in the >14-year-old group. Overall, peach was the most prevalent food eliciting an allergic reaction (10%). Food allergy diagnosis was reached using compatible clinical history and positive skin prick test to the involved food in 98% of cases. Globally, urticaria was the most frequently reported manifestation in our population (48.2%), followed by oral allergy syndrome (25.6%) and anaphylaxis (24.8%).

Conclusions: The Cibus project gives a full overview of the profile of food allergic patients in Catalonia and reinforces the predominance of plant food allergies in the Mediterranean area.

© 2016 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: maria.basagana.torrento@gmail.com (M. Basagaña).

<http://dx.doi.org/10.1016/j.aller.2016.06.004>

0301-0546/© 2016 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

Introduction

Food allergy affects around 6% of the European population¹ and regarding some recent studies, its prevalence worldwide has increased in the last decades.^{2,3} It has a great impact upon the patient's quality of life^{4,5} and represents a high economic burden to public health systems.⁶

Some risk factors have been identified, such as vitamin D deficiency, increased hygiene, or the age of exposure to foods, but the implication of other environmental factors still remains unclear.⁷ What seems evident is that the profile of food allergy appears to be different depending on the age and the geographical location, probably due to aeroallergen exposure and food habits.⁸

The main elicitors of food allergy worldwide are suggested to be cow's milk, egg, wheat, soy, peanut, tree nuts, fish, and shellfish^{1,9} but we must take into account that studies focused on investigating food allergy epidemiology in Europe are lacking and there is limited information on the prevalence and distribution of sensitisation to foods. Recent studies and reviews of the literature have provided very heterogeneous results.^{10–12}

The number of consultations in Allergy Units in Catalonia for food allergy has increased over the last years. According to data obtained from *Alergológica-1992 and 2005*, the consultations in Allergy Units in Catalonia and Balearic Islands for food allergy increased from 3.6% in 1992 to 6.05% in 2005.^{13,14} Thus, the "Cibus project" (Cibus is the Latin word for "food") was created to register the main culprit foods in our region and their clinical manifestations in order to outline the characteristics of our population.

Material and methods

Study design

A cross-sectional study was carried out in eight allergy outpatient clinics in Catalonia, Spain. Approval from the corresponding Ethics Committees was obtained. The recruitment period was from March 2008 to March 2010.

Patients included were those newly diagnosed of food allergy who agreed and signed the informed consent document (in the case of children the consent was obtained from their representatives). A website was created to report the data. Data collection included demographics, familiar and personal history of atopy (defined as having specific IgE (sIgE) to any common antigen as well as respiratory and/or skin disorders associated with IgE-mediated allergy), presence of other allergic diseases, food allergy manifestations, skin prick test (SPT) to food and respiratory allergens and results of sIgE measurements and/or oral food challenge (OFC) when available.

Diagnosis of food allergy

Food allergy diagnosis was established based on a detailed clinical history and a positive SPT and/or sIgE to the food/foods involved in the reaction/s. Clinical symptoms of the reported food allergies were classified as: oral allergy syndrome (OAS, defined as local

itching and/or angio-oedema of lips, tongue, palate and/or throat), urticaria and/or angio-oedema, anaphylaxis, dermatitis/eczema and/or gastrointestinal symptoms (GIS, including functional dyspepsia, crampy abdominal pain, nausea, vomiting and/or diarrhoea) occurring minutes to few hours after the food intake. Patients showing positive SPT and/or elevated food sIgE in the absence of any reported adverse reaction to that food were considered asymptomatic sensitisations.

Food allergy classification

For ease of interpretation of the results, foods reported were grouped as follows: egg, milk, wheat, other cereals (rice, oatmeal, rye, barley, maize), seeds (sesame, mustard, sunflower), legumes (lentil, chickpea, bean, soybean, pea, lupine), peanut, tree nuts (almond, hazelnut, walnut, chestnut, pinion, pistachio), vegetables (celery, onion, asparagus, green bean, lettuce, tomato, carrot, zucchini, spinach), fruits (peach, apple, pear, cherry, plum, pineapple, strawberry, fig, kiwi, avocado, banana, grape, melon, watermelon, orange, tangerine), meat (pork, beef, chicken), *Anisakis*, fish (hake, monkfish, sole, megrim, cod, tuna, salmon, sardine) and seafood (shrimp, mussel, clam, oyster, cuttlefish, squid).

In vivo and in vitro tests

Skin prick tests to standard panels of commercial airborne and food allergens were carried out. The standard airborne allergens panel included *Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, *Alternaria alternata*, *Cladosporium herbarum*, *Platanus acerifolia*, *Artemisia vulgaris*, *Lolium perenne* (Laboratorios LETI, Barcelona, Spain), dog and cat dander, *Cupressus arizonica*, *Olea europea*, *Parietaria Judaica* (Bial-Aristegui, Bilbao, Spain), *Penicillium sp* (Stallergenes, Antony, France). The standard food panel included milk, egg, peanut, hazelnut, walnut, soy, maize, wheat, shrimp, lettuce, tomato, green bean, kiwi, pineapple, banana, peach, mustard, hake, tuna (Laboratorios LETI) and *Anisakis simplex* (ALK-Abelló, Madrid, Spain). When food allergy was suspected to a food not included in the standard series, commercial extracts and/or fresh foods were used for skin testing. Fresh foods were tested by the prick-prick technique (P×P). All SPT were carried out on the volar side of the forearm using disposable prick lancets. SPT with saline and histamine 10 mg/mL were used as negative and positive control. A mean wheal diameter of 3 mm or greater obtained 15 minutes after puncture was considered a positive response.¹⁵

Food hypersensitivity was confirmed by sIgE measurements (UniCAP, Thermofisher, Uppsala, Sweden) in some centres. In these cases, sIgE levels >0.35 kU/L were regarded as positive.

Statistical analysis

SPSS software (version 18.0; SPSS, Chicago, IL, USA) was used for statistical analysis. For qualitative variables the frequency (percent) and its 95% confidence interval (95%

CI) was estimated. For quantitative variables, mean and standard deviation (SD) were calculated. Median, minimum and maximum values are given in those quantitative variables with large dispersion and atypical distribution. Qualitative variables were compared using the χ^2 test. Quantitative variables were compared using Student-*t* test. *P* values < 0.05 were considered significant.

Results

Patients

618 food allergic patients (mean age 30.51 years old (y.o.), SD 18.45) were included. Among them 155 were 14 years old or under. Full demographic data is presented in Table 1. In the population studied, frequency of familiar atopy, atopic dermatitis and asthma was higher in children (≤ 14 y.o.) compared to the >14 y.o. group ($P < 0.001$, $P < 0.001$ and $P = 0.004$, respectively) while rhinitis was significantly higher in the >14 y.o. group ($P < 0.001$). Regarding sensitisation to aeroallergens, the main sensitisers were house dust mites. Frequency of sensitisation to dog and cat dander, grass pollen, mugwort, olive tree and plane tree were significantly higher in the >14 y.o. group compared to the ≤ 14 y.o. group (Fig. 1).

Food allergy

Among the study population, a total of 1517 different food allergies were reported (2.45 ± 2.2 food allergies/patient), without statistical differences between the ≤ 14 y.o. and >14 y.o. groups.

In general, food allergies were markedly different between ≤ 14 y.o. and >14 y.o. groups. A full list of all food/food-group allergy frequencies is provided in Table 2. Egg and milk were the main elicitors in the younger group, while fruits and tree nuts were the most frequent in the older group. Frequency of fish allergy was higher in the ≤ 14 y.o. group compared to the >14 y.o. group ($P = 0.005$). On the contrary, frequency of seafood and *Anisakis* allergy were higher in the >14 y.o. group compared to the ≤ 14 y.o. group ($P = 0.003$ and $P < 0.001$, respectively).

A stratification of the food allergies in the ≤ 14 y.o. group was carried out, grouping them as shown in Fig. 2. Milk and egg relevance moved from 46% and 38% respectively in the youngest group (≤ 2 y.o.) to 1.3% and 2.8% in the >14 y.o. group.

On the contrary, fruits, peanut and tree nuts allergies increased progressively over the years. Other foods like seafood or *Anisakis* also underwent a moderate increase.

Regarding the food groups, the main elicitors were wheat (24%) for cereals, peanut (61%) for legumes, walnut (29%) for tree nuts, peach (35%) for fruits, lettuce (30%) for vegetables, hake (25%) for fish and shrimp (68%) for seafood. Overall, peach was the most prevalent food eliciting an allergic reaction (10%).

A specific analysis of nut allergy prevalence (including peanut, almond, hazelnut, walnut, chestnut, pinion, pistachio) showed that 92 subjects (43%) were allergic to only one nut, 51 (24%) to two different nuts, 34 (16%) to three different nuts and 35 (17%) were allergic to four or more




Figure 1 Frequency of sensitisation to aeroallergens. Statistical differences between groups are marked with * ($P < 0.05$) and ** ($P < 0.01$).

different nuts. Peanut, walnut and hazelnut was the most frequent association.

Diagnostic tools

SPT was the principal diagnostic tool used by the investigators. Food allergy diagnosis was reached using compatible clinical history and positive SPT to the involved food in the 98% of cases. SPT were negative in 24 cases (six milk, six cuttlefish, four clam, three shrimp, two banana, two watermelon, one salmon) and the diagnosis was obtained using other methods ($P \times P$, sIgE and/or OFC).

Also remarkable was that 33% of the positive SPTs were not associated with allergic symptoms. This percentage varied greatly depending on the food: For instance, egg, fish and *Anisakis* asymptomatic sensitisations were below 15%. On the contrary, cereals and seeds sensitisations had no clinical relevance at the time of the study in 77% and 68% respectively. sIgE was used to complement the diagnosis in 43% of cases, $P \times P$ in the 22%.

OFC was used to confirm the food elicitor in 33 allergic reactions (2%), 20 in the ≤ 14 y.o. group (milk, six; egg, five; lentil, chickpea, bean, pea, hazelnut, walnut, pinion, peach and melon, one) and 13 in the >14 y.o. group (almond and peach, two; egg, milk, soy, peanut, hazelnut, pistachio, onion, pineapple and hake, one).

Clinical manifestations

Globally, urticaria was the most frequently reported manifestation in our population (48.2%), followed by OAS (25.6%) and anaphylaxis (24.8%). Other symptoms like GIS, respiratory and eczema/dermatitis were scarcely reported (6.6%, 2.4% and 2.2% respectively).

The analysis of the four main symptoms (urticaria, OAS, anaphylaxis, GIS) by age groups (≤ 14 and >14 y.o.) showed that OAS and anaphylaxis frequency were significantly higher in the >14 y.o. group compared to the ≤ 14 y.o. group

Table 1 Demographic characteristics.

	≤14-year-old group (n= 155)	>14-year-old group (n= 463)	P
Mean age (SD)	6.95 (3.55)	38.4 (14.21)	
Sex female (%)	73 (47.1%)	264 (57%)	0.033
Familiar atopy	82 (52.9%)	124 (26.8%)	0.001
Asthma	53 (34.2%)	102 (22%)	0.004
Rhinitis	53 (34.2%)	247 (53.3%)	0.001
Atopic dermatitis	84 (54.2%)	46 (9.9%)	0.001

P values correspond to differences between both age groups. Significant differences are marked in bold.

Table 2 Food allergy frequency clustered age groups.

Food/food-group allergy	≤14-year-old group		>14-year-old group		P
	n	%	n	%	
Milk	48	30.97	6	1.30	0.001
Egg	50	32.26	13	2.81	0.001
Wheat	1	0.65	11	2.38	>0.05
Other cereals	3	1.94	8	1.7	>0.05
Legumes	9	5.81	15	3.2	0.001
Peanut	18	11.6	89	19.2	0.037
Tree nuts	31	20.00	172	37.15	0.001
Fruits	20	12.90	208	44.92	0.001
Vegetables	2	1.29	41	8.86	0.001
Fish	13	8.39	13	2.81	0.005
Seafood	9	5.81	69	14.90	0.003
Anisakis	1	0.65	53	11.45	0.001

Data is expressed as number of subjects allergic to a food or allergic to at least one of the foods within the corresponding food group. P values correspond to differences between both age groups: significant differences are marked in bold.




Figure 2 Food allergy frequency variation by age groups.

($P=0.016$ and $P<0.001$ respectively). The comparison of symptoms by age groups and food/food-groups showed that the higher frequency of anaphylaxis in the group of age >14 y.o. was because of tree nuts ($P<0.001$) and fruits

($P=0.043$). Although frequency of urticaria was not different between age groups, a separate analysis for foods showed that urticaria was more frequent in the ≤14 y.o. group for peanut ($P<0.001$), tree nuts ($P=0.006$) and seafood




Figure 3 Clinical manifestations of the main food/food groups elicitors of allergic reactions. GIS: Gastrointestinal symptoms; OAS: oral allergy syndrome.

($P=0.029$) groups compared to the >14 y.o. group. No differences were detected for egg, milk, *Anisakis* or fish allergy. The full distribution of symptoms by food/food-group and age group is shown in Fig. 3A and B.

Discussion

Prevalence of food allergies varies worldwide and geographical location is one of the main influencing factors, probably affected by aeroallergen exposure and food habits.^{11,16} Different sensitisation profiles can be possible even between the north and south regions of the same country.¹⁰ In Spain, the most recent epidemiological study in food allergy, *Alergológica-2005*,¹⁴ gives an overview of food allergies across the country, but there is little information about particularities of the different regions. The present study is, to our knowledge, the first epidemiological study on food allergies in Catalonia and the biggest one in terms of recruitment and number of reactions that has ever been described in Spain.

Regarding food/food groups involved in allergic reactions in our study, egg and milk are by far the main elicitors of food allergy in children, while fruits and tree nuts are the most frequent in the group >14 y.o. Other remarkable differences between both age groups are fish, seafood and *Anisakis*, being the first more frequent in children, unlike the other two. These data are consistent with those obtained from the

study *Alergológica-2005*¹⁴ and other studies performed in our geographical area^{17–19} as well as the data on the Spanish population obtained in the European Community Respiratory Health Survey (ECRHS) and Europrevall,^{11,20} but completely different from the food allergy patterns from the United States or United Kingdom, where peanut is by far the leading food.²¹ Differences in dietary habits might explain this difference. In fact, the most frequently involved nut in the allergic reactions of the studied population is walnut, which is usually associated with the Mediterranean diet. The individual food that triggered the largest number of reactions in the >14 y.o. group was peach. Peach is described as the main cause of food allergy in the adult population in countries like Spain or Italy, mainly due to its lipid transfer protein (LTP p 3).^{10,14} This protein is also responsible for multiple plant food-sensitisations and different clinical expressions, conforming the so-called LTP syndrome.^{22,23} Unfortunately, data about LTP sensitisation was not available in the present study, since at the time it was conducted (2008–2010) this diagnostic tool was not available in our allergy clinics.

The ECRHS study reported an unexpected no-correlation between a high sensitisation to foods and a high sensitisation to aeroallergens.²⁰ In our population, significant differences between aeroallergens sensitisation by age groups were detected for pollens. Grass, mugwort, olive and plane tree pollen sensitisations were more frequent in the older group. This finding correlates with a higher prevalence of allergy to plant foods in adulthood compared to childhood. In fact, mugwort and plane tree pollen sensitisations are clearly related to patients with plant food allergy,²⁴ and specifically to LTP sensitisation and might have a role in the development of plant food allergies.²⁵ Another point to remark is that the presence of mugwort pollen in Catalonia is scarce, so the high proportion of patients sensitised to this pollen might respond to a cross-reactive phenomenon related to LTPs, which does not exclude that they may have symptoms if exposed to higher concentrations of this pollen.²⁶ Even so, the main aeroallergen sensitisers in our population were the house dust mites, regardless of the age.

With regard to clinical manifestations of food allergy, the most frequently described in the literature are cutaneous reactions, followed by OAS, digestive symptoms and anaphylaxis.¹⁴ Similar results were found in our population; urticaria was the most frequently reported manifestation, followed by OAS and anaphylaxis. Only significant differences by age groups were detected for OAS and anaphylaxis, both more frequent in the >14 y.o. group. The increased frequency of anaphylaxis depended on peanut, nuts and seafood. These data are similar to those obtained in other epidemiological studies in patients from the Mediterranean area where milk and egg were the main responsible foods for anaphylaxis in children, while peach, vegetables and crustaceans were in adults.²⁷

In the majority of our patients, diagnosis was based on a positive case history and a positive SPT to the involved food. Clinical assessment is mandatory at all times, as positive SPTs alone can overestimate food allergy prevalence. In this study, 1/3 of the positive SPTs were not associated to an allergic reaction. Other methods of diagnosis (Pxp, specific IgE and/or OFC) were used when the SPT to the elicitor food was negative or as complementary diagnostic tools. Although OFC is the only conclusive tool in food allergy

diagnosis, in the present study it was only carried out in a very low percentage of patients. Double-blind placebo-controlled food challenge, the gold standard in food allergy, is a complex and hardly used technique that was not used by our clinicians. This aspect has already been reported in other studies, reflecting the distance between the usual methods of clinical practice and the general accepted scientific recommendations.²⁸⁻³⁰ Even so, the lack of OFC in this study should be considered a limitation itself.

In conclusion, the Cibus project provides one of the most extensive collections of newly diagnosed food allergy cases, with a complete evaluation of aeroallergens sensitisation that gives a full overview of the profile of food allergic patients in Catalonia. Given the similarity with other European and Spanish epidemiologic studies, it reinforces the predominance of plant food allergies in the Mediterranean area.

Sources of funding

A grant from SCAIC was obtained for the design and maintenance of the online database.

Conflict of interest

None of the authors have any conflict of interest regarding this paper.

Ethical disclosures

Confidentiality of data. Patients' data protection. The authors declare that they have followed the protocols of their work centre on the publication of patient data and that all the patients included in the study have received sufficient information and have given their informed consent in writing to participate in that study.

Right to privacy and informed consent. The authors have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence is in possession of this document.

Protection of human subjects and animals in research. The authors declare that no experiments were performed on humans or animals for this investigation.

Acknowledgments

The authors are grateful to SCAIC members who have voluntarily entered new cases of food allergy; we also thank Andrés Hidalgo for designing the online support for the database and the SCAIC Board for trusting in the project and funding the creation of the database.

References

- Nwaru BI, Hickstein L, Panesar SS, Roberts G, Muraro A, Sheikh A. Prevalence of common food allergies in Europe: a systematic review and meta-analysis. *Allergy*. 2014;69:992-1007.
- Ben-Shoshan M, Turnbull E, Clarke A. Food allergy: temporal trends and determinants. *Curr Allergy Asthma Rep*. 2012;12:346-72.
- Decker WW, Campbell RL, Manivannan V, Luke A, St Sauver JL, Weaver A, et al. The etiology and incidence of anaphylaxis in Rochester, Minnesota: a report from the Rochester Epidemiology Project. *J Allergy Clin Immunol*. 2008;122:1161-5.
- Avery NJ, King RM, Knight S, Hourihane JO. Assessment of quality of life in children with peanut allergy. *Paediatr Allergy Immunol*. 2003;14:378-82.
- Primeau MN, Kagan R, Joseph L, Lim H, Dufresne C, Duffy C, et al. The psychological burden of peanut allergy as perceived by adults with peanut allergy and the parents of peanut-allergic children. *Clin Exp Allergy*. 2000;30:1135-43.
- Patel DA, Holdford DA, Edwards E, Carroll NV. Estimating the economic burden of food-induced allergic reactions and anaphylaxis in the United States. *J Allergy Clin Immunol*. 2011;128:110-5.
- Moshe B-S, Turnbull E, Clarke A. Food allergy: temporal trends and determinants. *Curr Allergy Asthma Rep*. 2012;12:346-72.
- Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, et al., LEAP Study Team. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015;372:803-13.
- Allen JK, Koplin JJ. The epidemiology of IgE mediated food allergy and anaphylaxis. *Immunol Allergy Clin North Am*. 2012;32:35-50.
- Asero R, Antonicelli L, Arena A, Bommarito L, Caruso B, Colombo G, et al. Causes of food-induced anaphylaxis in Italian adults: a multi-centre study. *Int Arch Allergy Immunol*. 2009;150:271-7.
- Burney PG, Potts J, Kummeling I, Mills EN, Clausen M, Dubakiene R, et al. The prevalence and distribution of food sensitization in European adults. *Allergy*. 2014;69:365-71.
- Zuidmeer L, Goldhahn K, Rona RJ, Gislason D, Madsen C, Summers C, et al. The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol*. 2008;121:1210-8.
- Alergia a alimentos. In: *Alergológica. Factores epidemiológicos, clínicos y socioeconómicos de las enfermedades alérgicas*. Sociedad Española de Alergología e Inmunología Clínica & Alergia e Inmunología Abelló SA, editors. Madrid. NILO Industria gráfica. 1995. pp. 163-84.
- Fernández-Rivas M. Alergia a alimentos. In: *Alergológica 2005. Factores epidemiológicos, clínicos y socioeconómicos de las enfermedades alérgicas en España en 2005*. Madrid: Luzán5, S.A. de ediciones; 2006. pp. 227-53.
- Dreborg S, Frew A. Allergen standardization and skin tests. *EAACI position paper Allergy*. 1993;48:49-75.
- Rona RJ, Keil T, Summers C, Gislason D, Zuidmeer L, Sodergren E, et al. The prevalence of food allergy: a meta-analysis. *J Allergy Clin Immunol*. 2007;120:638-46.
- Crespo JF, Pascual C, Burks AW, Helm RM, Esteban MM. Frequency of food allergy in a pediatric population from Spain. *Pediatr Allergy Immunol*. 1995;6819:39-43.
- Joral A, Villas F, Garmendia J, Villareal O. Adverse reactions to food in adults. *J Investig Allergol Clin Immunol*. 1995;5:47-9.
- Alvarado MI, Pérez M. Study of food allergy in the Spanish population. *Allergol Immunopathol (Madr)*. 2006;34:185-93.
- Burney P, Summers C, Chinn S, Hooper R, Van Ree R, Lidholm J. Prevalence and distribution of sensitization to foods in the European Community Respiratory Health survey: a Europrevall analysis. *Allergy*. 2010;65:1182-8.
- Sicherer SH, Munoz-Furlong A, Sampson HA. Prevalence of peanut and tree nut allergy in the United States determined by means of a random digit dial telephone survey: a 5-year follow-up study. *J Allergy Clin Immunol*. 2003;112:1203-7.
- Pascal M, Muñoz-Cano R, Reina Z, Palacín A, Vilella R, Picado C, et al. Lipid transfer protein syndrome: clinical pattern, cofactor

- effect and profile of molecular sensitization to plant-foods and pollens. *Clin Exp Allergy*. 2012;42:1529–39.
23. Pastorello EA, Robino AM. Clinical role of lipid transfer proteins in food allergy. *Mol Nutr Food Res*. 2004;48:356–62.
 24. Cuesta-Herranz J, Barber D, Blanco C, Cistero-Bahima A, Crespo JF, Fernández-Rivas M, et al. Differences among pollen-allergic patients with and without plant food allergy. *Int Arch Allergy Immunol*. 2010;153:182–92.
 25. Palacin A, Gómez-Casado C, Rivas LA, Aguirre J, Tordesillas L, Bartra J, et al. Graph based study of allergen cross-reactivity of plant lipid transfer proteins (LTPs) using microarray in a multicenter study. *PLoS One*. 2012;7:e50799.
 26. Sánchez-López J, Tordesillas L, Pascal M, Muñoz-Cano R, Garrido M, Rueda M, et al. Role of Art v 3 in polinosis of patients allergic to Pru p 3. *J Allergy Clin Immunol*. 2014;133:1018–25.
 27. Rolla G, Mietta S, Raie A, Bussolino C, Nebiolo F, Galimberti M, et al. Incidence of food anaphylaxis in Piemonte region (Italy): data from registry of Center for Severe Allergic Reactions. *Intern Emerg Med*. 2013;8:615–20.
 28. Perry TT, Matsui EC, Kay Conover-Walker M, Wood RA. The relationship of allergen-specific IgE levels and oral food challenge outcome. *J Allergy Clin Immunol*. 2004;114:144–9.
 29. Vestege A, Mehl A, Rolinck-Werninghaus C, Staden U, Nocon M, Beyer K, et al. The predictive value of the skin prick test weal size for the outcome of oral food challenges. *Clin Exp Allergy*. 2005;35:1220–6.
 30. Bindslev-Jensen C, Ballmer-Weber BK, Bengtsson U, Blanco C, Ebner C, Houriane J, et al. Standardization of food challenges in patients with immediate reactions to foods—position paper from the European Academy of Allergology and Clinical Immunology. *Allergy*. 2004;59:690–7.