



ORIGINAL ARTICLE

## Five-assay microbiological system for the screening of antibiotic residues



Melisa Tumini, Orlando G. Nagel, Rafael L. Althaus\*

Facultad de Ciencias Veterinarias, Universidad Nacional del Litoral, R.P.L. Kreder 2805, 3080 Esperanza, Argentina

Received 24 July 2018; accepted 8 January 2019

Available online 2 May 2019

### KEYWORDS

Milk;  
Antibiotic;  
Detection;  
Classification;  
Bioassay;  
Microbiological  
system

**Abstract** A novel microbiological system in microtiter plates consisting of five bioassays is presented for the detection and classification of antibiotic residues in milk. The bioassays were optimized for the detection of beta-lactams (Bioassay B: *Geobacillus stearothermophilus*), macrolides (Bioassay M: *Bacillus megaterium* with fusidic acid), tetracyclines (Bioassay T: *B. megaterium* with chloramphenicol), quinolones (Bioassay Q: *Bacillus licheniformis*) and sulfamides (Bioassay QS: *B. licheniformis* with trimethoprim) at levels near the maximum residue limits (MRL). The response of each bioassay was interpreted visually (positive or negative) after 4–5.5 h of incubation. The system detects and classifies beta-lactams (5 µg/l of amoxicillin, 4 µg/l of ampicillin, 36 µg/l of cloxacillin, 22 µg/l of amoxicillin, 3 µg/l of penicillin, 114 µg/l of cephalexin, 89 µg/l of cefoperazone and 116 µg/l of ceftiofur), tetracyclines (98 µg/l of chlortetracycline, 92 µg/l of oxytetracycline and 88 µg/l of tetracycline), macrolides (33 µg/l of erythromycin, 44 µg/l of tilmicosin and 50 µg/l of tylosin), sulfonamides (76 µg/l of sulfadiazine, 85 µg/l of sulfadimethoxine, 77 µg/l of sulfamethoxazole and 87 µg/l of sulfathiazole) and quinolones (94 µg/l of ciprofloxacin, 98 µg/l of enrofloxacin and 79 µg/l marbofloxacin). In addition, the specificity values were high for B, T, Q (99.4%), M (98.8%) and QS (98.1%) bioassays. The control of antibiotics through this system can contribute to improving the quality and safety of dairy products.

© 2019 Published by Elsevier España, S.L.U. on behalf of Asociación Argentina de Microbiología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [ralthaus@fcv.unl.edu.ar](mailto:ralthaus@fcv.unl.edu.ar) (R.L. Althaus).

**PALABRAS CLAVE**

Leche;  
Antibiótico;  
Detección;  
Clasificación;  
Bioensayo;  
Sistema  
microbiológico

**Sistema microbiológico de cinco bioensayos para la detección de residuos de antibióticos**

**Resumen** Se presenta un novedoso sistema microbiológico en placas de microtitulación compuesto por 5 bioensayos para la detección y clasificación de residuos de antibióticos en leche. Los bioensayos fueron optimizados para la detección de betalactámicos (bioensayo B: *Geobacillus stearothermophilus*), macrólidos (bioensayo M: *Bacillus megaterium* con ácido fusídico), tetraciclinas (bioensayo T: *Bacillus megaterium* con cloranfenicol), quinolonas (bioensayo Q: *Bacillus licheniformis*) y sulfamidas (bioensayo QS: *Bacillus licheniformis* con trimetoprima), a niveles cercanos a los límites máximos de residuos (LMR). La respuesta de cada bioensayo se interpretó visualmente (positiva o negativa) después de 4 a 5,5 h de incubación. El sistema detecta y clasifica betalactámicos (5 µg/l de amoxicilina, 4 µg/l de ampicilina, 36 µg/l de cloxacilina, 22 µg/l de amoxicilina, 3 µg/l de penicilina, 114 µg/l de cefalexina, 89 µg/l de cefoperazona y 116 µg/l de ceftiofur), tetraciclinas (98 µg/l de clortetraciclina, 92 µg/l de oxitetraciclina y 88 µg/l de tetraciclina), macrólidos (33 µg/l de eritromicina, 44 µg/l de tilmicosina y 50 µg/l de tilosina), sulfamidas (76 µg/l de sulfadiazina, 85 µg/l de sulfadimetoxina, 77 µg/l de sulfametoxazol y 87 µg/l de sulfatiazol) y quinolonas (94 µg/l de ciprofloxacina, 98 µg/l de enrofloxacina y 79 µg/l de marbofloxacina). Además, los valores de especificidad fueron altos para los bioensayos B, T, Q (99,4%), M (98,8%) y QS (98,1%). El control de residuos de antibióticos mediante este sistema puede contribuir a mejorar la calidad e inocuidad de los productos lácteos.

© 2019 Publicado por Elsevier España, S.L.U. en nombre de Asociación Argentina de Microbiología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Introduction**

Various pathologies of milk-producing cattle, such as mastitis, metritis, pneumonia, enteritis and lameness<sup>26</sup>, are treated with penicillins<sup>6</sup>, ceftiofur<sup>16</sup>, oxytetracycline<sup>5,24</sup>, tylosin and enrofloxacin<sup>15</sup>, respectively.

Residues of these molecules could remain in the milk and cause adverse effects on consumer health, such as hypersensitivity reactions, allergy and resistance development<sup>17,22,8,1</sup>. In addition, the presence of these substances can delay acidification of fermented products, inhibit starter cultures and alter some properties involved in cheese ripening<sup>3,4</sup>.

In order to prevent the risks of antibiotic residues in milk, the European Community has established the maximum residue limits (MRL) and the procedures for their determination by Regulation EC 470/2009<sup>10</sup> and Regulation EC 37/2010<sup>11</sup>. Furthermore, in the USA, the Food and Drug Administration has determined the levels of tolerance of antibiotic residues in milk for consumer protection<sup>12</sup>.

Several screening tests have been developed for the control of antibiotic residues at permitted levels. In surveillance programs of veterinary drug residues, microbiological methods are widely used<sup>23</sup>, because they are inexpensive, allow large number of samples and do not require microbiology-trained staff. Moreover, these methods have detection limits similar to the MRL for beta-lactams, tetracyclines and sulfamides in milk; however, they do not detect residues of other antibiotics such as macrolides and quinolones<sup>14,2</sup>.

Thus, Nagel et al.<sup>20</sup> proposed bioassays that use several bacteria-tests with specific sensitivity for different groups of antibiotics used in the disease treatment of

dairy cattle. These bioassays integrate a Microbiological System with dichotomous responses (positive – negative) of easy interpretation in a maximum time of 6 h. This detection methodology detects and classifies residues in four antibiotic groups (beta-lactams, tetracyclines, sulfonamides and quinolones), but with cross-specificity to some macrolides (tylosin, tilmicosin, erythromycin) and aminoglycosides (neomycin).

Therefore, the aim of this study was to evaluate a Microbiological System in microtiter plates (MSmp) to detect and classify antibiotic residues in five groups (beta-lactams, tetracyclines, sulfonamides, quinolones and macrolides) to provide a dichotomous response (positive or negative) in less than 6 h, to be used routinely by milk quality control laboratories.

**Materials and methods****Design of a microbiological system in microtiter plates (MSmp)**

With the purpose of detecting and classifying antibiotic residues in milk, a microbiological system consisting of five bioassays was designed. This System is constituted by Bioassay B (beta-lactams), Bioassay T (tetracyclines), Bioassay M (macrolides), Bioassay Q (quinolones) and Bioassay QS (quinolones and sulfonamides). Table 1 shows culture media, concentration of sensitivity-enhancing agents, spore concentration of bacteria-tests, indicators, pH, incubation conditions for each bioassay that integrates the MSmp.

**Table 1** Composition of bioassays that integrate the microbiological system evaluated

Bioassays	Bacteria test	Agar medium	Incubation	Color response	
				Negative	Positive
B	<i>G. stearothermophilus</i> subsp. calidolactis C 953 (2 × 10 <sup>6</sup> spores/ml)	Brilliant black (100 mg/l) toluidine blue (270 mg/l), pH 8.2	64 °C, 4 h	Yellow	Black
T	<i>B. megaterium</i> ATCC 9885 (2.8 × 10 <sup>8</sup> spores/ml)	Brilliant black (200 mg/l) toluidine blue (10 mg/l), pH 8.5, chloramphenicol (2000 µg/l)	45 °C, 5 h	Yellow	Black
M	<i>B. megaterium</i> ATCC 9885 (2.8 × 10 <sup>8</sup> spores/ml)	Brilliant black (200 mg/l) toluidine blue (10 mg/l), pH 8.0, fusidic acid (200 µg/l)	45 °C, 5 h	Yellow	Black
Q	<i>B. licheniformis</i> ATCC 14580 (4.2 × 10 <sup>10</sup> spores/ml)	Triphenyl tetrazolium (200 mg/l) toluidine blue (10 mg/l), pH 7.0	45 °C, 5.5 h	Pink	Blue
QS	<i>B. licheniformis</i> ATCC 14580 (4.2 × 10 <sup>10</sup> spores/ml)	Triphenyl tetrazolium (200 mg/l) toluidine blue (10 mg/l), pH 7.0, trimethoprim (100 µg/l)	45 °C, 5.5 h	Pink	Blue

B: beta-lactams; T: tetracyclines; M: macrolides; Q: quinolones; QS: quinolones-sulfamides.

### Preparation of bioassays

Mueller Hinton Agar (38 g/l, Biokar<sup>®</sup>, Ref. 10272, France) fortified with glucose (10 g/l, Sigma Aldrich<sup>®</sup>, Ref. G8270, USA) was used and sterilized at 121 °C for 15 min. Then, each bioassay was added with redox indicators, spores of specific microorganisms, and sensitivity-improving substances (chloramphenicol, fusidic acid or trimethoprim), as described in Table 1. Subsequently, 100 µl of medium was dispensed into each of the 96 wells of the microtiter plate with an electronic dispenser (Eppendorf Research<sup>®</sup> Pro, Hamburg, Germany) and then, the plates were heat-sealed with aluminum foil coated with polypropylene and stored in a refrigerator (4 °C) until use.

### Antibiotic solutions and fortified samples

For the preparation of antimicrobial solutions, standard antibiotics from Sigma Chemical Co. (St. Louis, MO, USA) were used. Aqueous solutions for each antibiotic were prepared in 10 ml volumetric flasks at a concentration of 1000 mg/l at the time of analysis to avoid possible alteration of molecules and/or their properties.

### Dose-response curves

The dose-response curves were constructed for eight beta-lactams (amoxicillin, ampicillin, cloxacillin, oxacillin, penicillin 'G', cefoperazone, ceftiofur and cephalixin), three tetracyclines (chlortetracycline, oxytetracycline and tetracycline), three macrolides (erythromycin, tylosin and tilmicosin), three quinolones (ciprofloxacin, enrofloxacin and marbofloxacin), four sulfonamides (sulfadiazine, sulfadimethoxine, sulfamethoxazole and sulfathiazole) and

three aminoglycosides (kanamycin, neomycin and streptomycin), according to the Codex Alimentarius guidelines<sup>9</sup>.

For this purpose, 16 replicas of 12 concentrations were tested to obtain negative results in the first two concentrations and positive results in the last two concentrations tested. Then, 50 µl milk samples fortified with antibiotics were added to each bioassay well. The microplates were sealed with adhesive film and incubated under conditions of time and temperature to allow the color change of the control samples (Table 1).

### Interpretation of results

Bioassay responses were visually interpreted by three specialists independently. Each analysis was classified in terms of "positive" or "negative" when there were at least two coincident results.

Milk samples not containing or containing antibiotics at concentrations lower than the method detection limit will allow the growth of the test bacterium, accompanied by a change in the potential redox of the culture medium, manifested by a change in the coloration of the redox indicators. Thus, Bioassays B, M and T, which contained brilliant black and toluidine blue, changed to yellow (negative result), whereas Bioassays Q and QS, which contained triphenyl-tetrazolium and toluidine blue, changed to pink (negative result) after incubation (Table 1).

Conversely, when a milk sample contains antibiotic residues at levels above the detection limit of each molecule for a bioassay, it will not allow the growth of the microorganism, and thus there will be no change in the redox potential and the original color of the bioassay will remain. Thus, the persistence of the black color in Bioassays B, T and M indicated the presence of antibiotics in milk, whereas the

persistence of the blue color in Bioassays Q and QS also indicated the presence of antibiotics in milk.

### Calculation of detection limits

The frequencies of positive results were analyzed according to the following logistic regression model (StatGraphics Centurion X, 2007)<sup>25</sup>:

$$L_{ij} = \text{Logit} [P_{ij}] = \beta_0 + \beta_1[ATB]_i + \varepsilon_{ij} \quad (1)$$

where:  $L_{ij}$  = logistic linear model;  $[P_{ij}]$  = logit  $[Pp/(1 - Pp)]$ : (probability of a "positive" response/probability of a "negative" response),  $\beta_0$  and  $\beta_1$  = parameters calculated by the logistic model;  $[ATB]_i$  = antibiotic concentration ( $i = 1, 2, \dots, 12$  levels); and  $\varepsilon_{ij}$  = residual error.

The detection limits were calculated as the concentration producing 95% of positive frequencies<sup>7,9</sup>.

### Specificity of each bioassay

A total of 160 milk samples from Holstein Friesian cows 60–90 days postpartum with normal values of chemical composition (protein: 2.10–5.75 g/100 ml, fats: 2.38–6.03 g/100 ml, lactose: 3.63–5.17 g/100 ml, total solids: 9.31–12.86 g/100 ml), Bacterial Colony Count (BCC: 13.000–99.000/ml) and Somatic Cell Count (SCC: 2.000–392.000/ml) were used according to the Codex Alimentarius<sup>9</sup>. The chemical composition, SCC and BCC were determined using Milko Scan FT-120 (Foss Somatic, USA), Fossomatic 90 (Foss Electric, USA) and BactoScan 8.000S (Foss Electric, USA), respectively. Milk samples were analyzed with the five bioassays (B, T, M, Q and QS) in triplicate. For this purpose, 50  $\mu$ l of milk samples was added in each single well of each bioassay and then incubated in a water floating bath at a temperature-time for each bioassay (Table 1). Samples were analyzed simultaneously by the Delvotest<sup>®</sup> SP method (DSM, Delft, the Netherlands), which is considered a reference method. Then, specificity (%) was calculated using the following mathematical equation:

$$\text{Specificity} = \frac{\text{negative}}{\text{total samples}} \times 100 \quad (2)$$

## Results and discussion

### Detection limits of the bioassays

Table 2 shows the results obtained by applying the logistic regression model for the 24 antibiotics tested with each of the five bioassays (B, T, M, Q and QS). The adjustments obtained by the logistic regression model were good, since the concordance coefficients were between 98.7% and 99.4%.

Coefficient  $\beta_1$  indicates the slope of the dose–response curve. Higher values of this coefficient indicate good sensitivity of the bacteria-test to detect antibiotics in milk. Bioassay B had a higher coefficient  $\beta_1$  for beta-lactam antibiotics (between 0.0616 and 5.5301) in addition to tylosin (0.1458) and neomycin (0.0099). Bioassay T showed high values of  $\beta_1$  for the three tetracyclines studied ( $\beta_{1\text{Chlortetracycline}} = 0.1637$ ;  $\beta_{1\text{Oxitetra-cycline}} = 0.1258$ ;

$\beta_{1\text{Tetracycline}} = 0.0809$ ). Bioassay M had high values of this coefficient for macrolides ( $\beta_{1\text{Erythromycin}} = 0.2181$ ;  $\beta_{1\text{Tilmicosin}} = 0.1425$ ;  $\beta_{1\text{Tylosin}} = 0.1198$ ). Bioassay Q showed similar coefficients  $\beta_1$  for the three quinolones (between 0.1060 and 0.1106), and bioassay QS (with trimethoprim) also exhibited high coefficients  $\beta_1$  for sulfonamides (between 0.2052 and 0.2707).

The detection limits (DL) calculated for each molecule by the logistic regression model and the respective MRL<sup>9</sup> are summarized in Table 3. For beta-lactams, Bioassay B showed detection limits (DL) similar to their MRL (amoxicillin: 5  $\mu$ g/l, ampicillin: 4  $\mu$ g/l, cloxacillin 36  $\mu$ g/l, oxacillin: 22  $\mu$ g/l, penicillin 'G' 3  $\mu$ g/l, cephalixin 114  $\mu$ g/l, cefoperazone: 89  $\mu$ g/l, ceftiofur: 116  $\mu$ g/l). By contrast, in Bioassays M, T, Q and QS, beta-lactams had to be present in milk at levels above 5 MRL to produce positive results.

The detection limits for beta-lactams calculated in Bioassay B (Table 3) were lower than those reported by Nagel et al.<sup>20</sup> for a bioassay in microtiter plates using *Geobacillus stearothermophilus* (4  $\mu$ g/l of penicillin, 14  $\mu$ g/l of amoxicillin, 8  $\mu$ g/l of ampicillin, 49  $\mu$ g/l of cloxacillin, 25  $\mu$ g/l of oxacillin, 190  $\mu$ g/l of ceftiofur, 190  $\mu$ g/l of cephalixin and 140  $\mu$ g/l of cefoperazone). For assays in Petri dishes using *G. stearothermophilus*, Nouws et al.<sup>21</sup> reported lower detection levels (2  $\mu$ g/l of penicillin, 3  $\mu$ g/l of amoxicillin, 2  $\mu$ g/l of ampicillin, 15  $\mu$ g/l of cloxacillin, 20  $\mu$ g/l of oxacillin, 30  $\mu$ g/l of ceftiofur, 45  $\mu$ g/l of cephalixin, 30  $\mu$ g/l of cefoperazone). Similarly, Gaudin et al.<sup>13</sup> reported DL that were similar to those obtained in this work (Table 3) in Petri dishes (12–16  $\mu$ g/l of penicillin "G", 6–5  $\mu$ g/l of ampicillin, 16  $\mu$ g/l of amoxicillin, 50  $\mu$ g/l of cloxacillin, 30  $\mu$ g/l of oxacillin, 125  $\mu$ g/l of ceftiofur, 80  $\mu$ g/l of cefoperazone and 30  $\mu$ g/l of cephalixin).

With respect to Bioassay T, it should be noted that the DL obtained were similar to the MRL (100  $\mu$ g/l) for chlortetracycline (98  $\mu$ g/l) and oxytetracycline (92  $\mu$ g/l) and slightly higher for tetracyclines (128  $\mu$ g/l) due to the improvement in sensitivity of *B. megaterium* with chloramphenicol<sup>29</sup>. However, Bioassay M using the same microorganism without the addition of chloramphenicol showed positive results when tetracyclines were present at greater levels than 2 times the MRL (Table 3).

The detection limits of tetracyclines for Bioassay T (Table 3) were slightly lower than those calculated by Nagel et al.<sup>18,20</sup> when using a bioassay with *B. cereus* spores in microtiter plates (chlortetracycline: 330  $\mu$ g/l, oxytetracycline: 110  $\mu$ g/l, tetracycline: 110  $\mu$ g/l). In previous studies, Tumini et al.<sup>28</sup> found detection limits that were similar to those obtained in this work for a bioassay containing *B. pumilus* (chlortetracycline 117  $\mu$ g/l, oxytetracycline: 142  $\mu$ g/l, tetracycline: 105  $\mu$ g/l).

For assays in Petri dishes, Nouws et al.<sup>21</sup> obtained values lower than those calculated in this work when using *B. cereus* after 18 h incubation (10  $\mu$ g/l of chlortetracycline, 30  $\mu$ g/l of oxytetracycline and 30  $\mu$ g/l of tetracycline). Tsai and Kondo<sup>27</sup> and Gaudin et al.<sup>13</sup> reported higher levels than those calculated in this study (Table 3) when using assays in Petri dishes (chlortetracycline: 100–200  $\mu$ g/l, oxytetracycline: 200–250  $\mu$ g/l, tetracycline: 200–250  $\mu$ g/l).

With regard to macrolides, Bioassay M presented adequate detection limits for erythromycin (33  $\mu$ g/l), tilmicosin (49  $\mu$ g/l) and tylosin (50  $\mu$ g/l). The sensitivity to

**Table 2** Coefficients calculated by the Logistic regression model for the dose–response curve

Antibiotics	Bioassay B		Bioassay T		Bioassay M		Bioassay Q		Bioassay QS	
	$\beta_0$	$\beta_1$	$\beta_0$	$\beta_1$	$\beta_0$	$\beta_1$	$\beta_0$	$\beta_1$	$\beta_0$	$\beta_1$
<i>Beta-lactams</i>										
Amoxicillin	-6.3773	<b>1.9392</b>	-4.9124	0.0545	-4.9057	0.0537	-9.1554	0.0219	-8.7600	0.0261
Ampicillin	-8.8622	<b>3.0737</b>	-6.1354	0.0937	-6.1772	0.0924	-4.7193	0.0384	-4.4617	0.0377
Cloxacillin	-11.836	<b>0.4084</b>	-5.1654	0.0335	-5.1424	0.0389	-9.1554	0.0219	-8.3468	0.0206
Oxacillin	-16.313	<b>0.8706</b>	-4.8035	0.0339	-4.8032	0.0325	-7.2449	0.0267	-7.2674	0.0259
Penicillin ‘‘G’’	-15.299	<b>5.5301</b>	-10.916	0.0752	-10.955	0.0724	-5.0872	0.1476	-5.0768	0.1456
Cephalexin	-9.7361	<b>0.1112</b>	-4.8526	0.0120	-4.8602	0.0117	-10.237	0.0151	-10.364	0.0153
Cefoperazone	-5.9080	<b>0.0995</b>	-4.8658	0.0033	-4.8596	0.0032	-6.3529	0.0129	-6.3398	0.0127
Ceftiofur®	-4.2313	<b>0.0616</b>	-11.861	0.0239	-11.859	0.0233	-7.1114	0.0128	-7.1498	0.0127
<i>Aminoglycosides</i>										
Kanamycin	-18.733	0.0026	-15.398	0.0246	-15.332	0.0259	-7.3300	0.0017	-7.8288	0.0017
Neomycin	-10.903	0.0099	-2.9801	0.0061	-2.9754	0.0064	-8.8972	0.0002	-8.9971	0.0002
Streptomycin	-12.197	0.0045	-13.685	0.0276	-13.537	0.0262	-11.057	0.0002	-10.906	0.0002
<i>Macrolides</i>										
Erythromycin	-6.5414	0.0337	-5.6254	0.0907	-4.1801	<b>0.2181</b>	-5.5534	0.0079	-5.5484	0.0080
Tilmicosin	-5.7655	0.0262	-6.6341	0.0901	-4.0196	<b>0.1428</b>	-10.822	0.0146	-10.819	0.0141
Tylosin	-4.6664	0.1454	-7.4681	0.0998	-3.0152	<b>0.1198</b>	-8.7288	0.0205	-8.7291	0.0209
<i>Sulfamides</i>										
Sulfadiazine	-10.884	0.0007	-14.268	0.0060	-14.188	0.0059	-12.282	0.0102	-16.282	<b>0.2123</b>
Sulfadimethoxine	-16.567	0.0023	-18.394	0.0103	-18.321	0.0102	-11.982	0.0096	-21.825	<b>0.2541</b>
Sulfamethoxazole	-15.294	0.0044	-14.895	0.0074	-14.961	0.0075	-13.912	0.0120	-15.912	<b>0.2052</b>
Sulfathiazole	-19.432	0.0051	-13.701	0.0054	-13.698	0.0055	-11.709	0.0119	-23.709	<b>0.2707</b>
<i>Tetracyclines</i>										
Chlortetracycline	-4.6026	0.0012	-13.119	<b>0.1637</b>	-9.3559	0.0576	-11.647	0.0137	-11.723	0.0134
Oxytetracycline	-10.811	0.0072	-8.6151	<b>0.1258</b>	-10.999	0.0691	-7.3439	0.0121	-7.3774	0.0123
Tetracycline	-18.384	0.0124	-7.4310	<b>0.0809</b>	-10.174	0.0591	-7.1065	0.0088	-7.1163	0.0085
<i>Quinolones</i>										
Ciprofloxacin	-6.3695	0.0337	-15.356	0.0521	-15.329	0.0518	-7.0710	<b>0.1060</b>	-7.0712	<b>0.1097</b>
Enrofloxacin	-12.297	0.0037	-3.9754	0.0064	-3.9535	0.0067	-7.9332	<b>0.1106</b>	-7.9877	<b>0.1112</b>
Marbofloxacin	-7.3085	0.0015	-3.9744	0.0066	-3.9986	0.0067	-6.5049	<b>0.1091</b>	-6.5235	<b>0.1035</b>

B: beta-lactams; T: tetracyclines; M: macrolides; Q: quinolones; QS: quinolones-sulfonamides.  $\beta_0$  and  $\beta_1$ : parameters calculated by the logistic model.

$\beta_1$  coefficient values in bold for the same group of antibiotics indicate greater bioassay sensitivity compared to other bioassays of MSmp.

these compounds can be attributed to the incorporation of fusidic acid in the culture medium containing *B. megaterium* spores<sup>30</sup>. Conversely, macrolides must be present in milk at concentrations between 2 and 3 times the MRL to produce positive results in bioassay T (Table 3).

The detection limits of macrolides obtained by using Bioassay M (Table 3) were higher than those reported by Nouws et al.<sup>21</sup> for erythromycin (10 µg/l) and tilmicosin (6 µg/l) but lower than those reported for tylosin (100 µg/l) when using *Kocuria rhizophila* in Petri dishes. Moreover, Gaudin et al.<sup>13</sup> estimated similar DL for erythromycin (20–30 µg/l) and tilmicosin (25–50 µg/l) and higher ones for tylosin (150–200 µg/l) in a method in Petri dishes containing the same microorganism. However, both Nouws et al.<sup>21</sup> and Gaudin et al.<sup>13</sup> obtained a response after 18–24 h of incubation.

The residues of quinolones in milk were detected by Bioassays Q and QS, with DL similar to the MRL for

ciprofloxacin (Q: 94 µg/l QS: 91 µg/l), enrofloxacin (Q and QS: 98 µg/l) and marbofloxacin (Q: 87 µg/l QS: 91 µg/l) due to the good sensitivity of *B. licheniformis* for quinolones<sup>31</sup>.

For quinolone residues, the DL of bioassays Q and QS (Table 3) were lower than those reported by Nagel et al.<sup>20</sup> when using a bioassay in microtiter plates containing *B. subtilis* spores (150 µg/l of ciprofloxacin, 160 µg/l of enrofloxacin and 160 µg/l of marbofloxacin). By using bioassays in Petri dishes containing *Escherichia coli* cells and after 24 h of incubation, Nouws et al.<sup>21</sup> (4 µg/l of enrofloxacin and 5 µg/l marbofloxacin) and Gaudin et al.<sup>13</sup> detected low concentrations of quinolones in milk (10 µg/l of ciprofloxacin, 20 µg/l of enrofloxacin and 30 µg/l of marbofloxacin).

For sulfonamides, the addition of trimethoprim into the culture medium containing *B. licheniformis* spores (Bioassay QS) significantly improved the detection of sulfadiazine (76 µg/l), sulfadimethoxine (85 µg/l), sulfamethoxazole (77 µg/l) and sulfathiazole (87 µg/l) with respect to bioassay Q without trimethoprim (Table 3).

**Table 3** Detection limits ( $\mu\text{g/l}$ ) of microbiological system in microplates

Antibiotics	MRL <sup>a</sup>	Bioassays				
		B	TC	MC	Q	QS
<i>Beta-lactams</i>						
Amoxicillin	4	<b>5</b>	150	140	550	540
Ampicillin	4	<b>4</b>	97	100	200	190
Cloxacillin	30	<b>36</b>	242	208	550	550
Oxacillin	30	<b>22</b>	229	250	380	390
Penicillin "G"	4	<b>3</b>	184	200	54	55
Cephalexin	100	<b>114</b>	650	670	870	870
Cefoperazone	50	<b>89</b>	2360	2500	720	730
Ceftiofur <sup>®</sup>	100	<b>116</b>	500	500	780	795
<i>Aminoglycosides</i>						
Kanamycin	100	8300	620	590	5900	6020
Neomycin	1500	<b>1400</b>	<b>460</b>	<b>450</b>	59 300	59 500
Streptomycin	200	3400	490	500	70 100	69 200
<i>Macrolides</i>						
Erythromycin	40	280	94	<b>33</b>	1070	1100
Tilmicosin	50	350	106	<b>44</b>	700	760
Tylosin	50	<b>52</b>	104	<b>50</b>	450	400
<i>Sulfamides</i>						
Sulfadiazine	100	18 100	2400	2400	1500	<b>76</b>
Sulfadimethoxine	100	8200	1800	1800	1500	<b>85</b>
Sulfamethoxazole	100	4100	1900	1900	1400	<b>77</b>
Sulfathiazole	100	4400	2500	2500	1200	<b>87</b>
<i>Tetracyclines</i>						
Chlortetracycline	100	5900	<b>98</b>	214	1060	1100
Oxytetracycline	100	1800	<b>92</b>	202	850	840
Tetracycline	100	1700	<b>88</b>	222	1200	1100
<i>Quinolones</i>						
Ciprofloxacin	100	4100	290	300	<b>94</b>	<b>91</b>
Enrofloxacin	100	4000	590	560	<b>98</b>	<b>98</b>
Marbofloxacin	75	6800	1050	1100	<b>79</b>	<b>77</b>

B: beta-lactams; TC: tetracyclines; MC: macrolides; Q: quinolones; QS: quinolones-sulfonamides.

<sup>a</sup> MRL: maximum residue limit ( $\mu\text{g/l}$ ).

Detection limit values highlighted in bold for each bioassay indicate levels close to their respective MRL.

The detection limits for sulfonamides in milk calculated when using Bioassay QS (Table 3) were lower than those reported by Nagel et al.<sup>20</sup> for sulfadiazine (160  $\mu\text{g/l}$ ), sulfadimethoxine (260  $\mu\text{g/l}$ ), sulfamethoxazole (120  $\mu\text{g/l}$ ) and sulfathiazole (110  $\mu\text{g/l}$ ) when using a bioassay with *B. subtilis* and trimethoprim in the culture medium. However, in an assay in Petri dish with *B. subtilis* spores and requiring 18–24 h of incubation, Nouws et al.<sup>21</sup> obtained lower levels for residues of sulfadiazine (30  $\mu\text{g/l}$ ) and sulfadimethoxine (20  $\mu\text{g/l}$ ) than those reported in Table 3, while Gaudin et al.<sup>13</sup> obtained higher detection limits for sulfadiazine (100  $\mu\text{g/l}$ ), sulfadimethoxine (175  $\mu\text{g/l}$ ), sulfamethoxazole (100  $\mu\text{g/l}$ ) and sulfathiazole (1000  $\mu\text{g/l}$ ).

### Specificity of the bioassays


The results of specificity for each bioassay are detailed in Table 4. It can be observed that bioassays B, T and

**Table 4** Specificity of the microbiological system

Bioassays	Milk samples	Positive	Negative	Specificity (%)
B	160	1	159	99.4
T	160	1	159	99.4
M	160	2	158	98.8
Q	160	1	159	99.4
QS	160	3	157	98.1
Delvotest <sup>®</sup> SP	160	1	159	99.4

Specificity (%): (negative/total samples)  $\times$  100.

Q had high specificity (99.4%), as determined by the Delvotest<sup>®</sup> SP method, and that the specificity of bioassays M (98.8%) and QS (98.1%) was slightly lower than that of the other methods. It should be noted that Nagel et al.<sup>18,19</sup> and Tumini et al.<sup>28</sup> reported specificities



**Figure 1** Detection pattern (DL/MRL) for microbiological five-bioassay system. Circle 1 = 0.1 MRL, Circle 2 = MRL, Circle 3 = 10 MRL and Circle 4 = 100 MRL.

between 99.0–99.5% (*B. cereus*), 95.8–98.9% (*B. subtilis*) and 97.9–98.9% (*B. pumilus*) respectively when optimizing microbiological methods in microtiter plates.

**Cross-specificity of the five-bioassay system**

In order to simultaneously visualize the detection profiles of the five bioassays for the 24 antibiotics studied, we constructed Figure 1. The radial diagram was constructed in logarithmic scale due to the large amplitude in the range of the detection system (Table 3) since the DL were between 3 µg/l (penicillin with bioassay B) and 70 100 µg/l (streptomycin with bioassay Q). In addition, a logarithmic relative scale was chosen in terms of the MRL for each molecule (Log DL/MRL) because it allows to visualize the similarity or discrepancy between the DL and MRL of each antibiotic. In this diagram, circle 1 corresponds to Log DL/MRL = 0.1 values (the bioassay detects antibiotics below MRL levels), circle 2 corresponds to Log DL/MRL = 1 (limits equivalent to LMR detection), circle 3 indicates Log DL/MRL = 10 MRL and circle 4 indicates Log DL/MRL = 100 MRL (the bioassay does not detect the antibiotic).

It can be observed that Bioassay B (blue line) detected the eight beta-lactam antibiotics, as well as tylosin and neomycin; Bioassay T (red line) detected the three tetracyclines and neomycin; Bioassay M (green line) detected the three macrolides and neomycin; Bioassay Q (black line) detected only the three quinolones, and Bioassay QS (purple line) detected the four sulfonamides and the three quinolones. It is also noted that kanamycin and streptomycin were not detected by any of the five bioassays.

In summary, a simple visual analysis of the color of each bioassay easily allows to classify an antibiotic residue into

**Table 5** Interpretation of results of the microbiological system in microplates

Antibiotics	Bioassays				
	B	T	M	Q	QS
Free antibiotics	–	–	–	–	–
Beta-lactams	+ <sup>a</sup>	–	–	–	–
Tetracyclines	–	+ <sup>b</sup>	–	–	–
Macrolides	–	–	+ <sup>b</sup>	–	–
Quinolones	–	–	–	+	+
Sulfonamides	–	–	–	–	+

B: beta-lactams; T: tetracyclines; M: macrolides; Q: quinolones; QS: quinolones-sulfonamides.

<sup>a</sup> Neomycin or tylosin.

<sup>b</sup> Neomycin; + and –: positive and negative responses, respectively.

five categories. The simultaneous detection of quinolones (two bioassays), tylosin (two bioassays) and neomycin (three bioassays) provide a tool to ensure the presence of residues, thereby improving food security.

**Antibiotic classification by the five-bioassay system**

Table 5 summarizes the visual interpretation of the results obtained by the simultaneous analysis of the five bioassays. Responses are visually observed by the color change in each method. For Bioassays B, T and M, the yellow color indicates a negative result, whereas the black color indicates a positive result. For bioassays Q and QS, the pink color indicates a negative result, whereas blue indicates a positive result.

The presence of beta-lactams in milk was visualized by the persistence of the original black color in Bioassay B. Tetracycline residues were observed by the persistence of the black color in bioassay T, whereas macrolide residues were observed by the persistence of the black color in bioassay M. Sulfonamides were observed by the persistence of the original blue color in bioassay QS, while quinolone residues were evidenced by the persistence of the original blue color in bioassays Q and QS.

Table 5 shows that neomycin residues were evidenced by the persistence of the original colors of bioassays B, T and M, while tylosin residues were evidenced by the persistence of the original colors of bioassays B and M (Table 3). Finally, it should be noted that the milk samples that produced color change in the five bioassays were classified as negative samples either because they had no antibiotic or the antibiotics were below the detection limits of each bioassay.

## Conclusion

The microbiological system in microtiter plates consisting of Bioassays B (*G. stearothermophilus*), Bioassays T (*B. megaterium*), Bioassays M (*B. megaterium* and fucsidic acid), Bioassays Q (*B. licheniformis*) and Bioassays QS (*B. licheniformis* and trimetoprim) can detect a large amount of antibiotic residues in milk and allows to classify residues in beta-lactams, tetracyclines, macrolides, quinolones and sulfonamides. This system provides a simple dichotomous colorimetric response with visual interpretation and allows to analyze a large number of samples in a short period of incubation (4–5.5 h).

The simultaneous use of five bioassays represents a more efficient and rigorous control mechanism than the use of a single-bioassay microbiological system. Indeed, the routine implementation of this system in quality control microbiological laboratories may contribute to improving food security. Furthermore, in contrast to microbiological systems in Petri dishes, which should be used immediately after preparation, this microbiological system designed in a microtiter plate format heat-sealed with aluminized paper and containing sporulated bacteria allows the marketing, refrigerated transport and storage in a refrigerator at 4°C in the laboratory before being used. In the future, new bioassays using microorganisms with specific sensitivity to aminoglycosides should be developed to improve the detection profile and classification of antibiotic residues in milk.

## Conflict of interest

The authors declare that they have no conflicts of interest

## Funding

This research work was carried out as part of the CAI+D'16 Projects (PIC 50420150100113LI, Res. HCS 128/17, Universidad Nacional del Litoral, Santa Fe, Argentina).

## References

- Barra Caracciolo A, Topp E, Grenni P. Pharmaceuticals in the environment: biodegradation and effects on natural microbial communities. A review. *J Pharm Biomed Anal.* 2015;106:25–36.
- Beltrán MC, Berruga MI, Molina A, Althaus RL, Molina MP. Performance of the current microbial tests for screening antibiotic in sheep and goat milk. *Int Dairy J.* 2015;41:13–5.
- Berruga MI, Novés B, Molina MP, Román M, Molina A. Influence of cephalosporin on the coagulation time of yogurt made from ewe's milk. *Int J Dairy Tech.* 2008;61:372–8.
- Berruga MI, Beltrán MC, Noves B, Molina A, Molina MP. Effect of penicillins on the acidification of yogurt made from ewe's milk during the storage. In: International conference on antimicrobial research (ICAR2010). Science and technology against microbial pathogens. 2011. p. 145–9.
- Berry L, Read H, Walker L, Famula R. Clinical, histologic, and bacteriologic findings in dairy cows with digital dermatitis (footwarts) one month after topical treatment with lincomycin hydrochloride or oxytetracycline hydrochloride. *JAVMA.* 2010;237:555–60.
- Botsoglou N, Fletouris J. Drug residues in foods: pharmacology, food safety, and analysis. Series: food science and technology. New York, USA: Marcel Dekker, Inc.; 2001.
- Comunidad Económica Europea (CEE). Decisión 2002/657/CEE del Consejo del 12 de Agosto de 2002 por la que se aplica la Directiva 96/23/CE del consejo en cuanto al funcionamiento de los métodos analíticos y la interpretación de los resultados. *DOCE, L 73; 2002.* p. 30–1.
- Chanda R, Fincham R, Venter P. Review of the regulation of veterinary drugs and residues in South Africa. *Crit Rev Food Sci Nutr.* 2014;54:488–94. <http://dx.doi.org/10.1080/10408398.2011.588348>.
- Codex. Codex alimentarius commission, veterinary drug residues in food; 2016. Codex Veterinary Drug Residues in Food Online Database. Available at: <http://www.codexalimentarius.org/standards/vetdrugs/veterinary-drugs/en/> [accessed 04.01.16].
- European Union. Regulation (EC) No. 470/2009 of 6 lay 2009 laying down. Community procedures for the establishment of residue limits of pharmacologically active substances in foodstuffs of animal origin, repealing Council Regulation (EEC) No. 2377/90 and amending Directive 2001/82/EC of the European Parliament and of the Council and Regulation (EC) No. 726/2004 of the European Parliament and of the Council. *Off J.* 2009;L152:11–22.
- European Union. Regulation (EU) No. 37/2010 of 22 December 2009 on pharmacologically active substances and their classification regarding maximum residue limits in foodstuffs of animal origin. *Off J.* 2010;L15:1–72.
- Food and Drug Administration Center for Veterinary Medicine (FDA). MI-05-05: tolerance and/or safe levels of animal drug residues in milk; 2005. <http://www.fda.gov/Food/FoodSafety/ProductpecificInformation/MilkSafety/CodedMemoranda/MemorandaofInformation/ucm077350.htm> [accessed 14.11.11].
- Gaudin V, Maris P, Fuselier J, Ribouchon N, Cadieu P, Rault A. Validation of a microbiological method: the STAR protocol, a five-plate test for the screening of antibiotic residues in milk. *Food Addit Contam.* 2004;21:422–33.
- International Dairy Federation (IDF). Detecting antibiotics in milk – guidance on the application of screening and confirmatory methods in integrated dairy chain management. Bulletin No. 474. Brussels, Belgium: International Dairy Federation; 2014.



15. Jiménez Lozano E, Marqués I, Barrón D, Beltrán JL, Barbosa J. Determination of pKa values of quinolones from mobility and spectroscopic data obtained by capillary electrophoresis and a diode array detector. *Anal Chim Acta*. 2002;464:37–45.
16. Kaufmann TV, Westermann S, Drillich M, Plöntzke J, Heuwieser W. Systemic antibiotic treatment of clinical endometritis in dairy cows with ceftiofur or two doses of cloprostenol in a 14-d interval. *Anim Reprod Sci*. 2010;121:55–62.
17. Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clin Microbiol Rev*. 2011;24:718–33.
18. Nagel OG, Molina MP, Althaus RL. Optimization of bioassay for tetracycline detection in milk by means of chemometric techniques. *Lett Appl Microbiol*. 2011;52:245–52.
19. Nagel OG, Molina MP, Althaus RL. Use of chemometric techniques to design a microbiological method for sulfamide detection in milk. *Czech J Food Sci*. 2013;31:627–32.
20. Nagel O, Molina M, Althaus R. Microbiological system in microtitre plates for detection and classification of antibiotic residues in milk. *Int Dairy J*. 2013;32:150–5.
21. Nouws J, Van Egmond H, Shulders I, Loeffen G, Schouten J, Stegeman H. A microbiological assay system for assessment of raw milk exceeding EU maximum residue level. *Int Dairy J*. 1999;9:85–90.
22. Nölvak H, Truu M, Tiirik K, Oopkaup K, Sildvee T, Kaasik A, Mander T, Truu J. Dynamics of antibiotic resistance genes and their relationships with system treatment efficiency in a horizontal subsurface flow constructed wetland. *Sci Total Environ*. 2013;461–462:636–44.
23. Pikkemaat MG, Rapallini ML, Zuidema T, Elferink JW, Oostra-van Dijk S, Driessen-van Lankveld WD. Screening methods for the detection of antibiotic residues in slaughter animals: comparison of the European Union four-plate test, the Nouws antibiotic test and the Premi<sup>®</sup> Test (applied to muscle and kidney). *Food Add Contam*. 2011;28:26–34.
24. Stanton AL, Kelton DF, LeBlanc SJ, Millman ST, Wormuth J, Dingwell RT, Leslie KE. The effect of treatment with long-acting antibiotic at postweaning movement on respiratory disease and on growth in commercial dairy calves. *J Dairy Sci*. 2010;93:574–81.
25. Statgraphics Centurion, X.V. Version 15.2.05. Edición Multilingüe. StatPoint. Inc.; 2007.
26. Sawant A, Sordillo L, Jayarao B. A survey on antibiotic usage in dairy herds in Pennsylvania. *J Dairy Sci*. 2005;88:2991–9.
27. Tsai C, Kondo F. Improved agar diffusion method for detecting residual antimicrobial agents. *J Food Prot*. 2001;64:361–6.
28. Tumini M, Nagel O, Althaus R. Microbiological bioassay using *Bacillus pumilus* to detect tetracycline in milk. *J Dairy Res*. 2015;82:248–55.
29. Tumini M, Nagel O, Molina M, Althaus R. Novel bioassay using *Bacillus megaterium* to detect tetracycline in milk. *Rev Argent Microbiol*. 2016;48:143–6.
30. Tumini M, Nagel O, Molina M, Althaus R. Microbiological method using *Bacillus megaterium* with fusidic acid for detection of macrolides in milk. *Czech J Food Sci*. 2016;34:9–15.
31. Tumini M, Nagel O, Molina M, Althaus R. Microbiological assay with *Bacillus licheniformis* for the easy detection of quinolones in milk. *Int Dairy J*. 2016;64:9–14.