



BRIEF REPORT

## ***Brucella canis* Group 2 isolated in Argentina**



Eduardo Jorge Boeri<sup>a,\*</sup>, María Julia Madariaga<sup>a</sup>, María Luz Dominguez<sup>a</sup>,  
María Luisa Teijeiro<sup>a</sup>, Natalia Mercedes Fernandez<sup>a</sup>, Sebastián Alejandro Elena<sup>b</sup>,  
Marcos David Trangoni<sup>c</sup>

<sup>a</sup> Instituto de Zoonosis Luis Pasteur, Av. Diaz Velez 4821 (1405), Ciudad Autónoma de Buenos Aires, Argentina

<sup>b</sup> Laboratorio de Referencia de la OIE para Brucellosis, Dirección General de Laboratorio y Control Técnico (DiLab), Servicio Nacional de Sanidad y Calidad Agroalimentaria (SENASA), Martínez, 1640 Buenos Aires, Argentina

<sup>c</sup> Laboratorio de Brucella, Campylobacter y Microbiota, Instituto de Biotecnología-IABIMO, INTA-CONICET, CICVyA-CNIA INTA Hurlingham, Argentina

Received 12 September 2019; accepted 26 June 2020

Available online 8 November 2020

### KEYWORDS

MLVA;  
Diversity;  
Deletion;  
BMEI1435;  
*Brucella canis* Group 2

### PALABRAS CLAVE

MLVA;  
Diversidad;  
Delección;  
BMEI1435;  
*Brucella canis* Grupo 2

**Abstract** The aim of this study was to estimate the diversity and prevalence of both groups of *Brucella canis* 1 and 2 with and without deletion respectively in different areas of Argentina. A total of 104 bacterial cultures were typed as *B. canis* strains using the classical biotyping method. Two PCR assays were performed to confirm that all isolates were *B. canis* and not *Brucella suis*. The differentiation between groups 1 and 2 was achieved using another PCR assay and the diversity of *B. canis* isolates was assessed with four MLVA\_16 markers. All strains belonged to Group 2. Bruce 09 marker (MLVA\_16 assay) showed the greatest diversity. Only Group 2 of *B. canis* was identified among the strains evaluated. The markers chosen from the MLVA\_16 allowed us to detect genetic diversity among the strains of *B. canis* studied.

© 2020 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### *Brucella canis* Grupo 2 aislado en Argentina

**Resumen** El objetivo de este trabajo fue estimar la diversidad y la prevalencia de ambos grupos de *Brucella canis* 1 y 2 (con y sin delección, respectivamente) en diferentes áreas de Argentina. Un total de 104 cultivos bacterianos se tipificaron como cepas de *B. canis* usando biotipado clásico. Se realizaron dos ensayos de PCR para confirmar que todos los aislamientos eran *B. canis* y no *Brucella suis*. La diferenciación entre los grupos 1 y 2 se logró con otro ensayo de PCR, y la diversidad entre las cepas de *B. canis* se obtuvo mediante el empleo de cuatro marcadores del ensayo de MLVA\_16. Todas las cepas pertenecieron al grupo 2. El marcador Bruce

\* Corresponding author.

E-mail address: [eduardoboeri@gmail.com](mailto:eduardoboeri@gmail.com) (E.J. Boeri).

09 (ensayo MVLA-16) mostró la mayor diversidad. Sólo se halló el Grupo 2 de *B. canis* entre las cepas estudiadas. Los marcadores del MLVA\_16 permitieron detectar la presencia de diversidad genética entre las cepas de *B. canis* analizadas.  
 © 2020 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

*Brucella canis* is an etiological agent of canine brucellosis that has been found in many countries, such as the US, China and Mexico as well as in countries of South America, Asia and Europe. This zoonotic disease causes abortion in female dogs and epididymitis in male dogs as the main symptom. In addition, this disease causes important losses in breeding kennels because of reproductive failures<sup>5</sup>.

Bacterial isolation is the “gold standard” for the definitive diagnosis, despite the low sensitivity of this procedure regarding clinical samples<sup>2</sup>. In addition, the global scientific community uses the process of phenotypic characterization from isolates to distinguish the different species of the genus *Brucella*<sup>1</sup>.

Researchers have employed the polymerase chain reaction assay (PCR) in many studies for the molecular typing of *Brucella* species and biovars from bacteriological cultures<sup>10,14</sup>. For example, the Bruce-ladder assay (a multiplex PCR assay) is the only analysis accepted by the World Organization for Animal Health (OIE) for identification and typing *Brucella* species<sup>13</sup>. However, some *B. canis* strains can be identified erroneously as *B. suis*<sup>10</sup>. The researchers that reported this finding identified two groups of *B. canis* based on the presence or absence of a region of the BMEI1435 polysaccharide deacetylase gene in *B. canis*. Indeed, Group 1 presents a deletion (PCR amplifies a 607-bp fragment), whereas Group 2 lacks the deletion (PCR amplifies a 1674-bp fragment). López Goñi et al. replaced two primers of the PCR original protocol and obtained the Bruce-ladder v2.0 test<sup>14</sup>. This new test allows the differentiation of strains of *B. canis*, five biovars of *B. suis* and *B. neotomae*. In parallel, Kang et al.<sup>9</sup> also developed new primers from a specific region to avoid the problem of the original Bruce-ladder.

Another research group found that only Group 2 is present in Medellín city<sup>15</sup>. More recently, another study reported both groups in China<sup>4</sup>.

Multiple Locus Variable-number Tandem Repeat Analysis (MLVA) is a method that has been performed for the typing of bacterial species including *Mycobacterium tuberculosis*, *Bacillus anthracis* and others. A set of eight microsatellite loci has been proposed because they are highly discriminatory and efficient for distinguishing strains within a local outbreak. Although we cannot correctly predict the biovar or even the species of an isolate, this method is a useful tool especially in outbreaks. To date, different MLVA assays have been developed for typing *Brucella* strains with different kind of outcomes and utilities<sup>12</sup>.

To the best of our knowledge, in Argentina, there are no studies published on groups of *B. canis* isolates or any publication on *B. canis* MLVA studies. With this in mind, the aim

of this study was to estimate the prevalence of both groups of *B. canis* in different areas of Argentina from microbiological isolates of clinical dog samples. Other objectives were to compare the performance of two PCR assays in discriminating *B. canis* and *B. suis* strains and, finally, to estimate the diversity of *B. canis* using four markers of the MLVA\_16 assay.

The analyzed samples consisted of 104 *B. canis* isolates from clinical dog samples obtained between August 2009 and December 2018. The samples were collected from the Autonomous City of Buenos Aires and different areas of Argentina. Strains were typed following the recommendations described elsewhere<sup>1</sup>. Molecular typing confirmation of all *B. canis* strains and not *B. suis* was performed using two PCR assays<sup>9,14</sup>. DNA was extracted using a commercial kit, ROCHE High Pure PCR Template Preparation Kit (Roche Diagnostics, GmbH Roche Applied Science, Germany), following the manufacturer's instructions. A concentration between 50 and 70 ng/μl was used for further analyses after measuring DNA with ND-1000 NanoDrop. As a negative control of DNA extraction, we used ultrapure water. Two PCR assays (PCR1 and PCR2) were carried out to differentiate *B. canis* and *B. suis*. PCR1 and PCR2 were performed according to Lopez Goñi et al.<sup>14</sup> and Kang et al.<sup>9</sup> respectively. For both PCRs, *B. canis* strain RM6/66 and *B. suis* biovar 1 were used as positive controls. In addition, we used ultrapure water as a negative control. A third PCR (PCR3) was conducted according to Koylass et al.<sup>10</sup>. *B. canis* strain RM6/66 (Group 1) and *Brucella melitensis* biovar 1 (Group 2) were used as positive controls, respectively. All the products from PCR1, PCR2 and PCR3 were analyzed by electrophoresis in 1.5% agarose gel stained with ethidium bromide (0.5 μg/ml) and UV light visualization.

The markers used for the MLVA assay in this study were Bruce 04, Bruce 07, Bruce 09 and Bruce 16<sup>12</sup>. The selection of these markers was based on their greatest diversity among species of *B. canis*<sup>4,8</sup> (Table S1). A total of 15 strains were selected from various areas of Argentina: San Miguel de Tucumán (Province of Tucumán); Rio Cuarto (Province of Córdoba); Rosario (Province of Santa Fe); Río Grande (Province of Tierra del Fuego); General Pico (Province of La Pampa); Corrientes (Province of Corrientes); Ituzaingó, Tortuguitas, Paso del Rey, Del Viso (two strains), Villa Ballester, Banfield, Verónica (Province of Buenos Aires) and Villa Lugano (Autonomous City of Buenos Aires). This allowed an approximate representation of the different alleles (number of repeats) present in different areas of the country.

DNA amplification was conducted in tubes with a final volume of 25 μl per reaction. Each reaction contained DNA

(50 µg), each primer (forward/reverse; 0.5 µM), each of the four deoxynucleotide triphosphates (dNTPs; 200 µM), 1X buffer with magnesium chloride ( $MgCl_2$ , final concentration of 1.5 mM) (PROMEGA 5X Green GoTaq® Flexi Buffer Migration Pattern) and 1 U of Taq DNA Polymerase. *B. canis* strain RM6/66 and *B. melitensis* 16M were used as positive controls. Ultrapure water was used as negative control. The thermocycling conditions were as follows: a first denaturation at 94 °C for 3 min and 30 cycles of 94 °C for 30 s, 60 °C for 30 s and 72 °C for 50 s with a final extension of 72 °C for 5 min. Bionumerics software version 3.5 (Applied Maths, St-Martens-Latem, Belgium) was used to estimate the size of the PCR bands of the agarose gels. The quantification of the number of repeats of each marker was performed following the protocol described by Le Flèche et al.<sup>12</sup> The dendrogram was generated through the unweighted pair group method following arithmetic averages (UPGMA) and using the PAST software<sup>6</sup>. The Hunter-Gaston Diversity Index (HGDI)<sup>7</sup> was calculated using Epiccompare software version 1.0 ([www.ridom.de/epiccompare](http://www.ridom.de/epiccompare)) to elucidate the discriminatory power of the genotyping methods. Number 0 and 1 stand for "without diversity" and "extreme diversity", respectively. These results reflect the total of detected alleles. The PCR products were analyzed in 2% agarose gel in Tris acetate and EDTA (ethylenediaminetetraacetic acid) 1X buffer (TAE) with the addition of 0.5 µg/ml of ethidium bromide. An Applied Biosystems Veriti™ Thermal Cycler was used for the PCR reactions.

All 104 strain isolates were typed as *B. canis* according to the classical biotyping method. In this study, we evaluated two PCR assays for molecular typing of the evaluated bacteria that had been confirmed as *B. canis* strains by the classical method.

The two PCRs used to assess the species (PCR1 and PCR2) confirmed that all strains were *B. canis* (Table 1). Thus, both techniques yielded consistent results. PCR1 shows the molecular patterns between strains of *B. canis* and *B. suis*. PCR2 displays a 776-bp differential amplicon between species, i.e. *B. suis* contains the amplicon, whereas *B. canis* lacks it. In this study, all the evaluated strains were *B. canis* Group 2 (without deletion; 100% (CI 95% 0.95–1.00); Table 1, Fig. 1).

Subsequently, we performed an evaluation of the strains by MLVA\_16 by using four markers. For this purpose, we analyzed 15 strains from different regions of Argentina. The overall Diversity Index (HDGI) of the technique was 1.0 (CI 95% 1.0–1.0). The HDGI coefficient ranged from 0.705 to 0.895. Bruce 09 marker displayed the greatest diversity (0.895 CI 95% 0.816–0.974), followed by Bruce 16 (0.876 CI 95% 0.803–0.949), Bruce 07 (0.848 CI 95% 0.805–0.890) and Bruce 04 (0.705 CI 95% 0.550–0.859) (Table S2). Additionally, we identified 15 different molecular patterns (Bruce 04, Bruce 07, Bruce 09, Bruce16 respectively: 4-8-6-9, 6-5-8-5, 6-5-11-8, 6-5-11-9, 6-5-6-5, 6-6-8-9, 6-8-10-11, 6-9-8-8, 7-7-10-5, 8-6-7-9, 8-6-8-8, 8-8-4-12, 8-9-9-11, 8-9-5-5, 11-7-10-7) (Table S3). The molecular pattern detected in Del Viso (6-5-8-5) and the ones found in Banfield (6-8-10-11) were similar to those described in the city of Beijing and the Autonomous Region of Guanxi, Republic of China, respectively<sup>4</sup>. In addition, the molecular pattern found in La Pampa (6-6-8-9) was similar to those in Paju city, Gyeonggi Province, South Korea<sup>8</sup>.

In this study, we evaluated various *B. canis* strains from different regions of Argentina to assess the groups of strains existing in Argentina. All strains were identified using the classical biotyping method and molecular techniques. Among all the strains analyzed in this study, we only detected *B. canis* type 2 (without deletion of the BMEI1435 gene). This finding is consistent with a previous report from Colombia<sup>15</sup>. Similarly, most of the isolates analyzed in China belong to this type, although three isolates exhibited a deletion<sup>4</sup>. By contrast, Koylass et al., found both groups circulating in equal proportion in samples evaluated from Europe, South America and the United States. Specifically, the strains with the deletion came from Peru, Germany and the United States<sup>10</sup>. The pathogenicity of the strains with and without this deletion has not been studied yet. These data could trigger future research regarding the characteristics of the BMEI1435 gene. This gene has a hydrolase function and participates in the metabolism of carbohydrates. The variation of this gene may have an impact on virulence in the host, but this is still unclear. A recent study compared different *Brucella* genes and their virulence but did not include BMEI1435 in the analysis, perhaps due to the lack of consideration regarding pathogenicity at the date of the cited study<sup>3</sup>.

In the present study, we used four markers presented at panel 2 of the MLVA\_16 (Bruce 04, Bruce 07, Bruce 09 and Bruce 16) and based on the polymorphism found by other researchers<sup>4,8</sup>. We did not use the complete panel of MLVA\_16 and therefore we believe we cannot speak of "genotypes". Therefore, we defined them as molecular patterns instead. As a preliminary study, we used 15 strains from different regions of the country, which provided data on the diversity among circulating *B. canis* strains in specific areas of Argentina. It should be noted that the remaining markers of the full panel of the MLVA\_16 in the research of Di et al.<sup>4</sup> and Kang et al.<sup>8</sup> showed similar repeat numbers. These data suggest high homology of *B. canis* for these gene regions. Thus, the four markers used in the present study are suitable for evaluating *B. canis* genetic diversity. However, the use of the complete panel is very useful for other *Brucella* species such as *B. melitensis*, *B. suis* and *B. abortus*<sup>11</sup>.

The results from the dendrogram (Fig. 2) suggest great similarity between the strains of Villa Ballester and Del Viso (Province of Buenos Aires) and this result could be explained by the closeness of the locations (36 km). Although other strains, such as the strains of the city of Rosario (Province of Santa Fe) and Ituzaingó (Province of Buenos Aires) are somewhat further away (300 km), they also showed high similarity. In addition, the similarity between the strains of Rio Grande (Province of Tierra del Fuego) and the town of Banfield (Province of Buenos Aires), which are separated by 2800 km, is of particular interest. These findings suggest that dogs may have circulated from one location to another. This is quite usual as people often migrate with their dogs.

Finally, as a first preliminary study in Argentina, our results indicate the existence of a genetic diversity among circulating *B. canis* strains. In addition, the development of the complete MLVA\_16 panel should be done in the near future in order to broaden the knowledge of the circulating genotypes in Argentina.

Our results suggest that the only group of *B. canis* circulating in Argentina is Group 2 (without BMEI1435 gene


**Table 1** Description of *B. canis* strains and techniques used for their characterization and differentiation in groups 1 and 2.

Order	Isolation	Biotype	Bruce-ladder v2.0	Kang et al., 766 pb	Amplicon	Deletion	Year	Place
1	70150	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2014	Recoleta
2	70152	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2014	Recoleta
3	76531	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2015	P. del Rey <sup>a</sup>
4	55420	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	V. Lugano <sup>a</sup>
5	87447	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	La Pampa <sup>a</sup>
6	23653	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
7	91224	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Villa Crespo
8	23671	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
10	27269	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Almagro
11	23502	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2008	Caballito
12	47937	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2007	Almagro
13	25962	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2007	P. Chacabuco
14	51545	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	V. Ballester <sup>a</sup>
16	92151	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Barracas
18	47247	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
19	51080	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Mataderos
20	100887	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2017	Belgrano
21	48502	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	P. Madero
22	35241	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2011	Villa Lugano
24	1865	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2015	R. Grande
25	1789	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2014	Rosario <sup>a</sup>
26	1863	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	R. Grande
27	1864	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	R. Grande <sup>a</sup>
29	1790	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Belgrano
31	C28	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Pompeya
32	C32	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso <sup>a</sup>
33	C36	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
34	C40	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
35	C42	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso <sup>a</sup>
36	C1	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2015	Barracas
37	C5	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Caballito
39	C9	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Verónica <sup>a</sup>
40	C14	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Tucumán
41	C25	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Tortuguitas <sup>a</sup>
42	C26	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
43	C27	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Tortuguitas
44	C31	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
45	C33	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
46	C34	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
47	C37	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
48	C38	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
49	C39	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
50	C41	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
51	C43	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Del Viso
55	C10	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Castelar
57	C15	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Tucumán <sup>a</sup>
58	C16	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Ramos Mejía
61	C21	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	San Martín
62	C22	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Vicente López
63	C24	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2010	Olivos
67	C13	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
70	49080	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Caballito
71	74873	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Almagro
72	35241	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2011	Villa Lugano
73	47214	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2008	Caballito
74	47937	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2007	Almagro
75	74532	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Barracas
76	70218	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2015	Villa Urquiza
77	77794	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2016	Belgrano


Table 1 (Continued)

Order	Isolation	Biotype	Bruce-ladder v2.0	Kang et al., 766 pb	Amplicon	Deletion	Year	Place
78	47246	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
79	57327	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Barracas
80	79207	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2017	Banfield <sup>†</sup>
81	79341	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2017	Banfield
82	79894	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2017	V.del Parque
83	47131	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	P. del Rey
84	47224	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2009	Caballito
85	C48	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2011	Caballito
86	C49	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Caballito
87	C50	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Caballito
88	C51	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2011	Caballito
89	C52	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	Caballito
90	C53	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2014	Caballito
91	C54	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2011	Caballito
92	C55	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Caballito
93	C57	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	Caballito
94	C58	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2012	Caballito
95	C61	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	Rio cuarto <sup>†</sup>
96	C64	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2014	Verónica
97	C70	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2013	Barracas
98	82175	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	Ituzaingó
99	82122	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	P. Patricios
100	81296	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	P. Patricios
101	83961	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	Balvanera
102	Cepa 12	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	Corrientes <sup>†</sup>
103	82543	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	1674	No	2018	Ituzaingó <sup>†</sup>
104	RM6/66	<i>B. canis</i>	<i>B. canis</i>	<i>B. canis</i>	607	YES	2018	Reference

<sup>†</sup>: strains analyzed by MLVA.



**Figure 1** Electrophoresis on agarose gel of PCR products showing representative strains of Group 2. Lane 1: Negative control of PCR. Lane 2: Negative control of DNA extraction. Lane 3–7: profile of five strains evaluated. Lane 8: positive control of Group 2 (1674 bp). Line 9: positive control of Group 1 (607 bp). Line 10: molecular marker Dangsheng Biotech 100 bp DNA ladder plus.



**Figure 2** Dendrogram based on the different copy numbers of the tandem repeats of the 25 alleles found in the 15 strains studied by MLVA. The Euclidean similarity measure was used for the construction of the phylogenetic tree.

deletion). The two PCRs used for molecular typing yielded consistent results. Therefore, both are useful to discriminate between *B. canis* and *B. suis*. However, PCR1 should be used to evaluate biovars of *B. suis*. The use of the four markers chosen from the MLVA\_16 allowed us to identify the genetic diversity among the strains of *B. canis* circulating in Argentina. Thus, as a preliminary test, MLVA\_16 is useful for the genetic discrimination of this bacterium.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgment

We thank Dr. Gabriela Escobar for the provision of strains, Dr. Zumarraga Martín for processing agarose gels and Dr. Julia Sabio y García for critical reading of the manuscript. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ram.2020.06.012.

## References

1. Alton G, Jones S L., Angus S R., Verger S J. Bacteriological methods. In: Institut National de la Recherche Agronomique, editor. Techniques for the Brucellosis laboratory. Paris, France: INRA; 1988. p. 13–60.
2. Boeri E, Wanke M, Madariaga M, Teijeiro M, Trangoni M. Comparison of four polymerase chain reaction assays for the detection of *Brucella* spp. in clinical samples from dogs. Vet World. 2018;11:201–8.
3. Brambila-Tapia A, Armenta-Medina D, Rivera-Gomez N, Perez-Rueda E. Main functions and taxonomic distribution of virulence genes in *Brucella melitensis* 16M. PLoS One. 2014;9: e100349.
4. Di D, Cui B, Wang H, Zhao H, Piao D, Tian L, Tian G, Kang J, Mao X, Zhang X, Du P, Zhu L, Zhao Z, Mao L, Yao W, Guan P, Fan W, Jiang H. Genetic polymorphism characteristics of *Brucella canis* isolated in China. PLoS One. 2014;9:e84862.
5. Greene CE, Carmichael LE. Canine Brucellosis. In: Sykes J, Greene C, editors. Infectious diseases of the dog and cat. 4th ed. Saunders Elsevier; 2012. p. 398–411.
6. Hammer Ø, Harper D, Ryan P. Past: Paleontological statistics software package for education and data analysis, vol. 4, issue 1, art. 4. PalaeoElectronica; 2001. p. 9, 178 kb [online]. [http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)
7. Hunter P, Gaston M. Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. J Clin Microbiol. 1988;26:2465–6.
8. Kang S, Heo E, Cho D, Kim J, Kim Ji, Jung S, Her M. Genetic comparison of *Brucella canis* isolates by the MLVA assay in South Korea. J Vet Med Sci. 2011;73:779–86.
9. Kang S, Her M, Kim J, Kim Ji, Ko K, Ha Y, Jung S. Advanced multiplex PCR assay for differentiation of *Brucella* species. Appl Environ Microbiol. 2011;77:6726–8.
10. Koylass M, King A, Edwards-Smallbone J, Gopaul K, Perrett L, Whatmore A. Comparative performance of SNP typing and "Bruce-ladder" in the discrimination of *Brucella suis* and *Brucella canis*. Vet Microbiol. 2009;142:450–4.
11. Ledwaba M, Gomo C, Lekota K, Le Flèche P, Hassim A, Vergnaud G, Heerden H. Molecular characterization of *Brucella* species from Zimbabwe. PLoS Negl Trop Dis. 2019;13:e0007311.
12. Le Flèche P, Jacques I, Grayon M, Al Dahouk S, Bouchon P, Denoeud F, Nöckler K, Neubauer H, Guilloteau L, Vergnaud G. Evaluation and selection of tandem repeat loci for a *Brucella* MLVA typing assay. BMC Microbiol. 2006;6:9.
13. López-Goñi I, García-Yoldi D, Marín C, de Miguel M, Muñoz P, Blasco J, Jacques I, Grayon M, Cloeckaert A, Ferreira A, Cardoso R, Corrêa de Sá M, Walravens K, Albert D, Garin-Bastuji B. Evaluation of a multiplex PCR assay (Bruce-ladder) for molecular typing of all *Brucella* species, including the vaccine strains. J Clin Microbiol. 2008;46:3484–7.
14. Lopez-Goñi I, García-Yoldi D, Marín C, de Miguel M, Barquero-Calvo E, Guzman-Verri C, Albert D, Garin-Bastuji B. New Bruce-ladder multiplex PCR assay for the biovar typing of *Brucella suis* and the discrimination of *Brucella suis* and *Brucella canis*. Vet Microbiol. 2011;154:152–5.
15. Ortiz L, Muskus C, Sánchez M, Olivera M. Identification of *Brucella canis* Group 2 in Colombian kennels. Rev Colomb Cienc Pecu. 2012;25:615–9.