



Original article

A new endophytic species of *Microthecium* (Melanosporales, Sordariomycetes, Pezizomycotina, Ascomycota) from Mallorca (Balearic Islands, Spain)



Alberto M. Stchigel^a, José F. Cano-Lira^{a,*}, Ángel Pintos-Amengual^b

^a Universitat Rovira i Virgili (URV), Mycology Unit, Medical School, Sant Llorenç 21, 43201 Reus, Tarragona, Spain

^b Interdisciplinary Ecology Group University of Balearic Islands, Universitat de les Illes Balears, Ctra Valldemossa Km 7.5, Palma de Mallorca, Spain

ARTICLE INFO

Article history:

Received 17 March 2023

Accepted 23 October 2023

Available online 4 January 2024

Keywords:

Ascomycetous fungi

Crataegus

Endophytic fungi

Mediterranean

ABSTRACT

Background: The genus *Microthecium* contains 31 species worldwide distributed. Most of them are saprobic on soil and plant debris, but a few have been reported as mycoparasites on hypocrealean fungi. By contrast, this genus has never been reported as phytopathogenic, nor endophytic.

Aims: To isolate and identify endophytic fungi from Mediterranean herbaceous plants and trees in order to contribute to the knowledge of the hosts and their geographical location. The present work has been focused on the study of endophytic fungi of hawthorn (*Crataegus monogyna*).

Methods: The following steps were taken: i, isolation of the fungal strain from living stems of *C. monogyna*; ii, cultural and micro-morphological study, and iii, sequence comparison of different genetic markers by BLAST search with sequences deposited in GenBank.

Results: At the present work we describe a new species of the genus, *Microthecium pleomorphosporum*, isolated from living stems of *C. monogyna* in Mallorca (Balearic Islands, Spain). This fungus is characterized by the production of non-ostiolate perithecia and two sorts of ascospores (some smooth-walled, others delicately reticulated) bearing a germ pore at each end which are frequently ornamented by a surrounding donut-like structures, and a phialidic asexual morph and bulbils. The morphologically closest related species is *Microthecium tenuissimum*, which has bigger ascospores and lacks asexual reproduction. Phylogenetically, *M. pleomorphosporum* is close-related to other species of the genus, although no genetic marker that discriminates this new species from other phylogenetically closer ones could be elucidated as a gold standard.

Conclusions: *M. pleomorphosporum*, order Melanosporales, is reported here as the first endophytic species of *C. monogyna*.

© 2023 The Author(s). Published by Elsevier España, S.L.U. on behalf of Asociación Española de Micología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Una nueva especie endofítica de *Microthecium* (Melanosporales, Sordariomycetes, Pezizomycotina, Ascomycota) de Mallorca (Islas Baleares, España)

RESUMEN

Antecedentes: El género *Microthecium* incluye 31 especies de distribución cosmopolita. La gran mayoría de ellas son saprobias en el suelo y se encuentran sobre detritos vegetales, y tan solo unas pocas se han descrito como micoparásitas de hongos hipocreales. Por el contrario, nunca se había descrito a este género como fitopatógeno ni endofítico.

Objetivos: Aislar e identificar hongos endófitos de plantas herbáceas y árboles mediterráneos con el fin de contribuir al conocimiento de los hospederos y su situación geográfica. El presente trabajo se ha centrado en el estudio de los hongos endófitos del espino albar (*Crataegus monogyna*).

Palabras clave:

Hongos ascomicetos

Crataegus

Hongos endofíticos

Mediterráneo

* Corresponding author.

E-mail address: jose.cano@urv.cat (J.F. Cano-Lira).

Métodos: Se incluyeron diferentes procesos: (1) aislamiento de la cepa fúngica a partir de tallos vivos de *Crataegus monogyna*; (2) estudio cultural y micromorfológico y (3) comparación de las secuencias nucleotídicas de diferentes marcadores filogenéticamente informativos con secuencias de taxones conocidos depositadas en el *GenBank* mediante búsqueda BLAST.

Resultados: En el presente trabajo describimos una nueva especie para el género, *Microthecium pleomorphosporum*, aislada de tallos vivos de *C. monogyna* en Mallorca (Islas Baleares, España). Este hongo se caracteriza por la producción de peritecios no ostiolados y de dos tipos de ascosporas (unas con paredes lisas y otras con paredes delicadamente reticuladas), con un poro germinativo en cada extremo frecuentemente ornamentado por estructuras circundantes en forma de rosquilla, y un estado asexual que produce conidios fialídicos y bulbillos. La especie morfológicamente más cercana es *Microthecium tenuissimum*, que tiene ascosporas más grandes y carece de multiplicación asexual. Filogenéticamente, *Microthecium pleomorphosporum* está estrechamente emparentada con otras especies del género y no se ha podido establecer ningún marcador genético de referencia (*gold standard*) para discriminar esta nueva especie de otras filogenéticamente más próximas.

Conclusiones: *Microthecium pleomorphosporum*, del orden de los Melanosporales, puede ser la primera especie endofítica de *C. monogyna*.

© 2023 El Autor(s). Publicado por Elsevier España, S.L.U. en nombre de Asociación Española de Micología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The genus *Microthecium* Corda, recently resurrected by Marin-Felix et al.,⁹ contains species mostly producing yellowish-orange, orange-brown or reddish globose ascomata, brown, citriform to plataniform ascospores with smooth, reticulate, pitted or wrinkled walls, bearing a terminal apiculate or depressed germ pore at each end, and almost producing an asexual phialidic morph and bulbils *in vitro*. Currently, there are 31 accepted species in that genus,⁹ usually found on dung, plant debris, soil, monocotyledonous bulbs and Pinophyta, frequently as parasites of other fungal taxa² and never been previously described as endophytes.

During a survey on endophytic fungi of the Mediterranean flora, we isolated in pure culture a fungus morphologically similar to the genus *Microthecium*. After a molecular study and morphological comparison with the accepted species of the genus, we suggest erecting the new species *M. pleomorphosporum*.

Materials and methods

Fungal isolation

A fungus belonging to the order Melanosporales was isolated from living healthy stems of *Crataegus monogyna* Jacq. sampled in Menut (Mallorca, Balearic Islands, Spain). *C. monogyna* is known as “hawthorn” in English, receiving the common names of *espino albar* and *majuelo* in Spain, *arç blanc* in Catalonia and *cirerer de pastor* in the Balearic Islands. It is a spiny deciduous tree distributed mainly in the Mediterranean region, growing on all sorts of soils, from almost sea level to 2200 masl, 4–8 m in height, with a trunk not always unique, blackish-brown to grayish-brown bark. The dark greenish leaves, pedunculated 2–4 cm long, are lobed, denticulate, and alternate along the stem; the white flowers are grouped in corymbs, later producing red pommel fruits with a single seed.

The fungus was isolated from the stems after surface sterilization according to Crous et al.³ Briefly, the stems were cut in fragments of 2–4 cm long, submerged in 70% ethanol for 1 min, treated with NaOCl 2% for 2 min and, finally, with ethanol 70% for 1 min. After that, the sterilized stems were rinsed twice with sterile water, plated onto malt extract agar (MEA; Difco Inc., Detroit, USA) into 90 mm diameter disposable Petri dishes, and incubated at 25 °C. After two weeks some yellowish ascomata were observed under the stereomicroscope, and some of them were transferred, using a sterile needle, to 55 mm diam. Petri dishes, ones containing oatmeal agar (OA; oatmeal flakes, 30 g; agar-agar, 20 g; distilled water, 1 L) and others with potato dextrose agar (PDA; Pronadisa, Madrid, Spain). The Petri dishes were incubated at 15, 25 and 35 °C.

Morphological characterization

For cultural characterization, the isolate was grown for up to 30 days on OA, potato carrot agar (PCA; grated potatoes, 20 g; grated carrot, 20 g; agar-agar, 20 g; L-chloramphenicol, 100 mg; distilled water, 1 L), and PDA at 5, 10, 15, 20, 25, 30, 35 and 40 °C.⁹ Color notations in parentheses are from Kornerup and Wanscher.⁷ Vegetative and reproductive structures were examined under an Olympus BH-1 bright field microscope by direct mounting in lactic acid and water of the ascomata and/or microcultures grown on OA and PDA. Pictures were obtained with a Zeiss Axio Imager M1 brightfield microscope.

Molecular study

The DNA of our fungal isolate was extracted and purified directly from the colonies by means of the Fast DNA Kit protocol (MP Biomedicals, Solon, Ohio). The amplification of the internal transcribed spacer region (ITS) of the nuclear rDNA was carried out according to Magaña-Dueñas et al.⁸; genes' fragments encoding actin (*act*), translation elongation factor 1- α (*tef1*), RNA polymerase II subunit 2 (*rpb2*) and beta-tubulin (*tub2*) were also amplified according to Voigt and Wöstemeyer¹⁴ (*act*), Houbraken et al.⁵ (*tef1*), Sung et al.¹³ (*rpb2*) and Woudenberg et al.¹⁵ (*tub2*). BigDye Terminator 3.1 cycle sequencing kit (Applied Biosystems Inc., Foster City, California) was used to sequence both strands with a combination of the same primers used in the amplification. PCR products were purified and sequenced at MacroGen Europe (Amsterdam, The Netherlands) with a 3730XL DNA analyzer (Applied Biosystems), and the consensus sequences were obtained using SeqMan (version 7.0.0; DNASTAR, Madison, WI, USA). The sequences obtained were compared with other fungal sequences deposited at the National Center for Biotechnology Information (NCBI) database using the Basic Local Search Tool (BLAST; <https://blast.ncbi.nlm.nih.gov/Blast.cgi>). A maximum level of identity (MLI) $\geq 98\%$ was used for species-level identification.

Results

Molecular study

Based on the BLAST search in NCBI's GenBank nucleotide database using a fragment of the ITS nucleotide sequence of our strain FMR 18380 (272 bp, accession number OX383416.1), it was displayed a percentage identity greater than 98.78% for




Fig. 1. *Microthecium pleomorphosporum* (CBS 148652). (A) Colony after 14 days at 25 °C on PCA (front and reverse). (B) Colony after 14 days at 25 °C on PDA (front and reverse). (C and D) Translucent ascomata showing the peridial layers and eight ascospores within the asci. (E) Broad ascoma. (F) Detail of the peridial wall. (G–I) Smooth-walled ascospores. (J) Reticulate ascospores. (K) Donut-like structure surrounding the germ pore of an ascospore. (L and M) Bulbils. (N and P) Phialidic conidiogenous cells (white arrows) with catenate conidia.

Microthecium tenuissimum (D. García, Stchigel & Guarro) Y. Marín, Stchigel, Guarro & Cano (the morphologically nearest species) (CBS 112764, GenBank NR.161011.1; Identities = 266/269 (98.88%), 0 gaps), *Microthecium fimicola* (E.C. Hansen) Y. Marín, Stchigel, Guarro & Cano (CBS 967.97, GenBank MK926777.1; Identities = 266/170

(98.52%), one gap) and *Microthecium quadrangulatum* (D. García, Stchigel & Guarro) Y. Marín, Stchigel, Guarro & Cano (CBS 112763, GenBank MK926778.1; Identities = 266/270 (98.52%), one gap). When using the *act* sequence (820 bp, accession number OX383416.1), percentage identities were greater than 99%

for *Microthecium compresum* Udagawa & Cain (NBRC 8627, GenBank KP981525.1; Identities = 772/772 (100%), 0 gaps), *M. fimicola* (NBRC 8354, GenBank KP981528.1; Identities = 771/772 (99.87%), 0 gaps) and *Microthecium zobelii* Corda (NBRC 9442, GenBank KP981542.1; Identities = 770/772 (99.74%), 0 gaps); with the *tub2* sequence (476 bp, accession number OX383417.1) the species displayed were *M. tenuissimum* (CBS 112764, GenBank MK926880.1; Identities = 417/417 (100%), 0 gaps), *M. fimicola* (CBS 967.97, GenBank MK926877.1; Identities = 417/417 (100%), 0 gaps) and *M. zobelii* (CBS 341.73, GenBank MK926882.1; Identities = 416/417 (99.76%), 0 gaps); using *rpb2* sequence (640 pb, accession number OX383419.1) the species were *M. tenuissimum* (CBS 112764, GenBank MK876742.1; Identities = 527/527 (100%), 0 gaps), *M. fimicola* (CBS 967.97, GenBank MK876739.1; Identities = 526/527 (99.81%), 0 gaps) and *M. zobelii* (CBS 341.73, GenBank MK876744.1; Identities = 523/527 (99.24%), 0 gaps). Finally, with the *tef1* sequence (759 bp, accession number OX383418.1) the species displayed were *M. fimicola* (FMR 13148, GenBank KP981593.1; Identities = 748/750 (99.73%), 0 gaps), *Microthecium levitum* Udagawa & Cain (FMR 13884, GenBank KP981598.1; Identities = 742/750 (99.73%), 0 gaps) and *M. quadrangulatum* (CBS 112763, GenBank MK981599.1; Identities = 742/750 (99.33%), 0 gaps).

Taxonomy

Microthecium pleomorphosporum Stchigel, Pintos & Cano, sp. nov. Mycobank MB 841012 (Fig. 1).

Etymology: From Greek *πλέω*-, numerous, *-μορφή*-, form, and *-σπόρια*, spore, due to the variable shape of the ascospores.

Diagnosis: *M. pleomorphosporum* might be a new species withing the genus as it differs from the rest in producing two sorts of ascospores, smooth-walled and inconspicuously reticulated ones, plus a phialidic asexual morph and bulbils.

Morphological description: Hyphae are septate, hyaline, smooth- and thin-walled, anastomosing, frequently moniliform, 1–8 (–15) μm wide, that produce cylindrical, pale-orange arthroconidia in short chains when cultures are old. Sexual morph: *ascomata* perithecial, non-ostiolate, superficial to immersed and scattered on the culture medium, glabrous, translucent, pale-yellow to yellowish-orange at first, later becoming dark-brown due to the production of ascospores, opening when old by one or two undetermined sites at the *ascomata* wall, globose, 50–200 μm diam.; *ascomata* wall translucent, pale-yellow to yellowish-orange, 5–15 μm thick, composed by 2–6 layers of smooth- and thin-walled, flattened, pale-yellow to pale-orange cells of 5–30 μm diam., with *textura angularis* to *textura globu-*

losa; *asci* 8-spored, unitunicate, broadly ellipsoidal to sacciform, 32–35 \times 15–20 μm , soon evanescent, non-stipitate; unicellular ascospores, chocolate-brown, smooth-walled or delicately reticulate in part or in all their surface, citriform but flattened at one side, (13–) 15–18 \times 10–13 μm \times 8–9 μm , with a protruding germ pore surrounded by a donut-like structure derived from the outer wall of the ascospore. Asexual morph: conidiophores absent; phialides hyaline, smooth- and thin-walled, flask-shaped, 5–10 \times 2–3 μm , ventricose at the base or at the middle part, rarely at the upper part of the cell, arising laterally on the vegetative hyphae, delimited by a basal septum; conidia enteroblastic, unicellular, hyaline, smooth- and thin-walled, produced in short basipetal chains, globose to ellipsoidal, 2–3 \times 2 μm . Bulbils present, 4- to multicelled, yellowish to orange, up 30 μm diam, individual cells smooth- and thin-walled, more or less globose.

Cultural characteristics: Colonies on PCA grow rapidly, filling 90 mm diam. plates in 14 days at room temperature (22–25 °C). They have a submerged mycelium and aerial hyphae, granulose and brown (M. 6E6) due to the production of abundant ascomata; the reverse is grayish-brown (M. 6D4). Colonies on OA and PDA grow rapidly, filling 90 mm diam. plates in 14 days at room temperature; the color is dark-brown due to the production of abundant ascomata (M. 6E7). At 15 °C the colonies grow fast on all media, filling 65–90 mm diam. plates in 14 days, with abundant aerial mycelia, and ascomata abundantly produced. At 35 °C no growth was observed.

Holotype: Spain, Balearic Islands, Malla, Serra de la Tramuntana, Lluc, Menut, recovered from healthy stems of *C. monogyna* Jacq., 39°50'30.86"N, 2°54'03.63"E, altitude 522 msl, 05/IX/2020, collected by A. Pintos Amengual, identified by A. M. Stchigel, J. Cano and A. Pintos; holotype CBS H-27905, cultures ex-type FMR 18380 = AP5920 = CBS 148652.

Notes: *M. pleomorphosporum* is morphologically related to *M. tenuissimum* (D. García, Stchigel & Guarro) Y. Marín, Stchigel, Guarro & Cano.⁴ Both species produce two sorts of ascospores, smooth-walled and reticulate ones. However, they can be differentiated by the size of the ascospores [(13–) 15–18 \times 10–13 μm \times 8–9 μm in *M. pleomorphosporum* vs. 19–23 \times (12–) 14–15 (–17) \times 10–13 μm in *M. tenuissimum*]. *M. pleomorphosporum* produce, as well, the typical asexual morph of the genus (hyaline, flask-shaped single phialides arising laterally on the vegetative hyphae) and bulbils, which are absent in *M. tenuissimum*, and glabrous *ascomata* (bearing few short setae in *M. tenuissimum*).

Key to the species of *Microthecium* (based on Marin-Felix et al.⁹).

1. Sexual morph absent, only producing bulbils	<i>M. sepedonioides</i> (Preuss) Y. Marín, Stchigel, Guarro & Cano	2
Sexual morph present		3
2. Ascomata non-ostiolate		14
Ascomata ostiolate		4
3. Ascospores with an ornamented wall		8
Ascospores smooth-walled or nearly so		5
4. Ascospores pitted, with wing-like ridges	<i>M. foveolatum</i> Udagawa & Y. Horie	6
Ascospores coarsely reticulate		7
5. Asci 4-spored		7
Asci 8-spored		8
6. Ascospores (25–)28–34(–40) μm long	<i>M. beatonii</i> D. Hawksw.	9
Ascospores 22–28 μm long	<i>M. perplexum</i> D. Hawksw.	10
7. Ascospores 25–34 μm long	<i>M. episphaerium</i> (W. Phillips & Plowr.) Höhn	11
Ascospores 17–20 μm long	<i>M. retisporum</i> Udagawa & Cain	13
8. Ascomata always bigger than 120 μm diam		10
Ascomata from 50 μm diam		11
9. Ascospores shorter than 20 μm		11
Ascospores longer than 20 μm		12
10. Ascospores 15–19 \times 11–13 \times 8–9 μm , with the narrow faces coarsely reticulate and the others smooth	<i>M. compressum</i> Udagawa & Cain	12
Ascospores 10–17 \times 8–12 \times 9–10 μm , entirely smooth-walled	<i>M. levitum</i> Udagawa & Cain	12
11. Ascospores fusiform	<i>M. hypomyces</i> (Höhn.) Höhn.	12
Ascospores citriform		12
12. Ascospores 28–30 \times 12–13(–15) μm	<i>M. geopora</i> (W. Oberm.) Höhn.	12
Ascospores 18–25 \times 8.5–12 \times 6–9 μm	<i>M. zobelii</i> Corda	13
13. Ascospores 19–23 \times (12–)14–15(–17) \times 10–13 μm ; asexual morph absent	<i>M. tenuissimum</i> (D. García, Stchigel & Guarro) Y. Marín, Stchigel, Guarro & Cano	15
Ascospores (13–)15–18 \times 10–13 μm \times 8–9 μm ; asexual morph phialidic; bulbils produced	<i>M. pleomorphosporum</i> Stchigel, Pintos & Cano	15
14. Ascospores with wing-like appendages		16
Ascospores otherwise		16
15. Ascospores wrinkled, (12–)13–18 \times (7–)8–10 μm	<i>M. ciliatum</i> Udagawa & Takada	17
Ascospores pitted, (17–)20–22(–24) \times 12–14 \times 10–12 μm	<i>M. lenticulare</i> (Udagawa & T. Muroi) Y. Marín, Stchigel, Guarro & Cano	17
16. Ascospores ornamented		24
Ascospores smooth-walled		18
17. Ascospores punctate or punctate-reticulate		20
Ascospores reticulate or striate-reticulate		19
18. Ascospores punctate, ellipsoidal	<i>M. africanum</i> (J.C. Krug) Y. Marín, Stchigel, Guarro & Cano	19
Ascospores punctate-reticulate, ellipsoidal-fusiform		20
19. Ascospores delicately punctate; asexual morph and bulbils present	<i>M. japonicum</i> (Y. Horie, Udagawa & P.F. Cannon) Y. Marín, Stchigel, Guarro & Cano	21
Ascospores coarsely punctate; asexual morph and bulbils absent	<i>M. moreaui</i> (P.F. Cannon & D. Hawksw.) Y. Marín, Stchigel, Guarro & Cano	21
20. Ascospores striate-reticulate		22
Ascospores reticulate		22
21. Ascospores with inconspicuous ridges forming a very coarse reticulum, 18–22(–28) \times 9.5–11(–13) \times 8–9 μm	<i>M. micropertusum</i> (Y. Horie, Udagawa & P.F. Cannon) Y. Marín, Stchigel, Guarro & Cano	23
Ascospores without ridges or reticulum, 26–36 \times 13–17 μm	<i>M. mansonii</i> (Kirschst.) Y. Marín, Stchigel, Guarro & Cano	23
22. Ascospores with 4–6 prominent longitudinal ribs	<i>M. quadrangulare</i> (Dania García, Stchigel & Guarro) Y. Marín, Stchigel, Guarro & Cano	23
Ascospores without longitudinal ribs		23
23. Ascospores spindle-shaped, 19.5–22 \times 8.5–11 μm	<i>M. internum</i> (Tehon & G.L. Stout) Y. Marín, Stchigel, Guarro & Cano	26
Ascospores citriform to fusiform, 14–20 \times 10–17 μm	<i>M. fimicola</i> (E.C. Hansen) Y. Marín, Stchigel, Guarro & Cano	25
24. Crown of setae absent	<i>M. nectrioides</i> (Marchal) Y. Marín, Stchigel, Guarro & Cano	25
Crown of setae present		26
25. Ascospores citriform	<i>M. marchicum</i> (Lindau) Y. Marín, Stchigel, Guarro & Cano	27
Ascospores otherwise		29
26. Ascospores ellipsoid, often somewhat plataniiform		28
Ascospores otherwise		28
27. Bulbils present	<i>M. fallax</i> (Zukal) Y. Marín, Stchigel, Guarro & Cano	28
Bulbils absent		28
28. Ascospores 21–34 \times 11–17 μm	<i>M. brevirostre</i> (Fuckel) Y. Marín, Stchigel, Guarro & Cano	28
Ascospores 18–22 \times 9–11 μm	<i>M. fimbriatum</i> (Rostr.) Y. Marín, Stchigel, Guarro & Cano	29
29. Ascospores ellipsoid to fusiform	<i>M. fusisporum</i> (Petch) Y. Marín, Stchigel, Guarro & Cano	30
Ascospores ellipsoid to navicular		31
30. Ascospores (9.5–)11–12(–13) \times 4–4.5 μm	<i>M. pegleri</i> (D. Hawksw. & A. Henrici) Y. Marín, Stchigel, Guarro & Cano	31
Ascospores longer than 15 μm		31
31. Ascospores 16–24 \times 8–12 μm	<i>M. fayodi</i> (Vuill.) Y. Marín, Stchigel, Guarro & Cano	31
Ascospores 25–30 \times 11–15 μm	<i>M. brevirostratum</i> (Moreau) Y. Marín, Stchigel, Guarro & Cano	31

Discussion

Endophytic fungi live inside plants without causing disease, growing within the roots, stems and/or leaves. Some of them are capable of establishing a symbiotic relationship with the host, and have the ability to boost the plant growth and nutrient acquisition, increasing the host's resistance to drought and salinity, but also to the deleterious action of animals (insects, herbivorous mammals) and microbial pathogens.¹¹ Despite the fact that hundreds of articles have been published on this topic, and the increasing interest by the scientific community in the production of bioactive compounds by endophytic fungi,^{1,6,12} only a few studies have been carried out to know the biodiversity of endophytic fungi of the autochthonous Mediterranean plants. For this reason, it is not surprising that only *Aureobasidium* sp., *Chaetomium* sp., *Hormonema* sp. and *Torula herbarum* have been reported as endophytic fungi in *C. monogyna*.¹⁰

Most of the species of the genus *Microthecium* are saprobic on plant debris and soil, but few of them have been reported as mycoparasitic on *Fusarium* spp., i.e., *M. moreaui* (formerly *Persiciospora moreaui*), *M. mycoparasiticum* (syn. *Sphaerodes mycoparasitica*), *M. quadrangulatum* (syn. *Sphaerodes quadrangularis*) and *M. retisporum* (syn. *Sphaerodes retispora*).⁹ On the other hand, none of the species within the genus have been reported as plant pathogens or as endophytes. Consequently, *M. pleomorphosporum* is not only a new species, but the first endophytic species within its genus. Morphologically, *M. pleomorphosporum* is very alike to the nearest species, *M. tenuissimum*. However, both species are easily discriminated, because *M. pleomorphosporum* produces glabrous ascospores (these are sparsely setose in *M. tenuissimum*), which reach up to 200 µm in diam. (and up to 125 µm high in *M. tenuissimum*), ascospores smaller in size than *M. tenuissimum*, and has a phialidic asexual morph and bulbils, absent in *M. tenuissimum*.

Conclusions

M. pleomorphosporum is a new species of the genus, the first reported as a plant endophyte. Probably, new studies on endophytes in Mediterranean flora can increase the number of fungal taxa (including more members of the order Melanosporales) living into healthy plants, as well to know the role of these endophytes in preventing the development of plant diseases, due to the well known biotrophic activity of Melanosporales on plant pathogenic fungi. Like all other *Microthecium* species, *M. pleomorphosporum* is

impossible to discriminate/circumscribe at the phylogenetic level, but on the other hand it is easily recognizable by its strictly glabrous ascospores, the ornamentation of the ascospores (smooth-walled ones, and slightly reticulated others), and by the production of phialoconidia and bulbils.

Conflicts of interest

The authors declare having no conflict of interest.

References

- Ancheeva E, Daletos G, Proksch P. Bioactive secondary metabolites from endophytic fungi. *Curr Med Chem*. 2020;27:1836–54. <http://dx.doi.org/10.2174/0929867326666190916144709>.
- Cannon PF, Hawksworth DL. A re-evaluation of *Melanospora* Corda and similar Pyrenomycetes, with a revision of the British species. *Bot J Linn Soc*. 1982;84:115–60. <http://dx.doi.org/10.1111/j.1095-8339.1982.tb00363.x>.
- Crous PW, Verkley GJM, Groenewald JZ, Houbraken J. *Westerdijk Laboratory Manual Series No. 1. Fungal Biodiversity*; 2019 [ISBN: 9789491751165].
- García D, Stchigel AM, Guarro J. Two new species of *Sphaerodes* from Spanish soils. *Stud Mycol*. 2004;50:63–8.
- Houbraken J, Due M, Varga J, Meijer M, Frisvad JC, Samson RA. Polyphasic taxonomy of *Aspergillus* section *Usti*. *Stud Mycol*. 2007;59:107–28. <http://dx.doi.org/10.3114/sim.2007.59.12>.
- Jia M, Chen L, Xin H-L, Zheng C-J, Rahman K, Han T, et al. A friendly relationship between endophytic fungi and medicinal plants: a systematic review. *Front Microbiol*. 2016;7:906. <http://dx.doi.org/10.3389/fmicb.2016.00906>.
- Kornerup A, Wanscher JH. *Methuen handbook of colour*. 3rd edn London, England: Eyre Methuen; 1978. p. 252 [ISBN-13, 978-0413334008].
- Magaña-Dueñas V, Stchigel AM, Cano-Lira JF. New coelomycetous fungi from freshwater in Spain. *J Fungi*. 2021;7:368. <http://dx.doi.org/10.3390/jof7050368>.
- Marin-Felix Y, Guarro J, Cano-Lira JF, García D, Miller AN, Stchigel AM. *Melanospora* (Sordariomycetes, Ascomycota) and its relatives. *Mycoskeys*. 2018;44:81–122. <http://dx.doi.org/10.3897/mycokeys.44.29742>.
- Rashmi M. A worldwide list of endophytic fungi with notes on ecology and diversity. *Mycosphere*. 2019;10:798–1079. <http://dx.doi.org/10.5943/mycosphere/10/1/19>.
- Rodriguez RJ, White JF Jr, Arnold AE, Redman RS. Fungal endophytes: diversity and functional roles. *New Phytol*. 2009;182:314–30. <http://dx.doi.org/10.1111/j.1469-8137.2009.02773.x>.
- Strobel G. Special issue: fungal endophytes in plants. *J Fungi*. 2018;4:104. <http://dx.doi.org/10.3390/jof4030104>.
- Sung G-H, Sung J-M, Hywel-Jones NL, Spatafora JW. A multi-gene phylogeny of *Clavicipitaceae* (Ascomycota, Fungi): identification of localized incongruence using a combinatorial bootstrap approach. *Mol Phylogenet Evol*. 2007;44:1204–23. <http://dx.doi.org/10.1016/j.ympev.2007.03.011>.
- Voigt K, Wöstemeyer J. Reliable amplification of actin genes facilitates deep-level phylogeny. *Microbiol Res*. 2000;155:179–95. [http://dx.doi.org/10.1016/S0944-5013\(00\)80031-2](http://dx.doi.org/10.1016/S0944-5013(00)80031-2).
- Woudenberg JHC, Aveskamp MM, de Gruyter J, Spiers AG, Crous PW. Multiple *Didymella* teleomorphs are linked to the *Phoma clematidina* morphotype. *Persoonia*. 2009;22:56–62. <http://dx.doi.org/10.3767/003158509X427808>.