



## Genome Announcements

# Draft genome sequence of a GES-5-producing *Serratia marcescens* isolated in southern Brazil



Carolina Silva Nodari<sup>a,b,\*</sup>, Marina Siebert<sup>c</sup>, Ursula da Silveira Matte<sup>d,e</sup>,  
Afonso Luís Barth<sup>a,b</sup>

<sup>a</sup> Universidade Federal do Rio Grande do Sul, Programa de Pós-Graduação em Ciências Farmacêuticas, Porto Alegre, RS, Brazil

<sup>b</sup> Hospital de Clínicas de Porto Alegre, Laboratório de Pesquisa em Resistência Bacteriana (LABRESIS), Porto Alegre, RS, Brazil

<sup>c</sup> Hospital de Clínicas de Porto Alegre, Unidade de Análises Moleculares e de Proteínas, Porto Alegre, RS, Brazil

<sup>d</sup> Hospital de Clínicas de Porto Alegre, Centro de Pesquisa Experimental, Porto Alegre, RS, Brazil

<sup>e</sup> Universidade Federal do Rio Grande do Sul, Departamento de Genética, Porto Alegre, RS, Brazil

## ARTICLE INFO

## Article history:

Received 21 March 2016

Accepted 12 August 2016

Available online 23 November 2016

Associate Editor: Rodrigo Galhardo

## Keywords:

GES-5

*Serratia marcescens*

Whole-genome sequencing

## ABSTRACT

*Serratia marcescens* is a Gram-negative rod intrinsically resistant to polymyxins and usually associated with wound, respiratory and urinary tract infections. The whole genome of the first GES-5-producing *S. marcescens* isolated from a Brazilian patient was sequenced using Ion Torrent PGM System. Besides *bla*<sub>GES-5</sub>, we were able to identify genes encoding for other  $\beta$ -lactamases, for aminoglycoside modifying enzymes and for an efflux pump to tetracyclines.

© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Nowadays carbapenemase production is the main carbapenem resistance mechanism among *Enterobacteriaceae*.<sup>1</sup> The Guiana Extended-Spectrum (GES)  $\beta$ -lactamase family comprises several Ambler class A enzymes with distinguished  $\beta$ -lactam hydrolysis profiles. The original GES were classified as extended-spectrum  $\beta$ -lactamases (ESBL), but amino acid substitutions in the GES-type ESBLs enhanced their activity against carbapenems.<sup>2</sup> The GES-5 is the GES-carbapenemase which hydrolyses imipenem most efficiently.<sup>3</sup> Here we report the draft genome of the first GES-5-producing *Serratia marcescens* reported in Brazil.

As part of a surveillance study,<sup>4</sup> isolates with reduced susceptibility to carbapenems were submitted to Real Time

(RT) High Resolution Melting (HRM) Multiplex PCR with specific primers for *bla*<sub>NDM</sub>, *bla*<sub>KPC</sub>, *bla*<sub>VIM</sub>, *bla*<sub>GES</sub>, *bla*<sub>OXA-48-like</sub> and *bla*<sub>IMP</sub>.<sup>5</sup> One isolate, obtained from an ascitic fluid of a female patient in a tertiary hospital in Porto Alegre (Brazil) in October 2014, and identified as *S. marcescens* by the Vitek2 system, presented an amplicon with a Temperature of Melting (*T*<sub>m</sub>) similar to the *bla*<sub>GES</sub> positive control in the RT-PCR – 85.32 °C and 85.52 °C, respectively. Antimicrobial susceptibility was determined by Etest and the isolate presented high resistance levels to carbapenems (MIC > 32 mg/L) and polymyxins (MIC > 256 mg/L), but remained susceptible to fluoroquinolones and tigecycline.

\* Corresponding author.

E-mail: [carol.nodari@gmail.com](mailto:carol.nodari@gmail.com) (C.S. Nodari).

<http://dx.doi.org/10.1016/j.bjm.2016.08.002>

1517-8382/© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Whole genome sequencing (WGS) was performed using the Ion Torrent PGM™ system, with a 400-bp-read kit and a 316™ Chip. Library was previously obtained by enzymatic fragmentation. We obtained 997,155 reads with a mean read length of 232 bp. The reads were assembled in contigs using SPAdes.<sup>6</sup> The assembling revealed a 5,378,959 bp length genome, distributed in 208 contigs ( $\geq 500$  bp), with a total GC content of 59%.

The contigs were annotated using the NCBI pipeline<sup>7</sup> and manually curated using Artemis,<sup>8</sup> when necessary. We also submitted the contigs to ResFinder Database.<sup>9</sup> Annotation revealed 3657 CDS, as well as 70 tRNA and 14 rRNA genes. As expected, we were able to identify the presence of *bla*<sub>GES-5</sub> (locus tag AN414.25255) after annotation, as well as genes coding for other  $\beta$ -lactamases (*bla*<sub>CTX-M-2</sub> and *bla*<sub>OXA-2</sub>, locus tags AN414.24540 and AN414.25310, respectively), and aminoglycoside modifying enzymes (*aac*(3)-IIa and *aac*(6')-Ic, locus tags AN414.24600 and AN414.08220, respectively). We were also able to observe the presence of a gene coding for an efflux pump to tetracyclines (*tet*(41), locus tag AN414.07940). The location of the resistance determinants in the genome (chromosome or plasmid borne) was not determined. To the best of our knowledge, this is the first report of a GES-5-producing *S. marcescens* in Brazil. We also demonstrated that NGS platforms can be used as a valuable tool to evaluate resistance determinants among *Enterobacteriaceae*.

**Accession number:** This whole-genome shotgun project has been deposited at GenBank under the accession number LNZT00000000. The version described in this paper is in the first version, LNZT01000000. The BioProject ID is PRJNA294719.

---

## Conflicts of interest

The authors declare no conflicts of interest.

## REFERENCES

---

1. Nordmann P, Dortet L, Poirel L. Carbapenem resistance in *Enterobacteriaceae*: here is the storm! *Trends Mol Med*. 2012;18:263-272.
2. Walsh TR. Clinically significant carbapenemases: an update. *Curr Opin Infect Dis*. 2008;21:367-371.
3. Bae IK, Lee YN, Jeong SH, et al. Genetic and biochemical characterization of GES-5, an extended-spectrum class A beta-lactamase from *Klebsiella pneumoniae*. *Diagn Microbiol Infect Dis*. 2007;58:465-468.
4. Rozales FP, Ribeiro VB, Magagnin CM, et al. Emergence of NDM-1-producing *Enterobacteriaceae* in Porto Alegre, Brazil. *Int J Infect Dis*. 2014;25:79-81.
5. Monteiro J, Widen RH, Pignatari AC, et al. Rapid detection of carbapenemase genes by multiplex real-time PCR. *J Antimicrob Chemother*. 2012;67:906-909.
6. Bankevich A, Nurk S, Antipov D, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012;19:455-477.
7. Tatusova T, DiCuccio M, Badretdin A, et al. Prokaryotic genome annotation pipeline. In: *The NCBI Handbook [Internet]*. 2nd ed; 2013.
8. Carver T, Harris SR, Berriman M, et al. Artemis: an integrated platform for visualization and analysis of high-throughput sequence-based experimental data. *Bioinformatics*. 2012;28:464-469.
9. Zankari E, Hasman H, Cosentino S, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67:2640-2644.