



## Environmental Microbiology

# Heterologous expression of a rice metallothionein isoform (OsMTI-1b) in *Saccharomyces cerevisiae* enhances cadmium, hydrogen peroxide and ethanol tolerance

Zahra Ansarypour, Azar Shahpiri\*

Isfahan University of Technology, College of Agriculture, Department of Biotechnology, Isfahan, Iran

## ARTICLE INFO

## Article history:

Received 1 May 2016

Accepted 25 October 2016

Available online 4 February 2017

Associate Editor: Gisele Monteiro de Souza

## Keywords:

Cadmium

Ethanol

Hydrogen peroxide

Rice metallothionein

*S. cerevisiae*

## ABSTRACT

Metallothioneins are a superfamily of low-molecular-weight, cysteine (Cys)-rich proteins that are believed to play important roles in protection against metal toxicity and oxidative stress. The main purpose of this study was to investigate the effect of heterologous expression of a rice metallothionein isoform (OsMTI-1b) on the tolerance of *Saccharomyces cerevisiae* to Cd<sup>2+</sup>, H<sub>2</sub>O<sub>2</sub> and ethanol stress. The gene encoding OsMTI-1b was cloned into p426GPD as a yeast expression vector. The new construct was transformed to competent cells of *S. cerevisiae*. After verification of heterologous expression of OsMTI-1b, the new strain and control were grown under stress conditions. In comparison to control strain, the transformed *S. cerevisiae* cells expressing OsMTI-1b showed more tolerance to Cd<sup>2+</sup> and accumulated more Cd<sup>2+</sup> ions when they were grown in the medium containing CdCl<sub>2</sub>. In addition, the heterologous expression of GST-OsMTI-1b conferred H<sub>2</sub>O<sub>2</sub> and ethanol tolerance to *S. cerevisiae* cells. The results indicate that heterologous expression of plant MT isoforms can enhance the tolerance of *S. cerevisiae* to multiple stresses.

© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The yeast, *Saccharomyces cerevisiae*, has several properties which have established it as an important tool in the

expression of foreign proteins for research, industrial or medical use.<sup>1,2</sup> As a food organism, it is highly acceptable for the production of pharmaceutical proteins. In contrast, *Escherichia coli* has toxic cell wall pyrogens and mammalian cells may contain oncogenic or viral DNA, so that products

Abbreviations: MT, metallothionein; Os, *Oryza sativa*; GST, glutathione-S-transferase; ROS, reactive oxygen species.

\* Corresponding author at: Department of Biotechnology, College of Agriculture, Isfahan University of Technology, Isfahan 84156-83111, Iran.

E-mail: [a.shahpiri@cc.iut.ac.ir](mailto:a.shahpiri@cc.iut.ac.ir) (A. Shahpiri).

<http://dx.doi.org/10.1016/j.bjm.2016.10.024>

1517-8382/© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

from these organisms must be tested more extensively.<sup>3,4</sup> *S. cerevisiae* can be grown rapidly on simple media and to high cell density, its genetics are more advanced than any other eukaryote, and can be manipulated almost as readily as *E. coli*. As a eukaryote, *S. cerevisiae* is a suitable host organism for the high-level production of secreted as well as soluble cytosolic proteins.<sup>5</sup>

Industrial yeast strains during fermentation are exposed to various stresses such as osmotic shock, oxidative stress and toxicity of secondary metabolites which lead to the loss of biological products.<sup>6,7</sup> Molecular oxygen is relatively unreactive and harmless in its ground state, but can undergo partial reduction to form a number of reactive oxygen species (ROS), including the superoxide anion and hydrogen peroxide ( $H_2O_2$ ), which can further react to produce the highly reactive hydroxyl radical.<sup>8</sup> ROS are toxic agents that can damage a wide variety of cellular components resulting in lipid peroxidation, protein oxidation, and genetic damage through the DNA modification.<sup>8,9</sup> Yeast cells may be exposed to ROS generated by neutrophils and macrophages during immunological defense mechanisms and following exposure to numerous exogenous agents including xenobiotics, carcinogens, UV and ionizing radiation.<sup>7</sup>

*S. cerevisiae*, like all organisms, contains effective antioxidant defense mechanisms, which detoxify ROS as they are generated and maintain the intracellular redox environment in a reduced state.<sup>9</sup> In response to an oxidant challenge, yeast cells increase the synthesis of a number of antioxidant enzymes involved in the detoxification of ROS including glutathione reductase, superoxide dismutase, catalase, glutathione peroxidase and thioredoxins.<sup>9,10</sup> In addition to antioxidant enzymes, growing amount of evidences show a close relationship between oxidative stress and metallothioneins (MTs); the low molecular weight Cys-rich proteins which are ubiquitously present in eukaryotes and prokaryotes. In fact a variety of physical and chemical stresses associated with oxidative injury increase MTs synthesis.<sup>11-13</sup>

MTs belong to a superfamily of intracellular metal-binding proteins which is composed of 15 families.<sup>14,15</sup> MTs bind metals through the thiol groups of their Cys residues.<sup>14,16</sup> Induction of MTs synthesis by heavy metals is a very well-known phenomenon and accordingly MTs are regarded as biomarker to reflect heavy metal exposures in ecotoxicological assessment.<sup>15,17</sup> MTs have a strong effect in scavenging free radicals, which are produced under various stress conditions.<sup>13</sup> In Arabidopsis, it has been demonstrated that different types of MTs exhibit distinct and overlapping functions in maintaining the homeostasis of essential transition metals, detoxification of toxic metals, and protection against intracellular oxidative stresses.<sup>18-21</sup> Transgenic tobacco and yeast that overexpress MT from cotton displayed increased tolerance to environmental stresses, indicating the role of MT in response to abiotic stresses.<sup>22</sup>

Previously we heterologously expressed the isoform OsMTI-1b, a rice MT type 1, in *E. coli* as carboxyl-terminal extensions of glutathione-S-transferase (GST).<sup>15,23</sup> The cells expressing OsMTI-1b showed increased tolerance to  $Ni^{2+}$ ,

$Cd^{2+}$ , and  $Zn^{2+}$  and accumulated more metal ions compared with cells expressing GST alone. In addition, heterologous expression of OsMTI-1b conferred  $H_2O_2$  tolerance to *E. coli* cells.<sup>15</sup>

In this study, the gene encoding OsMTI-1b was heterologously expressed in *S. cerevisiae*. The tolerance of transgenic *S. cerevisiae* to cadmium, hydrogen peroxide as well as ethanol was examined and the accumulation of cadmium in the cells was determined.

## Materials and methods


### Cloning of genes encoding GST-OsMTI-1b

In order to clone the gene encoding GST-OsMTI-1b, the plasmid pET41a-OsMTI-1b was used as template.<sup>15</sup> The DNA fragment containing GST-OsMTI-1b was amplified using Pfu DNA polymerase (Thermo Scientific) in a reaction mixture containing template plasmid, deoxy nucleotides, reaction buffer and the primers 5'-ATATAAGCTTCGGATAACAATTCCTCT-3', which carries an HindIII restriction site at the 5' end (underlined) of forward primer, and 5'-ATTTCTCGAGTTAGCAGTTGCAAGGGTT-3' with a restriction site XhoI (underlined) at the 5' end of reverse primer. An addition of four bases was included at the 5' end in each oligonucleotide primer. The thermal profile was as follows: 1 cycle at 92 °C for 5 min; 35 cycles at 92 °C for 1 min; 63 °C for 1.5 min; 72 °C for 2 min and 1 cycle at 72 °C for 10 min. The PCR product was then digested with enzymes HindIII and XhoI and ligated using T4 ligase into p426GPD as a yeast expression vector,<sup>24</sup> after linearization with HindIII and XhoI. The ligation mix was used to transform competent *E. coli* cells (DH5 $\alpha$ ). Transformants were selected on LB medium supplemented with ampicillin (50 mg L<sup>-1</sup>). The resulting plasmid was termed GPD-GST-OsMTI-1b. The resulting plasmids were then verified by sequencing.

### Heterologous expression of OsMTI-1b in *S. cerevisiae*

The competent cells from *S. cerevisiae* strain CEN.PK 113-5D (5D) were provided by using lithium acetate method as previously described.<sup>25</sup> The new constructs GPD-GST-OsMTI-1b was transformed to competent cells of 5D to give new strain 5D-2527. A control strain was provided by transforming empty GPD to competent cells. Transformants were selected on plates containing SC-URA medium after incubation at 30 °C for 2-3 days.

The new strain 5D-2527 and control were grown in shake flasks containing 10 mL appropriate minimal medium [10%  $(NH_4)_2SO_4$ , 12%  $KH_2PO_4$ , 5%  $MgSO_4$ , trace metals 1 mL L<sup>-1</sup>, 2% glucose and vitamins] at 30 °C under vigorous agitation for overnight. Then 4 mL of the overnight cultures of cells were inoculated in 80 mL of minimal medium supplemented with 2% glucose. When the O.D.<sub>600</sub> of cultures were about 2.5 and 3.5, 1 mL samples of culture medium were harvested by centrifugation at 12,000  $\times g$  for 15 min, and frozen at -20 °C until use.



**Fig. 1 – Amino acid sequence and expression vector maps. (A) The amino acid sequence of OsMTI-1b. The Cys residues are shown as bold. (B) The map of GPD-GST-OsMTI-1b. The position of His.tag, S.tag and GST.tag is shown in gray box.**

#### Extraction and identification of recombinant proteins with western blot analyses

The extraction of total protein was performed as explained previously.<sup>26</sup> For extraction of total proteins, the frozen pellets were first pre-treated with 2.0 M LiAc for 10 min at room temperature. After centrifugation ( $12,000 \times g$  for 10 min,  $4^\circ\text{C}$ ), the remaining pellets were resuspended in 0.4 M NaOH for 5 min on ice, followed by centrifuging ( $12,000 \times g$  for 10 min,  $4^\circ\text{C}$ ), the remaining pellets were resuspended in 100  $\mu\text{L}$  SDS sample buffer (0.06 M Tris-HCl pH 6.8, 10% (v/v) glycerol, 2% (w/v) sodium dodecyl sulfate (SDS), 5% (v/v) 2-mercaptoethanol, 0.0025% (w/v) bromophenol blue) and boiled for 5 min. Then, the cell lysate was centrifuged to remove the cell debris. The total proteins recovered in supernatant phase were analyzed by 12% SDS-PAGE and stained by Coomassie Brilliant Blue R-250.<sup>27</sup>

For western blotting, proteins were transferred to a PVDF membrane (Roche Applied Science) according to the manufacturer's instructions. The blots were blocked overnight in blocking buffer containing 5% (w/v) skimmed milk in TBST (10  $\text{mmol L}^{-1}$  NaCl, 25  $\text{mmol L}^{-1}$  Tris-HCl, pH 7.5, 0.1% (v/v) Tween 20). The rabbit His-Tag antibody (GenScript), was diluted 1:1000 in TBST. The antigen-antibody interaction was carried out at room temperature for 1 h. The blots were washed ( $3 \times 10$  min) with TBST followed by ( $1 \times 10$  min) TBS (10  $\text{mmol L}^{-1}$  NaCl, 25  $\text{mmol L}^{-1}$  Tris-HCl, pH 7.5). Blots were then probed with goat-anti-rabbit IgG conjugated with horse radish peroxidase (GenScript) diluted 1:2000 in TBST as secondary antibody. The membranes were washed again, as explained before. After washing, the immune blots were developed using 0.5  $\text{mg mL}^{-1}$  diaminobenzidine in 50  $\text{mmol L}^{-1}$  Tris-HCl, pH 7 and 0.22% hydrogen peroxide.

#### Tolerance of *S. cerevisiae* cells to $\text{Cd}^{2+}$ , $\text{H}_2\text{O}_2$ and ethanol

Tolerance of new strain and control to metals,  $\text{H}_2\text{O}_2$  and ethanol in the growth medium was examined in different concentrations of  $\text{CdCl}_2 \cdot \text{H}_2\text{O}$  (0.3 and 0.9 mM),  $\text{H}_2\text{O}_2$  (1 and 3 mM) and ethanol (7 (v/v) % and 10 (v/v) %). For these analyses, 5 mL samples of the overnight cultures were inoculated in 80 mL of minimal medium supplemented with 2% glucose. The cultures were incubated at  $30^\circ\text{C}$  with shaking. The


culture media containing 5D-2527 or control strain were supplemented with cadmium at concentrations of 0.3 and 0.9 mM when O.D.<sub>600</sub> reached 3.3. The growth was monitored up to 14 h by O.D.<sub>600</sub> measurements. Each data represents the mean  $\pm$  SD obtained from three independent experiments with two replicates. For the analysis of cadmium in medium, cells from 10 mL of culture at 0 (T0) and 5 h after cadmium addition (T1) were precipitated by centrifugation at  $6000 \times g$  for 20 min. The supernatant was analyzed for  $\text{Cd}^{2+}$  using inductively coupled plasma atomic absorption spectroscopy (Perkin Elmer AAnalyst 700). The cadmium concentration changes in the medium of control and 5D-2527 between T1 and T0 ( $C_{T0}-C_{T1}$ ) were calculated. Each data is the mean  $\pm$  SD obtained from three independent experiments with two replicates. Differences at the 5% level were considered significant as analyzed by the paired student's t-test ( $p < 0.05$ ).

For analysis of  $\text{H}_2\text{O}_2$  and ethanol tolerance,  $\text{H}_2\text{O}_2$  and ethanol were added to the culture media after 7 h when O.D.<sub>600</sub> reached 3.6. The growth was monitored up to 14 h by O.D.<sub>600</sub> measurements. Each data represents the mean  $\pm$  SD obtained from three independent experiments with two replicates. Differences at the 5% level were considered significant as analyzed by the paired student's t-test ( $p < 0.05$ ).

## Results

### Heterologous expression of GST-OsMTI-1b in *S. cerevisiae*

The full-length coding sequence of OsMTI-1b consists of 219 bp encoding a protein with 72 amino acids and theoretical molecular weight (kDa)/pI of 7.13/5.08. The predicted protein consists of six C-X-C motifs equally distributed in N- and C-terminals, as other plant type 1 MT proteins (Fig. 1A). The protein was expressed as carboxyl-terminal extensions of GST.tag, 6 His.tag and S.tag which we entirely named GST in this study (Fig. 1B). The use of GST has the advantages of high solubility and stability against proteolytic degradation.<sup>28</sup> Therefore, the protein GST-OsMTI-1b is predicted to have theoretical molecular weight (kDa)/pI of 39.94/5.93. Western blot analysis showed a sharp protein band of expected molecular mass (Fig. 2, lanes 3 and 4).



**Fig. 2 – SDS-PAGE and western blot analysis of total proteins extracted from control strain (lanes 1 and 2) and 5D-2527 (lanes 3 and 4) when the O.D.<sub>600</sub> of cell cultures were 2.5 (lanes 1 and 3) and 3.5 (lanes 2 and 4). The purified GST-OsMTI-1b obtained from heterologous expression in *E. coli*<sup>15</sup> was run in lane P.**

#### Tolerance to H<sub>2</sub>O<sub>2</sub> and ethanol

The comparison of growth curves of strain 5D-2527 and control reveals that in the H<sub>2</sub>O<sub>2</sub>/ethanol free medium, the maximum O.D.<sub>600</sub> for control strain (O.D.<sub>max600</sub> 6.6 ± 0.2) is almost higher than that for the strain 5D-2527 (O.D.<sub>max600</sub> 6 ± 0.2). However, in the presence of 1 mM H<sub>2</sub>O<sub>2</sub> the maximum O.D.<sub>600</sub> of strain 5D-2527 (O.D.<sub>max</sub> 6.2 ± 0.24) was higher than that for the control strain (O.D.<sub>max600</sub> 4.6 ± 0.2, Fig. 3 A). In the presence of 3 mM H<sub>2</sub>O<sub>2</sub> the maximum O.D.<sub>600</sub> of strains 5D-2527 and control strain were 4.8 ± 0.2 and 3.8 ± 0.17, respectively. In addition, in the presence of 7 and 10% ethanol the maximum O.D.<sub>600</sub> values for the control strain were 4.6 ± 0.2 and 4.2 ± 0.2, respectively. These values were reached 5.6 ± 0.18 and 5.1 ± 0.25 for 5D-2527 in the same conditions (Fig. 3B). These results suggest that the heterologous expression of GST-OsMTI-1b confers H<sub>2</sub>O<sub>2</sub> and ethanol tolerance in *S. cerevisiae* cells.

#### The cells expressing OsMTI-1b have more tolerance to cadmium

In the presence of 0.3 and 0.9 mM Cd<sup>2+</sup> the maximum O.D.<sub>600</sub> for control strain were 3.2 ± 0.25 and 3.4 ± 0.2, respectively. These values reached 4 ± 0.2 and 4.7 ± 0.15 for the strain 5D-2527 when it was grown in the same conditions. These results suggest that the heterologous expression of GST-OsMTI-1b confers cadmium tolerance in *S. cerevisiae* cells.

To determine whether the enhanced tolerance of 5D-2527 to cadmium was due to increased accumulation of this metal ion in the cells, the concentrations of cadmium ions in the culture media inoculated with the 5D-2527 and control strains were determined by atomic absorption at 0 h (T<sub>0</sub>) and 5 h (T<sub>1</sub>) after the addition of cadmium to the cultures. The concentration of cadmium ions at T<sub>1</sub> was similar to that at T<sub>0</sub>

for the control strain whereas the concentrations of cadmium ions were decreased by 5.57 ppm at T<sub>1</sub> in the culture medium of 5D-2527 (Fig. 4).


#### Discussion

*S. cerevisiae* is the most important industrial yeast and the main microorganism employed in bioethanol production.<sup>29</sup> Industrial yeast strains during fermentation are exposed to various stresses such as osmotic shock, oxidative stress and toxicity of secondary metabolites which lead to the loss of biological products.<sup>6,29</sup> During oxidative stress in *S. cerevisiae* the protein Yap1p as a transcription factor plays a key role in activation of transcription of many genes including TRX2,<sup>30</sup> encoding thioredoxin, TRR1 – thioredoxin reductase,<sup>31</sup> GSH2 – glutathione synthase, GPX2 – glutathione peroxidase<sup>32</sup> and TSA1 – thioredoxin peroxidase<sup>33,34</sup> via binding with specific DNA sequences localized in their promoters.<sup>35</sup> Correspondingly the over expression of genes TRX2 and TRR1 enhanced the resistance of *S. cerevisiae* to oxidative stress.<sup>30,31</sup>

Ethanol production by *S. cerevisiae* is one of the first biotechnological commodities for many years. However the inhibitory effect of ethanol accumulation on yeast growth during fermentation is still an important challenge for bio-ethanol production.<sup>36</sup> Depending on yeast strain and condition, ethanol inhibition on yeast growth begins at concentrations of less than 5%.<sup>37</sup> The mechanisms that yeasts cope with ethanol stress are very complicated and not fully understood. The quantification of the growth behavior of all available single gene deletion strains of *S. cerevisiae* under ethanol stress revealed that the growth of 446 deletion strains was defective indicating that many genes are involved in the ethanol resistance in yeast. The deletion of some of these genes like PEX genes whose products are involved in peroxisome transport exhibited the severest sensitivity to ethanol.<sup>38</sup> Recently the generation of new transgenic *S. cerevisiae* strains has been successfully implemented to improve ethanol tolerance.<sup>39-42</sup> For instance, the overexpression of genes *INO1*, *DOG1*, *HAL1* and truncated form of *MSN2* in *S. cerevisiae* resulted in remarkably increased tolerance to high concentrations of ethanol.<sup>41</sup>

In the present work the gene encoding OsMTI-1b was transferred to *S. cerevisiae*. MTs are Cys-rich and low molecular weight proteins characterized as important intracellular factors in detoxification of heavy metals in the cells of prokaryotes and eukaryotes.<sup>14,43</sup> However the role of MTs in the protection of plant and animal cells against oxidative stress has been recently demonstrated.<sup>22,44,45</sup> The isoform OsMTI-1b is known as a rice MT type 1 due to the presence of six Cys residues in N- and C-terminals. Previously this isoform was heterologously expressed in *E. coli*. Transformed cells showed increased tolerance to heavy metals cadmium, zinc and nickel and accumulated more metals as compared to the control strain.<sup>15,23</sup> Furthermore, the heterologous expression of OsMTI-1b conferred hydrogen peroxide tolerance to *E. coli* cells.<sup>15</sup> The data presented here show that the heterologous expression of OsMTI-1b was successful when it was expressed as carboxy-terminal extension of GST. The *S. cerevisiae* cells






**Fig. 3 – Effects of heterologous expression of GST-OsMTI-1b on the tolerance of *S. cerevisiae* to (A) H<sub>2</sub>O<sub>2</sub> (1 and 3 mM), (B) ethanol (7% and 10%) and (C) Cd<sup>2+</sup> (0.3 and 0.9 mM). Each data is the mean ± SD obtained from three independent experiments with two replicates. Differences at the 5% level were considered significant as analyzed by the paired student's t-test ( $p < 0.05$ ).**

expressing GST-OsMTI-1b were able to remove efficiently cadmium from culture medium and showed more tolerance in the medium containing cadmium as compared to the control strain. In addition, the cells expressing GST-OsMTI-1b

conferred H<sub>2</sub>O<sub>2</sub> tolerance to *S. cerevisiae*. Similar results have been previously reported for transgenic yeasts overexpressing cotton MT type 3 (GhMT3) in response to H<sub>2</sub>O<sub>2</sub> stress<sup>22</sup> or MT type 3 from *Tamarix hispida* (Th MT3) in response to metal



**Fig. 4 – Cadmium concentration variation in the medium of control strain and strain 5D-2527 between T1 (5 h after addition CdCl<sub>2</sub> to medium) and T0 (starting point of addition CdCl<sub>2</sub>). Each data is the mean  $\pm$  SD obtained from three independent experiments with two replicates. Asterisks indicate statistically significant differences from the corresponding control as analyzed by the paired student's t-test ( $p < 0.05$ ).**

toxicity.<sup>45</sup> Here we also found that the expression of GST-OsMTI-1b enhances the resistance of transgenic *S. cerevisiae* against ethanol.

Taken together on the basis of our results, the heterologous expression of isoform OsMTI-1b conferred improved tolerance against H<sub>2</sub>O<sub>2</sub>, metals and ethanol. Therefore plant genes encoding MT isoforms may be useful for augmenting the genetic toolbox for generating new strains with multiple stress resistance which could be very efficient for biofuel production.

### Conflicts of interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

### Acknowledgment

Iran National Science Foundation is thanked for financial support (grant No. 91001342).

### REFERENCES

- Gellissen G, Hollenberg CP. Application of yeasts in gene expression studies: a comparison of *Saccharomyces cerevisiae*, *Hansenula polymorpha* and *Kluyveromyces lactis*. *Gene*. 1997;190:87–97.
- Schmidt FR. Recombinant expression systems in the pharmaceutical industry. *Appl Microbiol Biotechnol*. 2004;65:363–372.
- Terpe K. Overview of bacterial expression systems for heterologous protein production: from molecular and biochemical fundamentals to commercial systems. *Appl Microbiol Biotechnol*. 2006;72:211–222.
- Khan KH. Gene expression in mammalian cells and its applications. *Adv Pharm Bull*. 2013;3:257–263.
- Nasser MW, Pooja V, Abdin MZ, Jain SK. Evaluation of yeast as an expression system. *Indian J Biotechnol*. 2003;2:477–493.
- Gomez-Pastor R, Pérez-Torrado R, Cabisco E, Ros J, Matallana E. Engineered Trx2p industrial yeast strain protects glycolysis and fermentation proteins from oxidative carbonylation during biomass propagation. *Microb Cell Fact*. 2012;11:1–15.
- Halliwell B. Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Plant Physiol*. 2006;141:312–322.
- Jamieson DJ. Oxidative stress responses of the yeast *Saccharomyces cerevisiae*. *Yeast*. 1998;14:1511–1527.
- Morano K, Grant Ch, Moye-Rowley W. The response to heat shock and oxidative stress in *Saccharomyces cerevisiae*. *Genetics*. 2012;190:1157–1195.
- Grant CM, Maclver FH, Dawes IW. Mitochondrial function is required for resistance to oxidative stress in the yeast *Saccharomyces cerevisiae*. *FEBS Lett*. 1997;410:219–222.
- Bauman JW, Liu J, Liu YP, Klaassen CD. Increase in metallothionein produced by chemical that induced oxidative stress. *Toxicol Appl Pharmacol*. 1991;110:347–354.
- Hidalgo J, Company L, Borrás M, Garvey JS, Armario A. Metallothionein response to stress in rats: role in free radical scavenging. *Am J Physiol*. 1988;255:518–524.
- Kagi JHR. Over view of metallothionein. *Methods Enzymol*. 1991;205:613–626.
- Cobbett C, Goldsbrough P. Phytochelatin and metallothioneins: roles in heavy metal detoxification and homeostasis. *Annu Rev Plant Biol*. 2002;53:159–182.
- Mohammadi Nezhad R, Shahpiri A, Mirlohi A. Discrimination between two rice metallothionein isoforms belonging to type1 and type 4 in metal binding ability. *Biotechnol Appl Biochem*. 2013;60:275–282.
- Freisinger E. Structural features specific to plant metallothioneins. *J Biol Inorg Chem*. 2011;16:1035–1045.
- Wong CKC, Yeung HY, Cheung RYH, Yung KKL, Wong MH. Ecotoxicological assessment of persistent organic and heavy metal contamination in Hong Kong coastal sediment. *Arch Environ Contam Toxicol*. 2000;38:486–493.
- García-Hernández M, Murphy A, Taiz L. Metallothioneins 1 and 2 have distinct but overlapping expression patterns in *Arabidopsis*. *Plant Physiol*. 1998;118:387–397.
- Lee J, Shim D, Song WY, Hwang I, Lee Y. *Arabidopsis* metallothioneins 2a and 3 enhance resistance to cadmium when expressed in *Vicia faba* guard cells. *Plant Mol Biol*. 2004;54:805–815.
- Miller JD, Artega RN, Pell EJ. Senescence-associated gene expression during ozone-induced leaf senescence in *Arabidopsis*. *Plant Physiol*. 1999;120:1015–1024.
- Robinson NJ, Tommey AM, Kuske C, Jackson PJ. Plant metallothioneins. *Biochem J*. 1993;295:1–10.
- Xue T, Li X, Zhu W, Wu C, Yang G, Zheng C. Cotton metallothionein GhMT3a, a reactive oxygen species scavenger, increased tolerance against abiotic stress in transgenic tobacco and yeast. *J Exp Bot*. 2009;60:339–349.
- Mohammadi Nezhad R, Shahpiri A, Mirlohi A. Heterologous expression and metal-binding characterization of a type 1 metallothionein isoform (OsMTI-1b) from rice (*Oryza sativa*). *Protein J*. 2013;32:131–137.
- Mumberg D, Muller R, Funk M. Yeast vectors for the controlled expression of heterologous proteins in different genetic backgrounds. *Gene*. 1995;156:119–122.
- Gietz RD, Woods RA. Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol*. 2002;350:87–96.
- Zhang T, Lei J, Yang H, Xu K, Wang R, Zhang Zh. An improved method for whole protein extraction from yeast *Saccharomyces cerevisiae*. *Yeast*. 2011;28:795–798.

27. Sambrook J, Fritsch EF, Maniatis T. *Maniatis Molecular Cloning: A Laboratory Manual*. 2nd ed. New York: Cold Spring Harbor Laboratory Press; 1989.
28. Esposito DC, Chatterjee DK. Enhancement of soluble protein expression through the use of fusion tags. *Curr Opin Biotechnol*. 2006;17:353–358.
29. Matsushika A, Inoue H, Kodaki T, Sawayama Sh. Ethanol production from xylose in engineered *Saccharomyces cerevisiae* strains: current state and perspectives. *Appl Microbiol Biotechnol*. 2009;84:37–53.
30. Kuge S, Jones N. YAP1 dependent activation of TRX2 is essential for the response of *Saccharomyces cerevisiae* to oxidative stress by hydroperoxides. *EMBO J*. 1994;13:655–664.
31. Morgan BA, Banks GR, Toone WM, Raitt D, Kuge S, Johnston LH. The Skn7 response regulator controls gene expression in the oxidative stress response of the budding yeast *Saccharomyces cerevisiae*. *EMBO J*. 1997;16:1035–1044.
32. Sugiyama K, Izawa S, Inoue Y. The Yap1p-dependent induction of glutathione synthesis in heat shock response of *Saccharomyces cerevisiae*. *J Biol Chem*. 2000;275:15535–15540.
33. Grant CM, Collinson LP, Roe J-H, Dawes IW. Yeast glutathione reductase is required for protection against oxidative stress and is a target gene for yAP-1 transcriptional regulation. *Mol Microbiol*. 1996;21:171–179.
34. Inoue Y, Matsuda T, Sugiyama K, Izawa S, Kimura A. Genetic analysis of glutathione peroxidase in oxidative stress response of *Saccharomyces cerevisiae*. *J Biol Chem*. 1999;274:27002–27009.
35. Toone WM, Jones N. AP-1 transcription factors in yeast. *Curr Opin Genet Dev*. 1999;9:55–61.
36. Stanley D, Bandara A, Fraser S, Chambers PJ, Stanley GA. The ethanol stress response and ethanol tolerance of *Saccharomyces cerevisiae*. *J Appl Microbiol*. 2010;109:13–24.
37. Pina C, Couto JA, Hogg T. Inferring ethanol tolerance of *Saccharomyces* and non-*Saccharomyces* yeasts by progressive inactivation. *Biotechnol Lett*. 2004;26:1521–1527.
38. Yoshikawa K, Tanaka T, Furusawa C, Nagahisa K, Hirasawa T, Shimizu H. Comprehensive phenotypic analysis for identification of genes affecting growth under ethanol stress in *Saccharomyces cerevisiae*. *FEMS Yeast Res*. 2009;9:32–44.
39. Alper H, Moxley J, Nevoigt E, Fink GR, Stephanopoulos G. Engineering yeast transcription machinery for improved ethanol tolerance and production. *Science*. 2006;314:1565–1568.
40. Hisashi Y, Hitoshi I, Hiroshi U. Disruption of URA7 and GAL6 improves the ethanol tolerance and fermentation capacity of *Saccharomyces cerevisiae*. *Yeast*. 2007;24:551–560.
41. Hong M, Lee K, Yu B, et al. Identification of gene targets eliciting improved alcohol tolerance in *Saccharomyces cerevisiae* through inverse metabolic engineering. *J Biotechnol*. 2010;149:52–59.
42. Watanabe M, Watanabe D, Akao T, Shimoi H. Overexpression of MSN2 in a sake yeast strain promotes ethanol tolerance and increases ethanol production in sake brewing. *J Biosci Bioeng*. 2009;107:516–518.
43. Palmiter RD. The elusive function of metallothioneins. *Proc Natl Acad Sci USA*. 1998;95:8428–8430.
44. Coyle P, Philcox JC, Carey LC, Roife AM. Metallothionein: the multi purpose protein. *Cell Mol Life Sci*. 2002;59:627–647.
45. Yang J, Wang Y, Yang C, Li C, Liu G. *Tamarix hispida* metallothionein-like ThMT3, a reactive oxygen species scavenger, increases tolerance against Cd<sup>2+</sup>, Zn<sup>2+</sup>, Cu<sup>2+</sup> and NaCl in transgenic yeast. *Mol Biol*. 2011;38:1567–1574.