



Environmental Microbiology

Screening of freshwater fungi for decolorizing multiple synthetic dyes



Panpan Yang^{a,b}, Wenzhao Shi^a, Hongkai Wang^{a,*}, Hongmei Liu^b

^a Zhejiang University, Biotechnology Institute, State Key Laboratory of Rice Biology, Hangzhou, Zhejiang, China

^b Shandong Agricultural University, College of Life Science, State Key Laboratory of Crop Biology, Tai'an, Shandong, China

ARTICLE INFO

Article history:

Received 22 May 2015

Accepted 6 March 2016

Available online 4 July 2016

Associate Editor: Cynthia Canêdo da Silva

Keywords:

Freshwater fungi

Biodegradation

Synthetic dyes

Decolorization

ABSTRACT

The biodegradation of synthetic dyes by fungi is emerging as an effective and promising approach. In the present study, freshwater fungal strains isolated from submerged woods were screened for the decolorization of 7 synthetic dyes. Subsequently, 13 isolates with high decolorization capability were assessed in a liquid system; they belonged to 9 different fungal species. Several strains exhibited a highly effective decolorization of multiple types of dyes. New absorbance peaks appeared after the treatment with 3 fungal strains, which suggests that a biotransformation process occurred through fungal biodegradation. These results showed the unexploited and valuable capability of freshwater fungi for the treatment of dye-containing effluents. The ability of certain fungi to decolorize dyes is reported here for the first time.

© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Synthetic dyes are widely used for coloring the products of several industries such as textiles, leather, cosmetics, paper, printing materials, and plastics. It is estimated that 1–2% of dye production is lost, and 5–10% is discharged to the environment when the dyes are used.^{1,2} Several dyes and the chemicals used to produce them are often toxic, carcinogenic or even explosive.^{3,4} Effluents from industries that use various dyes are considered as pollutants that can cause severe environmental problems as well as medical and esthetic problems.² The decolorization of this industrial waste is a challenging task because certain dyes are resistant to degradation.⁴

Physical and chemical decolorizing methods have been developed to remove dyes from wastewater; however, several

of them have disadvantages such as high costs and/or limited applicability.^{2,5} Studies on the capability of microorganisms to perform dye decolorization has received increasing attention because the use of microorganisms is considered a cost effective and environmentally friendly method for removing organic dyes from wastewater before they are discharged.^{6,7} Likewise, the capabilities and mechanisms of decolorization by bacteria have been studied.^{8,9} However, the application of bacteria was limited due to the narrow substrate range of various degrading bacteria.¹⁰ Moreover, to complete the degradation of dyes, different groups of bacteria are required in an alternation process from anaerobic to aerobic conditions.^{2,7}

Research on the fungal decolorization of dye wastewater has been performed in recent years.^{11,12} Several fungi with the capability to decolorize a wide range of dyes have been

* Corresponding author.

E-mail: hkwang@zju.edu.cn (H. Wang).

<http://dx.doi.org/10.1016/j.bjm.2016.06.010>

1517-8382/© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

reported. For example, the white-rot fungi and brown-rot fungi are well-studied fungi groups with decolorization abilities.¹² Other fungi, such as *Aspergillus niger*, *Rhizopus arrhizus*, and *R. oryzae*, can also decolorize and/or absorb diverse dyes and possess excellent color removal capabilities.^{13,14}

The mechanism of fungal decolorization mainly involves two aspects, biodegradation and biosorption.¹⁵ The biodegradation capability of fungi is due to their extracellular, non-specific and non-selective enzyme system.⁶ Fungal enzyme production depends on nutrient limitations, and their subsequent dye decolorization ability is achieved depending on the growth conditions.¹⁶ Considering the complex environmental factors involved in the dye wastewater conditions, the screening of more fungi is necessary for use in dye decolorization.

Aquatic fungi are the main decomposers of aquatic ecosystems and play crucial roles in the cycling of nutrients.¹⁷ In addition, a unique characteristic of fungi is their ability to produce several non-specific enzymes.¹⁸ These non-specific extracellular and/or exoenzymes enable the aquatic fungi to attack structurally diverse organic compounds that correspond to different pollutant classes.^{19,20} Hence, these fungi may serve as a new resource to treat wastewater. However, little research has been performed on the process of decolorizing wastewater by freshwater fungi,¹⁰ and more fungi with potential for the biodegradation of synthetic dyes need to be explored. In the present study, several freshwater fungi were isolated from streams in the Zhejiang Province, PR China, and their ability of decolorizing multiple synthetic dyes was evaluated.

Materials and methods

Dyes and media

Seven dyes were used for screening the decolorization ability of the fungal isolates. All the dyes were dissolved in distilled water at a concentration of 10 g/L; then, they were filtered and sterilized using a 0.22 µm-diameter bacteria filter before being tested for use. All the dyes were purchased from Sigma-Aldrich. The molecular structures of the dyes are listed in Table 1.








Water agar medium (WA; WA1.5% agar) was used for the fungal isolation; potato-carrot-agar medium (PCA; 5% potato, 5% carrot, 2% agar) was used for the identification of fungal isolates; potato-dextrose-agar medium (PDA; 20% potato, 2% dextrose, 2% agar) was used to grow the fungal cultures and for the DNA extraction; and malt-agar medium (MEA; 2% malt extract, 1.5% agar) was used for screening the decolorization by fungi.

The liquid MEA medium used for the decolorization test was the same as that described above but without agar. The PDA medium was used for colony growth, and the genomic DNA isolation was prepared following the method of Wong-sawas et al.²¹

Fungal strain isolation from streams

Submerged wood samples were randomly collected from streams in the Zhejiang Province; then, they were brought

Table 1 – Structures of the synthetic dyes used in the present study and the wavelength of maximum absorption – λ_{max} (nm).

Dye	Structure	λ_{max} (nm)
Acid Blue 40		620
Reactive Red 11		525
Acid Blue 193		580
Acid Blue 62		590
Acid Blue 113		630
Reactive Blue 74		590
Acid Red 73		500

back to laboratory and placed separately in snap lock plastic bags with sterile moist paper towels, incubated at room temperature and examined periodically during a three-month period. Single spore isolations were obtained using the methods described by Choi et al.^{22,23}

Dye decolorization on solid media

The solid media were prepared with MEA medium and the addition of each dye to a total concentration of 50 mg/L. A mycelium plug derived from the edge of fungal strains grown on WA medium plates for 4 days at 25 °C were transferred to the center of a solid medium plate and inoculated at 25 °C for 1 week. The formation of decolorized zones under or around the developing mycelia was monitored for dye decolorization. All the agar plate assays were performed in triplicate.

Morphological identification of fungal strains with decolorization capability

Pure cultures of fungi with decolorization capabilities were incubated on PCA, WA and PDA media at 25 °C for 2 weeks for their identification. The morphological characters of fungi were examined and recorded in detail under an Olympus BX-51 microscope equipped with an Olympus DP-50 digital camera system.

Molecular identification of the fungal strains tested

The total genomic DNA was extracted from the cultured mycelia with a modified CTAB method protocol.²¹ The complete internal transcribed spacers 1 and 2 and the 5.8S rDNA regions of the isolates tested were amplified by PCR using ITS1 and ITS4 primers. The 18S rDNA region of isolate JX-43 was obtained by PCR using the NS1 and NS4 primers because there is no ITS sequence of this species in GenBank. The PCR reaction protocol and sequencing process was performed as described by Wongsawas et al.²¹

Dye decolorization in liquid systems

The fungal strains that showed a decolorized zone on the solid media were picked out for further testing in liquid media. The concentration of each dye was supplemented to obtain 50 mg/L in a previously autoclaved liquid medium. A plug derived from the edge of the selected fungal strains on PDA was transferred to a 10 mL liquid medium that contained the respective dye and inoculated at 25 °C for 1 week. Then, the liquid medium that contained fungal mycelia was centrifuged at 5000 rpm for 10 min, and aliquots of the supernatants were transferred into new tubes and diluted as necessary for the absorbance readings. For each dye, untreated liquid medium was used as a control. The absorbance spectrum of each dye was examined using a U-2001 two-beam spectrophotometer using a wavelength range from 200 to 900 nm, and the maximum absorbance wavelength was determined according to the single peak with the strongest absorption. The absorbance determinations were measured at the maximum absorbance wavelength on a GENios + microplate reader. The standard curves of each dye were performed at the maximum absorbance wavelength, and the dye concentration ranged from 0 to 50 mg/L. The dye decolorization of samples was

expressed as a percentage, as follows: $D = (AC - AA)/AC \times 100$, where D is the decolorization rate (%), and AC and AA are the concentrations measured for the control and the treated sample, respectively.²⁴

The dye biosorption of fungal mycelia was evaluated as follows: after the centrifugal process described above, the fungal mycelia in each centrifuge tube were put on a filter paper (Whatman) to remove the additional liquid; then, they were transferred to a new centrifuge tube that contained 10 mL ethanol and 10 mL fresh liquid medium. These tubes were placed on a rotary shaker (160 rpm) at room temperature for 20 min and then centrifuged at 5000 rpm for 10 min. The control for each dye was performed by adding 10 mL ethanol into 10 mL liquid medium that contained 50 mg/L of each dye. The dye biosorption rate was expressed as a percentage, as follows: $A = AM/AD \times 100$, where A is the dye adsorption rate of each fungal mycelia (%), and AM and AD are the concentrations measured in the treatment and control, respectively.

The assays were performed in triplicate, the experimental error was calculated as the standard deviation (SD) and, in all the cases, the SD was below 3%.

Results

Freshwater fungal strain isolation and screening of the dye decolorization ability on agar plates

A total of 92 strains of freshwater fungi were isolated from decayed woody samples collected from streams in the Zhejiang Province, PR China. The decolorization capability of these fungal strains was assessed on dye-containing MEA plates with 7 types of synthetic dyes inoculated at 25 °C for 1 week.

Thirteen fungal strains that caused either a partial or full decolorization were screened out of 92 strains of freshwater fungi isolated from the decayed wood samples. The dye




Fig. 1 – Decolorization of 7 dyes by fungal strain C-1 on MEA medium that contained 50 mg/L of different dyes inoculated during 7 days. (A) Reactive Red 11; (B) Acid Blue 40; (C) Acid Blue 62; (D) Acid Blue 113; (E) Acid Blue 193; (F) Acid Red 73; (G) Reactive Blue 74.

decolorization results on agar plates showed that these fungal strains had the ability to decompose more than 3 types of dyes. Likewise, different fungi had different dye decolorization abilities. For example, strain C-1 had the capability to decompose all 7 different types of dyes tested in these experiments (Fig. 1).

Identification of the fungi with potential for dye decomposition

The identification of the 13 fungal strains was performed based on morphological characters and on the ITS rDNA sequence (Table 2). Based on the recorded morphological data (not shown) from 3 strains (DX-8, I-5 and C-1), it appears that they belong to the species *Myrothecium verrucaria*, with slightly different cultural characteristics (Fig. 2) and different abilities to decompose dyes (Table 3). The ITS regions of strain C-1 were sequenced, and a preliminary sequencing analysis confirmed that the taxon belonged to the *Myrothecium* species. The fungal strain I-6c was identified as *Colletotrichum dematum*. The fungal strain I-101 was similar to *Corynespora cassiicola*, while strains H-6 and Sy06 possessed morphology similar to that of *Dictyosporium zhejiangensis*. The morphology of I-10 and BD-42 were consistent with that of *Fusarium thapsinum*.²⁵ The fungal strain JX-62 had the morphological characters of *Alternaria alternata*.

The morphological characteristics of J-1 indicate a close resemblance to *Plectosporium tabacinum*. The ITS sequences further confirmed such an affinity. Strain JX-43 was similar to *Acrogenospora sphaerocephala*; however, the ITS sequence of this strain did not reveal high sequence similarities because no ITS sequence data of this species were found in the GenBank database. Then, we sequenced the SSU sequence of strain JX-43, and a blast search result indicated that it had 99% identity with *Farlowiella carmichaeliana*, the teleomorph of *A. sphaerocephala*. The D-1 strain could not be identified based on morphology because it was unable to sporulate; however, the ITS sequences show a close similarity to *Ceriporia lacerata*, known as the white-rot fungus.²⁶

All 7 types of dyes were used for the decolorization test in liquid media. The wavelengths of maximum absorption by



Fig. 2 – Colonial characterization of 3 strains of *Myrothecium verrucaria* incubated on PDA for 5 days at 25 °C. DX-8 produced abundant spores on the culture, but I-5 had no spores on the colony; the sporulation ability of C-1 was between that of DX-8 and I-5.

each dye are listed in Table 1. The 13 fungal strains that were significantly active in dye decolorization in solid plates were used in the liquid decolorization system assay. All the tested strains showed a highly efficient decolorization of a variety of dyes (Table 3). The decolorizing extent of the dyes by the different strains exhibited an individual pattern based on the dye type and the fungal strains, even those that belonged to the same fungal species. Each strain had the ability of decolorize multiple dyes and, at least, they decolorized one dye with high efficiency. For example, strain D-1 could decolorize all the tested dyes with a rate of up to 50%. Strains JX-43 and I-10 were able to decolorize the Reactive Red 11 and Acid Blue 113 dyes almost completely; however, only 34.1% and 41.3% decolorization rates were achieved with Reactive Blue 74, respectively.

Table 2 – Fungal strains used for the decolorization assay after the screening test and their blast results obtained from GenBank.

Strain number	Species name	Accession number of the tested isolates	Identity	Accession with high Ident in the GenBank
C-1	<i>Myrothecium verrucaria</i>	JQ801304	100%	FJ235085
DX-8	<i>Myrothecium verrucaria</i>	–	–	–
I-5	<i>Myrothecium verrucaria</i>	–	–	–
I-6C	<i>Colletotrichum dematum</i>	JQ801303	100%	AB046608
I-101	<i>Corynespora cassiicola</i>	JQ801302	99%	GU461298
Sy06	<i>Dictyosporium zhejiangensis</i>	KF931342	99%	FJ456893
H-6	<i>Dictyosporium zhejiangensis</i>	–	–	–
J-1	<i>Plectosporium tabacinum</i>	–	99%	KC845227
I-10	<i>Fusarium thapsinum</i>	–	–	–
DB-42	<i>Fusarium thapsinum</i>	KF813067	100%	JQ363735
JX-62	<i>Alternaria alternata</i>	KF813070	100%	JN681160
Jx-43	<i>Acrogenospora sphaerocephala</i>	KF836060	93%	AY265337
D-1	<i>Ceriporia lacerata</i>	KF78284	99%	FJ462746

Table 3 – Decolorization rates (%) of 11 dyes after the treatment with 13 fungal strains for 7 days in liquid media.

Dye	Acid Blue 40	Reactive Red 11	Acid Blue 193	Acid Blue 62	Acid Blue 113	Reactive Blue 74	Acid Red 73
<i>Myrothecium verrucaria</i> C-1	46.50 ± 0.15	50.64 ± 0.18	46.05 ± 0.32	67.09 ± 0.33	79.83 ± 1.06	66.74 ± 0.46	63.23 ± 0.52
<i>M. verrucaria</i> DX-8	45.41 ± 0.55	56.28 ± 0.46	41.47 ± 1.05	68.79 ± 0.23	77.59 ± 0.44	60.96 ± 0.96	61.54 ± 0.42
<i>M. verrucaria</i> I-5	48.57 ± 0.32	60.38 ± 0.34	41.46 ± 0.56	67.65 ± 0.24	79.51 ± 0.32	65.53 ± 0.73	58.00 ± 0.44
<i>Colletotrichum dematium</i> I-6c	69.60 ± 0.21	86.02 ± 0.29	53.58 ± 0.61	91.59 ± 0.32	85.86 ± 0.92	79.71 ± 0.86	54.96 ± 0.16
<i>Corynespora cassiicola</i> I-101	24.59 ± 0.33	44.39 ± 0.20	25.96 ± 0.12	48.69 ± 0.21	48.21 ± 0.82	51.74 ± 0.56	30.28 ± 0.86
<i>Dictyosporium zhejiangensis</i> Sy06	77.67 ± 0.90	62.56 ± 0.46	44.78 ± 0.11	61.55 ± 0.87	77.87 ± 0.23	66.43 ± 0.22	56.34 ± 0.63
<i>D. zhejiangensis</i> H-6	75.53 ± 0.24	64.02 ± 0.24	46.00 ± 1.12	58.83 ± 0.45	77.41 ± 0.72	64.04 ± 0.22	55.23 ± 0.57
<i>Plectosporium tabacinum</i> J-1	76.44 ± 1.23	95.96 ± 0.87	31.64 ± 0.79	50.16 ± 0.56	65.09 ± 0.65	56.95 ± 2.22	67.12 ± 0.33
<i>Fusarium thapsinum</i> I-10	93.97 ± 0.46	100 ± 0.00	92.93 ± 0.43	60.04 ± 0.68	98.64 ± 0.24	34.12 ± 0.33	51.76 ± 0.65
<i>F. thapsinum</i> DB-42	90.86 ± 0.94	92.81 ± 0.72	90.58 ± 0.61	66.21 ± 0.89	96.16 ± 0.36	40.29 ± 0.83	55.55 ± 0.54
<i>Alternaria alternata</i> Jx-62	49.16 ± 1.22	86.79 ± 0.23	45.71 ± 0.35	51.25 ± 0.34	91.11 ± 0.46	46.28 ± 0.74	66.43 ± 0.22
<i>Acrogenospora sphaerocephala</i> JX-43	89.84 ± 0.61	99.5 ± 0.21	78.00 ± 0.88	69.97 ± 0.67	100.00 ± 0.00	41.30 ± 0.56	61.98 ± 0.45
<i>Ceriporia lacerata</i> D-1	64.86 ± 0.34	67.12 ± 0.62	82.37 ± 0.22	70.98 ± 0.41	87.38 ± 0.12	69.64 ± 0.18	76.16 ± 0.72

Certain isolates of this study had a decolorization efficiency higher than that of the white-rot fungus *Ceriporia lacerata*. The obtained results are summarized in Table 3.

New absorbance peaks appeared after the treatment of dyes with the fungal strains. A new absorbance peak was shown when Acid Blue 62 was treated with sy06 and I-6C respectively, which has a maximum absorbance wavelength at 590 nm (Fig. 3).

In this study, the dye biosorption by the fungal strains was evaluated (Table 4). Different strains exhibited different extents of dye absorbance depending on the type of dyes. Strain I-101 had the ability of adsorbing 6 out of 7 dyes. In addition, 8 out of 13 strains were able to adsorb Acid Red 73, whereas the dye Acid Blue 193 exhibited a null adsorption by

all the fungal strains used in this study. Strain I-101 has the highest rate of biosorption among the tested fungal strains, and the efficiency of dye biosorption was more than 20% in 5 types of dyes. However, at least 4 strains, C-1, DX-8, I-5, DB-42, did not show any biosorption ability.

Discussion

The presence of dyes in industrial products currently results in wastewater pollution, and these pollutants need to be treated before they are discharged.²⁷ The ability of microorganisms to perform dye degradation has received considerable attention.^{7,16} The effluents from the textile industry are extremely variable in composition; even a small structural difference can affect the decolorization process.^{2,28} There is still a need for studies to discover the potential of fungal isolates for the improvement of dye wastewater because certain isolates cannot be suitable for a variety of dyes in wastewater conditions.¹ The fungal isolates in this research provide a new avenue for approaching the biodegradation of dyes in the future.

The fungi used for dye biodegradation were mainly isolated from forest ecosystems, such as white-rot and brown-rot fungi. Although these fungi have been shown to decolorize dyes in liquid fermentations, their enzyme production was shown to be unreliable. This outcome is mainly due to the unfamiliar environment of liquid fermentations for these forest fungi.¹² New isolates of fungi used to decolorize water conditions will probably have the efficacy of bioremediation of dyes because these fungi have the ability to secrete extracellular non-specific enzymes that contribute to the carbon recycling in liquid environments. Junghanns et al.¹⁰ showed that certain isolates of aquatic fungi had a high and unexploited potential for the treatment of dye-containing wastewaters, even at high concentrations. In the present study, 13 isolates of freshwater fungi showed a high efficiency for the decolorization of multiple dyes, which should provide advantages for the treatment of effluents with complex dye




Fig. 3 – Spectral curves of Acid Blue 62 treated with 4 fungal strains after 7 days. The new absorbance peak appeared at 500 nm when the dye was treated with the Sy06 and I-6c strains. CK0: ME broth medium. CK1: ME broth medium with 50 mg/L Acid Blue 62.

Table 4 – Biosorption rates (%) of 11 dyes after the treatment with 13 fungal strains for 7 days in liquid media.

Dye	Acid Blue 40	Reactive Red 11	Acid Blue 193	Acid Blue 62	Acid Blue 113	Reactive Blue 74	Acid Red 73
<i>Myrothecium verrucaria</i> C-1	0	0	0	0	0	0	0
<i>M. verrucaria</i> DX-8	0	0	0	0	0	0	0
<i>M. verrucaria</i> I-5	0	0	0	0	0	0	0
<i>Colletotrichum dematium</i> I-6c	8.58 ± 0.93	0	0	26.50 ± 0.64	1.40 ± 0.32	2.36 ± 0.35	14.03 ± 0.58
<i>Corynespora cassiicola</i> I-101	20.10 ± 0.36	29.01 ± 1.41	0	30.10 ± 0.73	26.17 ± 0.85	20.87 ± 0.88	1.12 ± 0.12
<i>Dictyosporium zhejiangensis</i> Sy06	0	0	0	0	0	0	32.0 ± 0.99
<i>D. zhejiangensis</i> H-6	0	0	0	0	0	0	41.84 ± 2.41
<i>Plectosporium tabacinum</i> J-1	0	0	0	0	0	0	10.03 ± 1.88
<i>Fusarium thapsinum</i> I-10	0	0	0	0	0	0	8.03 ± 2.46
<i>F. thapsinum</i> DB-42	0	0	0	0	0	0	0
<i>Alternaria alternata</i> Jx-62	2.22 ± 0.44	0	0	0	0	0	24.44 ± 1.53
<i>Acrogenospora sphaerocephala</i> JX-43	0	0	0	0	0	0	12.03 ± 0.66
<i>Ceriporia lacerata</i> D-1	0	0	0	4.78 ± 0.88	0	0	0

compositions. The decolorization abilities of several fungal species in decolorizing synthetic dyes, such as *A. sphaerocephala*, *D. zhejiangensis*, and *P. tabacinum*, have never been reported before.

The mechanisms of fungal dye removal are categorized into the following three groups: bioaccumulation, bioabsorption and biodegradation.^{7,16} Bioaccumulation is referred to the accumulation of pollutants by actively growing cells through their metabolism.¹⁶ Biosorption can occur in either living or dead biomass; amino, carboxyl, thiol, lipid and phosphate groups that exist in the fungal cell walls are responsible for binding the dye molecules in the biosorption process.^{16,29} The main mechanism of decolorization by fungi is biodegradation because the fungi can produce various non-specific extracellular and intracellular enzymes involved in the dye decolorization process, such as laccase, manganese peroxidase (MnP), manganese independent peroxidase (MIP), lignin peroxidase (LiP), tyrosinase, and others.^{1,16,29} The biodegradation ability of fungi through enzymatic activities was confirmed by using purified enzymes for decolorization.^{11,16} Moreover, there are several factors that can influence the rate of fungal dye biodegradation, such as environmental parameters, the dye molecule structures, and the relative contributions of enzymes involved in the decolorization, which may be different for each fungus.^{1,16,30} Further research is needed to understand the factors that influence the decolorization rate with the studied fungi.

Studies on the bio-decolorization of textile dye effluents by fungi have been reported even in the presence of salts and high dye concentration using the white-rot fungi,³¹ *Aspergillus fumigatus* and *Phanerochaete sordid*.^{28,32} In the present study, one isolate, D-1, which was identified as the white-rot fungus *C. lacerata*,²⁶ also showed a powerful ability to decolorize multiple dyes. Certain fungal isolates in this research, such as *C. dematum* strain I-6c, *F. thapsinum* strain DB-42, *F. thapsinum* strain I-10, and *A. sphaerocephala* strain JX-43, showed a higher potential for dye decolorization than the D-1 strain, which indicates their suitability for the treatment of dye-polluted

water. The assessment of the removal abilities of multiple dyes by certain fungal strains indicates that several enzymes were involved in the decolorization process although the enzymes contributing to dye decolourisation remain to be defined, for the dye biosorption contributing to decolorisation by mycelia was very low. Moreover, the high frequency appearance of new absorbance peaks after fungal treatment demonstrated the roles of the enzymes involved in both the decomposition and the biotransformation of dyes. This characterization of fungi revealed that they may be useful for the treatment of dye-containing wastewaters.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

This research was supported by the Major Science & Technology Project of the Xinjiang Uygur Autonomous Region (no. 201130102) and the Natural Science Foundation of the Zhejiang Province (LY15C140001).

REFERENCES

- Forgacs E, Cserhati T, Oros G. Removal of synthetic dyes from wastewaters: a review. *Environ Int.* 2004;30(7):953–971.
- Khan R, Bhawana P, Fulekar MH. Microbial decolorization and degradation of synthetic dyes: a review. *Rev Environ Sci Bio-Technol.* 2013;12(1):75–97.
- Attia AA, Girgis BS, Fathy NA. Removal of methylene blue by carbons derived from peach stones by H_3PO_4 activation: batch and column studies. *Dyes Pigm.* 2008;76(1):282–289.
- Sharma P, Kaur H, Sharma M, Sahore V. A review on applicability of naturally available adsorbents for the removal of hazardous dyes from aqueous waste. *Environ Monit Assess.* 2011;183(1–4):151–195.

5. Ali H. Biodegradation of synthetic dyes – a review. *Water Air Soil Pollut.* 2010;213:251–273.
6. Baccar R, Blanquez P, Bouzid J, Feki M, Attiya H, Sarra M. Decolorization of a tannery dye: from fungal screening to bioreactor application. *Biochem Eng J.* 2011;56(3):184–189.
7. Banat IM, Nigam P, Singh D, Marchant R. Microbial decolorization of textile-dye-containing effluents: a review. *Bioresour Technol.* 1996;58(3):217–227.
8. Saratale RG, Saratale GD, Chang JS, Govindwar SP. Bacterial decolorization and degradation of azo dyes: a review. *J Taiwan Inst Chem Eng.* 2011;42(1):138–157.
9. Singh P, Iyengar L, Pandey A. Bacterial decolorization and degradation of azo dyes. *Microb Degrad Xenobiot.* 2012;101–133.
10. Junghanns C, Krauss G, Schlosser D. Potential of aquatic fungi derived from diverse freshwater environments to decolourise synthetic azo and anthraquinone dyes. *Bioresour Technol.* 2008;99(5):1225–1235.
11. Shi L, Yu H, Dong T, et al. Biochemical and molecular characterization of a novel laccase from selective lignin-degrading white-rot fungus *Echinodontium taxodii* 2538. *Process Biochem.* 2014;49:1097–1106.
12. Singh AP, Singh T. Biotechnological applications of wood-rotting fungi: a review. *Biomass Bioenergy.* 2014;62:198–206.
13. Fu Y, Viraraghavan T. Fungal decolorization of dye wastewater: a review. *Bioresour Technol.* 2001;79:251–262.
14. Srinivasan A, Viraraghavan T. Decolorization of dye wastewaters by biosorbents: a review. *J Environ Manage.* 2010;91(10):1915–1929.
15. Vanhulle S, Trovaslet M, Enaud E, et al. Decolorization, cytotoxicity, and genotoxicity reduction during a combined ozonation/fungal treatment of dye-contaminated wastewater. *Environ Sci Technol.* 2008;42(2):584–589.
16. Kaushik P, Malik A. Fungal dye decolorization: recent advances and future potential. *Environ Int.* 2009;35(1):127–141.
17. Duarte S, Pascoal C, Cassio F, Barlocher F. Aquatic hyphomycete diversity and identity affect leaf litter decomposition in microcosms. *Oecologia.* 2006;147(4):658–666.
18. Bucher VV, Pointing SB, Hyde KD, Reddy CA. Production of wood decay enzymes, loss of mass, and lignin solubilization in wood by diverse tropical freshwater fungi. *Microb Ecol.* 2004;48(3):331–337.
19. Abdel-Raheem AM, Ali EH. Lignocellolytic enzyme production by aquatic hyphomycetes species isolated from the Nile's delta region. *Mycopathologia.* 2004;157(3):277–286.
20. Harms H, Schlosser D, Wick LY. Untapped potential: exploiting fungi in bioremediation of hazardous chemicals. *Nat Rev Microbiol.* 2011;9(3):177–192.
21. Wongsawas M, Wang HK, Hyde KD, Lin FC. *Dictyosporium zhejiangense* sp. nov., a new freshwater anamorphic fungus from China. *Cryptogam Mycol.* 2009;30(4):355–362.
22. Choi YW, Hyde KD, Ho WH. Single spore isolation of fungi. *Fungal Divers.* 1999;3:29–38.
23. Choi YW, Hyde KD, Ho WH, Mcgraw AC. Single spore isolation of fungi Single spore isolation of fungi. *Fungal Divers.* 1999;3:29–38.
24. Rosales E, Pazos M, Sanroman MA. Comparative efficiencies of the decolourisation of leather dyes by enzymatic and electrochemical treatments. *Desalination.* 2011;278(1–3):312–317.
25. Leslie JF, Summerell BA. *The Fusarium laboratory manual.* Ames, IA, USA: Blackwell Publishing Professional; 2006.
26. Lee JW, Gwak KS, Park JY, et al. Biological pretreatment of softwood *Pinus densiflora* by three white rot fungi. *J Microbiol.* 2007;45(6):485–491.
27. Mohorcic M, Teodorovic S, Golob V, Friedrich J. Fungal and enzymatic decolourisation of artificial textile dye baths. *Chemosphere.* 2006;63(10):1709–1717.
28. Jin XC, Liu GQ, Xu ZH, Tao WY. Decolorization of a dye industry effluent by *Aspergillus fumigatus* XC6. *Appl Microbiol Biotechnol.* 2007;74(1):239–243.
29. Kyzas GZ, Fu J, Matis KA. The change from past to future for adsorbent materials in treatment of dyeing wastewaters. *Materials.* 2013;6:5131–5158.
30. Güldönmez G. Influence of surfactants on dye removal and growth of *Aspergillus versicolor* – an effective way to decolorize textile dye. *Clean Soil Air Water.* 2014;42(7):917–922.
31. Faraco V, Pezzella C, Miele A, Giardina P, Sannia G. Bio-remediation of colored industrial wastewaters by the white-rot fungi *Phanerochaete chrysosporium* and *Pleurotus ostreatus* and their enzymes. *Biodegradation.* 2009;20(2):209–220.
32. Harazono K, Nakamura K. Decolorization of mixtures of different reactive textile dyes by the white-rot basidiomycete *Phanerochaete sordida* and inhibitory effect of polyvinyl alcohol. *Chemosphere.* 2005;59(1):63–68.