



Environmental Microbiology

Endophytic *Bacillus* strains enhance pearl millet growth and nutrient uptake under low-P



Vitória Palhares Ribeiro^a, Ivanildo Evódio Marriel^b, Sylvia Morais de Sousa^b, Ubiraci Gomes de Paula Lana^b, Bianca Braz Mattos^c, Christiane Abreu de Oliveira^{b,*}, Eliane Aparecida Gomes^{b,**}

^a Universidade Federal de São João del Rei, Sete Lagoas, MG, Brazil

^b Embrapa Milho e Sorgo, Sete Lagoas, MG, Brazil

^c Embrapa Solos, Rio de Janeiro, RJ, Brazil

ARTICLE INFO

Article history:

Received 16 March 2018

Accepted 22 June 2018

Available online 14 August 2018

Associate Editor: Ieda Mendes

Keywords:

Endophytic

Indole-acetic acid

Phosphate solubilizing

Plant growth promotion

Siderophores

ABSTRACT

Bacterial endophytes are considered to have a beneficial effect on host plants, improving their growth by different mechanisms. The objective of this study was to investigate the capacity of four endophytic *Bacillus* strains to solubilize iron phosphate (Fe-P), produce siderophores and indole-acetic acid (IAA) *in vitro*, and to evaluate their plant growth promotion ability in greenhouse conditions by inoculation into pearl millet cultivated in a P-deficient soils without P fertilization, with Araxá rock phosphate or soluble triple superphosphate. All strains solubilized Fe-P and three of them produced carboxylate-type siderophores and high levels of IAA in the presence of tryptophan. Positive effect of inoculation of some of these strains on shoot and root dry weight and the N P K content of plants cultivated in soil with no P fertilization might result from the synergistic combination of multiple plant growth promoting (PGP) traits. Specifically, while B1923 enhanced shoot and root dry weight and root N P content of plants cultivated with no P added, B2084 and B2088 strains showed positive performance on biomass production and accumulation of N P K in the shoot, indicating that they have higher potential to be microbial biofertilizer candidates for commercial applications in the absence of fertilization.

© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Embrapa Milho e Sorgo, Sete Lagoas, MG 35701-970, Brazil.

** Corresponding author at: Embrapa Milho e Sorgo, Sete Lagoas, MG 35701-970, Brazil.

E-mails: christiane.paiva@embrapa.br (C.A. Oliveira), eliane.a.gomes@embrapa.br (E.A. Gomes).

<https://doi.org/10.1016/j.bjm.2018.06.005>

1517-8382/© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Most of Brazilian soils, classified as Oxisols,¹ show low natural fertility, low pH, and high Al saturation.² Oxisols have low P availability due to high adsorption of P and the formation of Fe and Al phosphates. The most abundant and least soluble phosphate fraction in pasture, forest, and agricultural Oxisols is iron phosphate (Fe-P), followed by aluminum phosphate (Al-P) and calcium phosphate (Ca-P).^{3–6}

It is well documented that naturally occurring plant growth-promoting bacteria (PGPB) can act in insoluble phosphates form and release a soluble P in the soil solution readily absorbed by plant roots.^{4,7,8} In addition, PGPB can mitigate abiotic and biotic stresses on plants, including nutrient limitations, heat and drought, exposure to pollutants and antagonistic effect against phytopathogenic microorganisms. The use of PGPB is a favorable strategy in both environmental and economic aspects,^{7,8} presenting positive results for various crops, including common beans,⁹ wheat,^{7,10} maize,^{11,12} peanuts,¹² soybean,¹³ eggplants, tomatoes, and peppers.¹⁴

Most research on PGP microorganisms associated with plants is focused on rhizobacteria; however, there is an increasing interest in the diversity and role of endophytic bacteria for PGP.¹⁵ If bacteria are reintroduced in the endophytic form, a more stable relationship can be established between the plant and bacteria, which results in a more efficient promotion of plant growth.^{16,17}

Several rhizospheric or endophytic bacteria belonging to *Pseudomonas*, *Azospirillum*, *Azotobacter*, *Burkholderia*, *Bacillus*, *Enterobacter*, *Rhizobium*, and *Agrobacterium* have been reported as PGP microorganisms.¹⁸ Among them, *Bacillus* is frequently reported as a potential biofertilizer due to its multi-functional PGP traits, such as phosphate solubilization,^{13,14} indoleacetic acid (IAA) production,^{10,13} siderophore (iron chelator) production,¹⁹ and biocontrol ability against plant pathogens.²⁰ In addition, biofertilizers containing *Bacillus* strains are considered important because of their spore-forming capacity, allowing their adaptation to extreme abiotic conditions, such as extreme temperatures, pH, or pesticide exposure.¹⁴ *Bacillus* species have shown to positively affect soybean seed germination by enhancing significantly the root and shoot length, or number of lateral root of the seedling. This effect was related to the production of phytohormone and siderophore and the capability of these bacteria to solubilize phosphate.¹³ It has also been reported that *Bacillus* strains significantly promoted seed germination and growth of three test plants, tomato, pepper and eggplant, compared to control plants and the authors attributed this effect to plant-growth promoting traits.¹⁴ Other studies also revealed that *Bacillus* have improved plant growth under drought stress¹⁵ and produced a variety of compounds that can be used for the management of a broad range of plant pathogens.¹⁶

Previously, in our laboratory, endophytic *Bacillus* strains isolated from the roots, leaves, and sap of maize showed Ca-P-solubilizing and organic acid-producing ability *in vitro*.²¹ Bacterial strains that solubilize P-Ca showed potential as PGPs in soils fertilized with hydroxyapatite rock phosphate (RP)

rich in Ca.²² In the present study, four of these *Bacillus* isolates were selected, representing high (B1920 and B2088) and low (B2084 and B1923) Ca-P-solubilizing ability, to characterize their capacity to produce siderophores and IAA and solubilize Fe-P. Moreover, a greenhouse experiment was conducted to investigate the effects of endophytic *Bacillus* on pearl millet growth and nutrient uptake.

Materials and methods

Screening of Fe-P solubilizing ability

Four endophytic *Bacillus* strains, B1920, B1923, B2084, and B2088, previously isolated from maize sap, roots, and leaves,²¹ were analyzed for Fe-P solubilization in liquid culture. The strains were reactivated on PDA plates (200 gL⁻¹ of potato, 20 gL⁻¹ of dextrose and 15 gL⁻¹ of agar), using the method of streaking for obtaining pure colonies. One colony of each strain was transferred to trypticase soy broth, incubated overnight at 28 °C and subsequently 5 × 10⁷ cells mL⁻¹ of the bacterial suspension was transferred, in triplicate, to 100 mL of National Botanical Research Institute's phosphate growth (NBRIP) medium²² supplemented with FePO₄ (5 gL⁻¹) and incubated at 28 °C for 9 days at 120 rpm. After incubation, the cultures were centrifuged at 5000 × g for 10 min, the supernatant was filtered using Whatman no. 42 paper, the soluble P concentration was determined,²³ and the pH was measured.

Siderophore production

The isolates were assayed for siderophore production using chrome azurol sulphonate (CAS) agar plate assay,²⁴ modified by Pérez-Miranda et al.²⁵ Briefly, 10 µL of an overnight culture (aprox. 10⁷ cell mL⁻¹) of each isolate grown in a LB medium was transferred to an agar nutrient medium and incubated at 37 °C for 16 h. After, the siderophore production was detected using an overlay technique, in which a modified CAS medium was applied over the plates containing the cultivated microorganism and incubated in the dark at 25 °C for four days. A change in color in the overlaid medium indicates the type of siderophore produced, i.e., from blue to purple for siderophores of the catechol type, to orange for siderophores of the hydroxamate type, or to light yellow for siderophores of the carboxylate type.

Indole-acetic acid (IAA) production

Tryptophan-dependent IAA production was measured using the colorimetric method by Patten and Glick.²⁶ Briefly, single bacterial colonies of each endophytic strain were inoculated into TSB medium supplemented with 1.0 mg mL⁻¹ of L-tryptophan as an IAA precursor and incubated at 30 °C for 5 days at 100 rpm in the dark. After centrifugation at 5500 × g for 10 min, a 0.1 mL aliquot of the supernatant was mixed with 0.1 mL of Salkowsky's reagent²⁷ and incubated at room temperature for 20 min in the dark. The level of IAA present in the culture supernatant was determined in triplicate by colorimetric measurement at 540 nm and compared with a standard curve.

Greenhouse experiment

A pot experiment was carried out in a greenhouse to evaluate the effects of the *Bacillus* strains on shoot and root biomass (dry weight) and N, P, and K mobilization using pearl millet BRS1501 (*Pennisetum glaucum*) as the test crop. Pearl millet was chosen because presents fast development with short life cycle responding particularly well to nutrient deficiency with high capacity of absorption and cycling of nutrients. The experiment consisted of a factorial of three P treatments and four bacteria strains, as well as an uninoculated control treatment consisting of seed coated with sterilized mineral coal and cassava starch gum without bacteria, arranged in a completely randomized design with four replicates. The experiment was performed using 4 kg of red clay-texture oxisol, pH 5.2 (soil to water ratio of 1:2.5 (w/v)), containing 0.4 cmolc dm⁻³ Al, 2.5 cmolc dm⁻³ Ca, 0.2 cmolc dm⁻³ Mg, and 30 mg dm⁻³ K, cation exchange capacity of 11.8 cmolc dm⁻³, base saturation of 23.2%, and clay content of 74 dag kg⁻¹. The original content of available P in the soil was 2.2 mg dm⁻³.

Approximately 10 days before planting, fertilization was carried out with a nutrient solution without P²⁸ (285.8 mg dm⁻³ NH₄NO₃, 382.4 mg dm⁻³ KCl, 123.9 mg dm⁻³ (NH₄)₂SO₄, 2.9 mg dm⁻³ H₃BO₃, 7.9 mg dm⁻³ CuSO₄·5H₂O, 9.2 mg dm⁻³ MnSO₄·H₂O, 8.3 mg dm⁻³ ZnCl₂, 0.5 mg dm⁻³ (NH₄)₆Mo₇O₂₄·4H₂O), which was replaced by a half-strength of the same nutrient solution 20 days after sowing. The P sources were triple superphosphate (TSP) and Araxá rock phosphate (RP), both at 300 mg P dm⁻³ soil and a treatment with no P added. Twenty seeds of pearl millet, coated with appropriate bacterial inocula, were sown in each pot, and the seedlings were later thinned to eight plants per pot. The bacterial inocula were prepared as follows: cells from 50 mL cultures incubated for 96 h in LB medium were harvested by centrifugation at 10,000 × g for 10 min, resuspended in a 0.85% (w/v) NaCl solution and the optical densities were adjusted to 1.0 absorbance at 540 nm corresponding to 10⁸ cells mL⁻¹. Subsequently, the suspensions were added to the sterilized mineral coal (inoculum carrier) in the proportion of 10⁹ cells g⁻¹ of mineral coal. The inoculant (bacteria + mineral coal) was pelletized onto millet seeds with 4% (w/w) cassava starch gum at a final concentration of 10⁸ cells seed⁻¹.

Harvesting was carried out 50 days after planting, and roots and shoots were separated and dried in a forced air circulation oven at the temperature of 65 °C until constant weight in order to obtain dry matter. Then, the plant material was ground in a Wiley mill, and chemical analyses were conducted for determining the N P K concentration in pearl millet shoots and roots in the Laboratory of Plant Chemical Analysis at Embrapa Milho e Sorgo using ICP-OES.²⁹ The N P K content was calculated by multiplying the N P K concentration with the dry weight, which was performed separately for roots and shoots.

Statistical analyses

Data were subjected to variance analysis using the SISVAR software, version 5.6,³⁰ and the biological means were compared by the least significant difference (LSD) test at 5% probability.

Results

PGP potential in vitro

A high variability of Fe-P solubilization among the strains was observed after incubation at 28 °C (Table 1). In the absence of bacteria (control), the concentration of soluble P in the growth medium was very low, and no significant decrease was observed in pH after 9 days. Solubilized Fe-P measurements ranged from 34.25 to 52.22 mg L⁻¹, and the highest solubilization was achieved by the B1923 and B1920 strains, while the lowest was achieved by the B2084 and B2088 strains. All strains that solubilized Fe-P reduced the pH of the media compared to the control treatment, with a significant and negative correlation between the amount of soluble P and the final pH of the media ($r = -0.84$; $p < 0.05$). In addition to their phosphate-solubilizing potential, all strains, except B1920, produced the carboxylate type of siderophores when grown on CAS medium agar plates (Table 1).

The bacterial strains were able to produce IAA in the presence of L-tryptophan with a significant difference, and the yield ranged from 3.2 (B1920) to 61.6 mg L⁻¹ (B1923) (Table 1).

PGP potential in the greenhouse

Plants fertilized with soluble phosphate significantly produced the highest shoot and root biomass and N P K content. In general, no significant differences were observed between the biomass and nutrient content of plants cultivated with no P added or RP fertilization (Tables 2 and 3).

Bacterial isolates B1923, B2084, and B2088 significantly enhanced millet shoot dry biomass when grown with no P fertilization (P0) compared to the control treatment. Positive effects were also observed on shoot N P K content of plants inoculated with B2084 and B2088 strains and cultivated under P0 (Table 2). Overall, in average B2084 and B2088 significantly increased shoot biomass by around 55% and N P K content by 30, 50, and 70%, respectively, compared to control treatment with no P fertilization.

On the other hand, bacterial inoculation did not increase the shoot biomass or N P K content of plants cultivated with RP and TSP. Similar results were observed for root biomass and N P content, wherein the isolates B1923 and B2088 outperformed the others when the plants were cultivated under P0 and RP, respectively (Table 3). The relative increase in root biomass, N P content was 100, 72, and 89%, respectively, for plants inoculated with B1923 and cultivated under P0 and 66, 64, and 68%, respectively, for the same parameters in plants inoculated with B2088, but fertilized with RP, when compared to the control treatment (Fig. S1 and Table 3).

Discussion

The efficiency of phosphate solubilization by microorganisms depends on the nature of the P source and the organisms involved in the process. We selected from a previous screening *in vitro*, four *Bacillus* isolates that contrasted for Ca-P solubilization²¹ to be characterized for their capacity to

Table 1 – Isolate, local of origin, species, Fe-P solubilized, pH, IAA and siderophore production by four endophytic *Bacillus* strains after 9 days of growing at 28 °C.

Isolate	Origin	Species	Fe-P ^b (mg L ⁻¹)	pH ^c	IAA (µg mL ⁻¹)	Siderophore
Control ^a	–	–	0.00 c ^d	4.42 a	0.0 e	–
B1920	Sap	<i>Bacillus subtilis</i>	50.06 a	3.47 b	3.2 d	Negative
B1923	Sap	<i>B. pumilus</i>	52.22 a	3.50 b	61.6 a	Carboxylate
B2084	Leave	<i>B. subtilis</i>	34.56 b	3.43 b	24.4 c	Carboxylate
B2088	Root	<i>B. subtilis</i>	34.25 b	3.28 c	55.8 b	Carboxylate

^a Culture medium without bacteria.

^b P solubilized after 9 days of growing.

^c pH in water after 9 days of growing.

^d Means followed by the same letters do not differ significantly by the least significant difference (LSD) test at 5% ($p < 0.05$). The values are means of three or four replicates.

Table 2 – Shoot biomass (dry weight) and N P K content of pearl millet inoculated with endophytic *Bacillus* strains after 50 days of cultivation under greenhouse conditions with no P added (P0), Araxá rock phosphate (RP) and soluble P (TSP). Results are means SD (n = 4).

Treatment	Shoot											
	Dry weight (g pot ⁻¹)			N (mg pot ⁻¹)			P (mg pot ⁻¹)			K (mg pot ⁻¹)		
	P0	RP	TSP	P0	RP	TSP	P0	RP	TSP	P0	RP	TSP
B0 ^a	5.6 Aa ^b	7.9 Aa	19.9 Ba	277.2 Aa	308.5 Aab	472.8 Ba	20.6 Aa	24.9 Aa	37.3 Ba	240.3 Aa	355.5 Aa	848.0 Ba
B1920	6.7 Aab	7.1 Aa	17.5 Ba	302.6 Aab	317.2 Aab	451.8 Ba	24.7 Aab	24.3 Aa	34.0 Ba	298.1 Aab	362.2 Aa	764.5 Ba
B1923	7.8 Abc	8.3 Aa	19.7 Ba	274.2 Aa	386.8 Bb	427.8 Ba	25.6 Aab	29.6 Aba	34.3 Ba	356.0 Aab	390.9 Aa	881.6 Ba
B2084	8.7 Ac	7.9 Aa	19.2 Ba	351.4 Ab	354.9 Aab	418.3 Aa	29.5 ABb	24.9 Aa	35.0 Ba	406.9 Ab	379.8 Aa	828.2 Ba
B2088	8.6 Ac	8.6 Aa	19.9 Ba	362.2 Ab	290.1 Aa	387.7 Aa	30.8 ABb	24.7 Aa	34.5 Ba	405.8 Ab	396.0 Aa	797.7 Ba

^a B0 – uninoculated control.

^b Means followed by the same lower case letters in a column and capital letters on the lines do not differ significantly by the LSD test ($p < 0.05$).

Table 3 – Root biomass (dry weight) and N P K content of pearl millet inoculated with endophytic *Bacillus* strains after 50 days of cultivation under greenhouse conditions with no P added (P0), Araxá rock phosphate (RP) and soluble P (TSP). Results are means SD (n = 4).

Treatment	Root											
	Dry weight (g pot ⁻¹)			N (mg pot ⁻¹)			P (mg pot ⁻¹)			K (mg pot ⁻¹)		
	P0	RP	TSP	P0	RP	TSP	P0	RP	TSP	P0	RP	TSP
B0 ^a	1.2 Aa ^b	1.5 Aa	3.5 Ba	26.4 Aa	28.4 Aa	49.3 Ba	1.9 Aa	2.2 Aa	4.2 Ba	22.2 Aa	23.1 Aa	43.2 Ba
B1920	1.7 ABab	1.3 Aa	2.9 Ba	35.9 Aab	27.6 Aa	45.4 Aa	2.6 Aab	1.9 Aa	4.2 Aa	28.3 Aa	24.0 Aa	35.0 Aa
B1923	2.4 Ab	1.5 Aa	3.6 Ba	45.5 Ab	31.3 Aa	56.7 Ba	3.6 ABb	2.3 Aa	4.78 Ba	29.9 Aa	27.9 Aa	45.9 Aa
B2084	1.8 Aab	1.7 Aa	3.6 Ba	37.3 Aab	37.5 Aab	56.1 Ba	2.7 Aab	2.7 Aab	4.1 Aa	29.7 Aa	32.7 Aa	45.3 Ba
B2088	1.7 Aab	2.5 Bb	3.7 Ca	35.6 Aab	46.6 ABb	55.5 Ba	2.6 Aab	3.7 ABb	4.7 Ba	34.4 Aa	38.4 Aa	43.3 Aa

^a B0 – uninoculated control.

^b Means followed by the same lower case letters in a column and capital letters on the lines do not differ significantly by the LSD test ($p < 0.05$).

solubilize Fe-P, that is the most abundant and least soluble P fraction in Brazilian Oxisols.^{3,4} In addition, we assessed siderophore and IAA production, followed by a greenhouse test to assess their effect on plant growth promotion.

Our results of soluble P released by *Bacillus* grown on a medium containing FePO₄ varied from 34.25 to 52.22 mg L⁻¹ and were lower than the previously reported values for Ca-P solubilization by the same isolates, which ranged from 112 to 179 mg L⁻¹.²¹ Similar results, reported previously with different bacterial species, have shown that Fe-P and Al-P are less soluble than Ca-P.^{31,32} The highest Fe-P solubilization has been documented in rhizobacteria and fungi;^{4,32,33} however, our isolates showed higher solubilization than reported for endophytic bacteria isolated from peanuts, that ranged from

13.9 ± 0.3 to 37.4 ± 2.0 mg L⁻¹.¹² In our study, the inoculated strains decreased the growth medium pH and simultaneously increased the concentration of soluble phosphate obtained from FePO₄. Abreu et al.²¹ measured organic acids secreted by the same bacteria, and the results suggested that culture medium acidification might be one of the mechanisms involved on Ca-P solubilization. These organic acids can chelate cations associated with phosphate, thus making P available to plants.^{8,34,35} However, the mechanisms behind Fe-P solubilization remain largely unknown. Highly weathered Brazilian soils under different agrosystems predominantly consist of Fe and Al oxides, independent of the vegetation, facilitating the formation of Fe-P and Al-P. This makes P often low in the soil, limiting crop yield,⁴ emphasizing the

importance of screening for soil microorganisms that exhibit a Fe-P solubilization ability.

The effects of inoculating the four bacterial strains onto pearl millet grown with different P-sources under greenhouse conditions varied, depending on P treatment and the strain used. Compared to uninoculated controls grown with no P added, the strains B1923, B2084, and B2088 promoted increase in dry weight of shoots, and two of them, B2084 and B2088, increased shoot N P K content. Interestingly, B2084 and B2088 produced high concentrations of gluconic acid (324 and 171 mM, respectively) and solubilized phosphate, with a reduction of pH, when cultivated in Ca-P²¹ or Fe-P (Table 1) media. Gluconic acid production has been frequently described as the most efficient mechanism related to P solubilization by bacteria. For instance, *Pisum sativum* inoculated with strains of endophytic *Pseudomonas fluorescens*, which is capable of producing gluconic acid (14–169 mM) and solubilizing phosphate, presented higher growth compared to uninoculated plants.³⁶ These strains probably exude gluconic acid into the rhizosphere of the inoculated plants, thus playing an important role in the solubilization of natural P fixed in soil particles and allowing the plant to subsequently assimilate the soluble P.

On the other hand, although the inoculation of plants with strains B1923, B2084, and B2088 and cultivated with Araxá RP increased shoot dry weight, this effect was not statistically significant, indicating that the amount of P solubilized by these bacteria was probably not sufficient to significantly increase the shoot parameters evaluated in our short-term experiment. Long-term field experiments are likely to show more clearly the effect of RP solubilization by these bacteria on P accumulation in grain and shoot biomass.

In our study, two *Bacillus* strains also caused a significant increase in root dry weight and N P content in plants cultivated in soil without P added (B1923) or amended with RP (B2088). The difference between the performance of the two isolates in the greenhouse experiment was probably a consequence of their different P-solubilizing mechanisms, measured *in vitro*, since B2088 is an efficient Ca-P solubilizer²¹ and B1923 showed higher Fe-P solubilization (Table 1). This characteristic could be important in the selection of bacteria to be used as inoculants for plants cultivated in different soils or soils amended with RP. It is well known that the rate of RP solubilization is influenced by different physicochemical characteristics that depend on the source material and particle size. Araxá is an RP of igneous or metamorphic origin that exhibits a high crystallization level and low citric acid solubility,³⁷ which makes it difficult to solubilize, particularly in the first year of cultivation. In addition, fluoride present in Araxá RP limited the solubilization by *Aspergillus niger* by negatively affecting metabolic processes involved in phosphate solubilization, such as the growth of the fungus, its citric acid production, and medium acidification.³⁸

Most of the Fe-P-solubilizing strains also produced siderophores (Table 1), which are iron-binding compounds that can chelate ferric iron and thereby make it available for microbial and plant cells. It is known that siderophores have the potential for plant pathogen biocontrol,^{13,39} as well as the potential to solubilize iron from minerals, thus increasing the

availability of P to plants.^{40,41–43} Recently, Ghosh et al. (2016)⁴⁴ reported high siderophore production by three *Burkholderia* strains that produced soluble phosphate when ferric phosphate was used as the sole phosphate source. However, the specific contribution of siderophores to P solubilization mechanisms in soil is yet to be elucidated.⁴⁵

In addition to the siderophore-producing ability, an *in vitro* screening revealed that all *Bacillus* isolates, when grown with tryptophan in the culture medium, were able to produce IAA, in a range between 3.2 and 61.6 $\mu\text{g mL}^{-1}$, revealing a substantial variability among isolates (Table 1). Other species of *Bacillus* and *Lactobacillus* presented similar results, with reported IAA production between 30 and 60 $\mu\text{g mL}^{-1}$ ¹⁰ and 0.81 and 86.82 $\mu\text{g mL}^{-1}$.¹³ IAA production is widespread among bacteria and has been used as a criterion for selecting effective PGP bacteria.⁴⁶ Bacteria that produce IAA and promote growth were described for different crops such as maize, peanut, wheat, soybean, and rice.^{46–49} In general, high levels of bacterial IAA are associated with lateral and adventitious root formation, while low IAA levels often stimulate root elongation.⁵⁰ An extended root surface combined with a more efficient P solubilization ability improves nutrient acquisition and water uptake. IAA-producing bacteria also stimulate shoot growth; however, it is not clear whether IAA has a direct effect on shoot growth or an indirect effect, i.e., by stimulating the root system and consequently enhancing nutrient and water uptake.⁵¹

Most PGPB endophytes are facultative, changing between the free-living and endophytic stages depending on different factors such as the soil, microbial competitors, and plant nutrients.⁵² This facultative endophytic lifestyle enables multi-traits PGP, resulting in more competitive bacteria than rhizospheric microorganism as they can be inoculated in seeds and spread systemically through the plant reducing the need of continuous inoculations.⁵³

In conclusion, we observed a positive effect of the inoculation of three endophytic *Bacillus* (B1923, B2084 and B2088) on pearl millet growth and nutrient uptake, indicating that these strains have potential to be used as plant inoculants in highly weathered tropical soils. B1923 strain was able to produce IAA and solubilize Fe-P *in vitro* and this trait was expressed under phosphate limiting conditions in the soil (no P added) resulting in enhanced shoot and root dry weight and root N P content of pearl millet plants. Strains B2084 and B2088, despite of solubilize less Fe-P than B1923, showed positive performance on biomass production and accumulation of nutrients N P K in the shoot, which represents the most relevant response of these microorganisms in the productive environment. Based on this, we indicated that the strains B2084 and B2088 have higher potential to be microbial biofertilizer candidates for commercial applications in the absence of fertilization.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by Fapemig (Grant No. APQ-01867-14), MPCPAgro (CNPq 465133/2014-4, Capes) and Embrapa (Grant No. 01.13.05.001.02-04).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.bjm.2018.06.005](https://doi.org/10.1016/j.bjm.2018.06.005).

REFERENCES

- Santos HG, Jacomine PKT, Anjos LHC, et al. *Sistema Brasileiro de Classificação de Solos*. 3. ed. Brasília, DF: Embrapa; 2013:353.
- Fageria NK, Baligar VC. Improving nutrient use efficiency of annual crops in Brazilian acid soils for sustainable crop production. *Commun Soil Sci Plant Anal*. 2001;32:1303–1319.
- Ahn PM. Tropical soils and fertilizer use. In: *Intermediate Tropical Agriculture Series*. Malaysia: Longman Scientific and Technical; 1993:3.
- Barroso CB, Nahas E. The status of soil phosphate fractions and the ability of fungi to dissolve hardy soluble phosphates. *Appl Soil Ecol*. 2005;29:73–83.
- Withers PJA, Rodrigues M, Soltangheisi A, et al. Transitions to sustainable management of phosphorus in Brazilian agriculture. *Sci Rep*. 2018;8:2537.
- Novais RF, Smyth TJ. *Fósforo em Solo e Planta em Condições Tropicais*. Viçosa: UFV; 1999:399.
- Singh H, Reddy MS. Effect of inoculation with phosphate solubilizing fungus on growth and nutrient uptake of wheat and maize plants fertilized with rock phosphate in alkaline soils. *Eur J Soil Biol*. 2011;47:30–34.
- Gomes EA, Silva UC, Marriel IE, Oliveira CA, Lana UGP. Rock phosphate solubilizing microorganisms isolated from maize rhizosphere soil. *Rev Bras Milho Sorgo*. 2014;13:69–81.
- Collavino MM, Sansberro PA, Mroginski LA, Aguilar OM. Comparison of in vitro solubilization activity of diverse phosphate-solubilizing bacteria native to acid soil and their ability to promote *Phaseolus vulgaris* growth. *Biol Fertil Soils*. 2010;46:727–738.
- Mohite B. Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *J Soil Sci Plant Nutr*. 2013;13:638–649.
- Arruda L, Beneduzi A, Martins A, et al. Screening of rhizobacteria isolated from maize (*Zea mays* L) in Rio Grande do Sul State (south Brazil) and analysis of their potential to improve plant growth. *Appl Soil Ecol*. 2013;63:15–22.
- Anzuay MS, Ciancio MGR, Ludueña LM, et al. Growth promotion of peanut (*Arachis hypogaea* L.) and maize (*Zea mays* L.) plants by single and mixed cultures of efficient phosphate solubilizing bacteria that are tolerant to abiotic stress and pesticides. *Microbiol Res*. 2017;199:98–109.
- Wahyudi AT, Astuti RP, Widayawati A, Meryandini A, Nawangsih AA. Characterization of *Bacillus* sp. strains isolated from rhizosphere of soybean plants for their use as potential plant growth for promoting Rhizobacteria. *J Microbiol Antimicrob*. 2011;3:34–40.
- Bahadır PS, Liaqat F, Eltem R. Plant growth promoting properties of phosphate solubilizing *Bacillus* species isolated from the Aegean Region of Turkey. *Turk J Bot*. 2018;42:1–14.
- Malfanova N, Franzil L, Lugtenberg B, Chebotar V, Ongena M. Cyclic lipopeptide profile of the plant-beneficial endophytic bacterium *Bacillus subtilis* HC8. *Arch Microbiol*. 2012;194:893–899.
- Hardoim PR, Van Overbeek LS, Van Elsas JD. Properties of bacterial endophytes and their proposed role in plant growth. *Trends Microbiol*. 2008;16:463–547.
- Bacon CW, Hinton DM. *Bacillus mojavensis*: its endophytic nature, the surfactants, and their role in the plant response to infection by *Fusarium verticillioides*. In: Maheshwari DK, ed. *Bacteria in Agrobiology: Plant Growth Responses*. Springer-Verlag Berlin; 2011:21–39.
- Vessey JK. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil*. 2003;255:571–586.
- Bjelić D, Marinković J, Tintor B, Mrkovački N. Antifungal and plant growth promoting activities of indigenous rhizobacteria isolated from maize (*Zea mays* L.) rhizosphere. *Commun Soil Sci Plant Anal*. 2018, <http://dx.doi.org/10.1080/00103624.2017.1421650>.
- Lu X, Zhou D, Chen X, Zhang J, Huang H, Wei L. Isolation and characterization of *Bacillus altitudinis* JSCX-1 as a new potential biocontrol agent against *Phytophthora sojae* in soybean [*Glycine max* (L.) Merr.]. *Plant Soil*. 2017;413:1–14.
- Abreu CS, Figueiredo JE, Oliveira CA, et al. Maize endophytic bacteria as mineral phosphate solubilizers. *Genet Mol Res*. 2017, <http://dx.doi.org/10.4238/gmr16019294>.
- Nautiyal CS. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiol Lett*. 1999;170:265–270.
- Murphy J, Riley JP. A modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta*. 1962;27:31–36.
- Schwyn B, Neilands JB. Universal chemical assay for the detection and determination of siderophores. *Analytical Biochem*. 1987;160:47–56.
- Perez-Miranda S, Cabriol N, George-Téllez R, Zamudio-Rivera LS, Fernández FJ. O-CAS, a fast and universal method for siderophore detection. *J Microbiol Methods*. 2007;70:127–131.
- Patten CL, Glick BR. Bacterial biosynthesis of indole-3-acetic acid. *Canadian J Microbiol*. 1996;42:207–220.
- Loper JE, Schroth MN. Influence of bacterial sources of indole-3-acetic acid on root elongation of sugar beet. *Phytopathology*. 1986;76:386–389.
- Somasegaran P, Hoben HJ. *Methods in Legume – Rhizobium Technology*. University of Hawaii Niftal. Biological Nitrogen Fixation; 1985:510.
- Nogueira ARA, Souza GB. *Manual de Laboratórios: Solo, Água, Nutrição Vegetal, Nutrição Animal e Alimentos*. São Carlos: Embrapa Pecuária Sudeste; 2005:313.
- Ferreira DF. Sisvar: a computer statistical analysis system. *Ciênc Agrotec*. 2011;35:1039–1042.
- Bashan Y, Kamnev AA, De-Bashan LE. Tricalcium phosphate is inappropriate as a universal selection factor for isolating and testing phosphate-solubilizing bacteria that enhance plant growth: a proposal for an alternative procedure. *Biol Fertil Soils*. 2013;49:465–479.
- Rinu K, Malviya MK, Sati P, Tiwari SC, Pandey A. Response of cold-tolerant *Aspergillus* spp. to solubilization of Fe and Al phosphate in presence of different nutritional sources. *ISRN Soil Sci*. 2013. Article 598541.
- Spagnoletti FN, Tobar NE, Fernández Di Pardo A, Chiochio VM, Lavado RS. Dark septate endophytes present different potential to solubilize calcium, iron and aluminum phosphates. *Appl Soil Ecol*. 2017;111:25–32.
- Barroso CB, Nahas E. Solubilização do fosfato de ferro em meio de cultura. *Pesq Agropec Bras*. 2008;43:529–535.
- Oliveira CA, Alves VM, Marriel IE, et al. Phosphate solubilizing microorganisms isolated from rhizosphere of

- maize cultivated in an oxisol of the Brazilian Cerrado Biome. *Soil Biol Biochem.* 2009;41:1782–1787.
36. Oteino N, Lally RD, Kiwanuka S, et al. Plant growth promotion induced by phosphate solubilizing endophytic *Pseudomonas* isolates. *Front Microbiol.* 2015. Article 745.
 37. Kliemann HJ, Lima DV. Eficiência agrônômica de fosfatos naturais e sua influência no fósforo disponível em dois solos de cerrado. *Pesq Agropec Trop.* 2001;31:111–119.
 38. Mendes GO, Vassilev NB, Bonduki VHA, Silva IR, Ribeiro JI, Costa MD. Inhibition of *Aspergillus niger* phosphate solubilization by fluoride released from rock phosphate. *Appl Environ Microbiol.* 2013;79:4906–4913.
 39. Ahmed E, Holmström SJM. Siderophores in environmental research: roles and applications. *Microb Biotechnol.* 2014;7:196–208.
 40. Vassilev N, Medina A, Azcón R, Vassileva M. Microbial solubilization of rock phosphate on media containing agro-industrial wastes and effect of the resulting products on plant growth and P uptake. *Plant Soil.* 2006;287: 77–84.
 41. Caballero-Mellado J, Onofre-Lemus J, Estrada-de los Santos P, Martínez-Aguilar L. The tomato rhizosphere, an environment rich in nitrogen-fixing Burkholderia species with capabilities of interest for agriculture and bioremediation. *Appl Environ Microbiol.* 2007;73:5308–5319.
 42. Hamdali H, Bouizgarne B, Hafidi M, Lebrihi A, Virolle MJ, Ouhdouch Y. Screening for rock phosphate-solubilizing Actinomycetes from Moroccan phosphate mines. *Appl Soil Ecol.* 2008;38:12–19.
 43. Santos S, Neto IFF, Machado MD, Soares HMVM, Soares EV. Siderophore Production by *Bacillus megaterium*: effect of growth phase and cultural conditions. *Appl Biochem Biotechnol.* 2014;172:549–560.
 44. Ghosh R, Barman S, Mukherjee R, Mandal NC. Role of phosphate solubilizing *Burkholderia* spp. for successful colonization and growth promotion of *Lycopodium cernuum* L. (Lycopodiaceae) in lateritic belt of Birbhum district of West Bengal, India. *Microbiol Res.* 2016;183:80–91.
 45. Sharma SB, Sayyed RZ, Trivedi MH, Gobi TA. Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *SpringerPlus.* 2013;2:587.
 46. Ali B, Sabri AN, Ljung K, Hasnain S. Auxin production by plant associated bacteria: impact on endogenous IAA content and growth of *Triticum aestivum* L. *Lett Appl Microbiol.* 2009;48:542–547.
 47. Spaepen S, Dobbelaere S, Croonenborghs A, Vanderleyden J. Effects of *Azospirillum brasilense* indole-3-acetic acid production on inoculated wheat plants. *Plant Soil.* 2008;312:15–23.
 48. Mehnaz S, Kowalik S, Reynolds B, Lazarovits G. Growth promoting effects of corn (*Zea mays*) bacterial isolates under greenhouse and field conditions. *Soil Biol Biochem.* 2010;42:1848–1856.
 49. Majeed A, Abbas MK, Hameed S, Imran A, Rahim N. Isolation and characterization of plant growth-promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion. *Soil Biol Biochem.* 2015;42: 1848–1856.
 50. Patten CL, Glick BR. Role of *Pseudomonas putida* indoleacetic acid in development of the host plant root system. *Appl Environ Microbiol.* 2002;68:3795–3801.
 51. Dodd IC, Zinovkina NY, Safronova VI, Belimov AA. Rhizobacterial mediation of plant hormone status. *Ann Appl Biol.* 2010;157:361–379.
 52. Miliute I, Buzaitė O, Baniulis D, Stanys V. Bacterial endophytes in agricultural crops and their role in stress tolerance: a review. *Zemdirbyste - Agriculture.* 2015;102:465–478.
 53. Johnston-Monje D, Raizada MN. Conservation and diversity of seed associated endophytes in *Zea* across boundaries of evolution, ethnography and ecology. *PLoS ONE.* 2011;6:e20396.