

## Hepatitis E virus: new faces of an old infection

Mar Riveiro-Barciela,\* Francisco Rodríguez-Frías,\*\*,\*\*\* María Buti\*,\*\*\*

\* Liver Unit, Department of Internal Medicine, Vall d'Hebron Hospital, Autonomous University of Barcelona, Barcelona, Spain.

\*\* Department of Biochemistry, Vall d'Hebron Hospital. Barcelona. Spain.

\*\*\* Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBERehd), Instituto Carlos III, Spain.

### ABSTRACT

Hepatitis E virus is one of the most common causes of acute hepatitis worldwide, with the majority of cases occurring in Asia. In recent years, however, an increasing number of acute and chronic hepatitis E virus infections have been reported in industrialized countries. The importance of this infection resides in the associated morbidity and mortality. In acute cases, a high mortality rate has been reported in patients with previously undiagnosed alcoholic liver disease. Hepatitis E infection can become chronic in immunocompromised patients, such as solid organ transplant recipients, patients receiving chemotherapy, and HIV-infected patients, and lead to the development of hepatic fibrosis and cirrhosis. Hence, treatment strategies involving reductions in immunosuppressive regimens and therapy with ribavirin or peg-interferon have been evaluated. In terms of prevention, a promising new vaccine was recently licensed in China, although its efficacy is uncertain and potential adverse effects in risk groups such as chronic liver disease patients and pregnant women require investigation. In conclusion, physicians should be aware of hepatitis E as a cause of both acute and chronic hepatitis in immunocompromised patients. The best treatment option for HEV infection remains to be defined, but both ribavirin and peg-interferon may have a role in therapy for this condition.

**Key words.** Epidemiology. Diagnosis. Manifestations. Treatment. Vaccine.

### INTRODUCTION

Hepatitis E virus (HEV) is a common cause of acute hepatitis infection worldwide.<sup>1</sup> In developed countries, however, HEV is responsible for a small number of cases, being hepatitis A virus the most prevalent etiology.<sup>2-3</sup> HEV infection was first described in 1980 through analysis of blood samples from patients affected by an acute hepatitis epidemic in New Delhi (India) during 1955 and 1956, spread by contaminated water.<sup>4</sup> The causal agent in this epidemic, the “enteric non-A hepatitis virus” was identified in 1983<sup>5</sup> and named E, because of its enteric and endemic characteristics.

Although most HEV hepatitis cases occurring in industrialized countries are imported from areas where this infection is endemic, an increasing number of native cases have been reported. HEV infection in developed countries is generally caused by genotype 3 or 4, and is mainly diagnosed by serological testing in patients older than 60 years with unexplainable acute hepatitis.<sup>6</sup> In the last few years, HEV has also been identified as a cause of chronic liver disease in immunocompromised patients, such as solid organ transplant recipients,<sup>7-23</sup> individuals infected by the human immunodeficiency virus (HIV),<sup>24-34</sup> and patients with hematologic disorders receiving chemotherapy.<sup>35-37</sup> In addition, it has been found to cause acute decompensation in patients with chronic liver disease.<sup>38-40</sup>

HEV is a non-enveloped virus 27 to 34 nm in size, with a 7.2-kb positive-sense single-stranded RNA genome. It contains 3 open reading frames (ORFs) as well as 5' and 3' untranslated regions: ORF1 encodes non-structural proteins, ORF2 encodes the capsid protein, and ORF3 encodes a small phospho-

Correspondence and reprint request: Dr. María Buti. Psg.  
Vall d'Hebrón 119-129. Barcelona, 08035, Spain.  
Tel: (+34) 93 274 61 40. Fax: (+34) 93 274 60 68  
E-mail: mbuti@vhebron.net

*Manuscript received: June 30, 2013.*

*Manuscript accepted: July 25, 2013.*




Figure 1. Phylogenetic tree of HEV.

protein.<sup>41-42</sup> HEV is the unique member of the *Hepevirus* genus and *Hepeviridae* family.<sup>2</sup> Its resistance to inactivation by the acid and slightly alkaline conditions of the intestinal tract explains why its transmission mainly occurs via the fecal-oral route.

HEV comprises five genotypes, all belonging to one serotype.<sup>43</sup> Genotypes 1 and 2 are strictly human and are associated with large outbreaks transmitted by contaminated water in countries where HEV infection is endemic, such as Asia for genotype 1 and Africa and Mexico for genotype 2.<sup>44</sup> Genotype 3 and 4 can infect humans, but also other mammals, mainly pigs and deer. Genotype 5 is of avian origin and has only been described in birds. Figure 1 shows the phylogenetic tree of rabbit HEV.

## EPIDEMIOLOGY CHANGE IN DEVELOPED COUNTRIES

The first cases of HEV infection in industrialized countries were imported from developing areas in Asia and Africa. Since 1997, when the similarity between swine and human HEV strains was first described,<sup>45</sup> an increasing number of native HEV infections have been reported in the USA, Europe,

and developed Asian-Pacific countries (Japan, Taiwan, Hong Kong, Australia). Although most of these cases have involved genotype 3 strains, some genotype 4 infections have been recently reported in France.<sup>46-47</sup>

In contrast to tropical countries, where HEV infection is fecal-orally transmitted by contaminated water, in industrialized countries, zoonotic HEV transmission is considered the main source of infection. There are also reported cases of transmission by blood transfusion<sup>48-51</sup> and liver transplantation.<sup>52</sup> HEV3 and HEV4 infections have been related to ingestion of raw or undercooked meats, such as pig liver sausages or game meats,<sup>46,53,54</sup> but the complete spectrum of animals that are HEV reservoirs is unknown. In the last years, several studies have characterized new HEV genotypes in rabbits in France,<sup>55</sup> rats in Germany<sup>56,57</sup> and the United States,<sup>58</sup> farm rabbits in China,<sup>59,60</sup> and wild boars in Japan.<sup>61,62</sup>

A study performed in Spain revealed a higher prevalence of anti-HEV antibodies in swine industry workers in comparison to the general population (18.8 vs. 4.1%,  $p = 0.03$ ).<sup>63</sup> These data suggest that swine HEV infection promotes a high prevalence of HEV antibodies in individuals directly and frequently exposed to these animals, and exemplifies the zoonotic feature of this infection. HEV is endemic in swine and there is a high anti-HEV prevalence worldwide (almost 100% in the USA and Mexico, 90% in New Zealand, 46% in Laos, and 98% in Spain); furthermore, it shares genotypes 3 and 4, which are responsible for human infections in non-endemic areas.<sup>5</sup>

A study performed in Barcelona (Spain) showed that detection of HEV-RNA in urban sewage samples has remained stable (around 30%) despite overall improvements in sanitation. However, during the same 5- to 10-year period, the presence of hepatitis A virus dramatically decreased (from 57.4 to 3.1%). This observation highlights the importance of animal reservoirs of HEV, which act as external sources of infection.<sup>64</sup> Studies on the presence of HEV in the pork food chain, revealed a 10 and 6% HEV detection rate in pork sausage at point of sale in the United Kingdom<sup>65</sup> and Spain,<sup>66</sup> respectively, and a rate in meat samples of 3% in the Czech Republic and 6% in Italy.<sup>66</sup> The elevated detection of HEV in 60% of floor and working surfaces and 57 to 71% of hands and aprons of workers dissecting pigs, emphasizes the high risk for workers and highlights the need to improve preventive measures in these industries to avoid HEV transmission.<sup>66</sup>

## HEPATITIS E DIAGNOSIS: CLINICAL PRESENTATION AND EXTRAHEPATIC MANIFESTATIONS


In endemic countries, HEV infection by genotypes 1 and 2 mainly affects children and young adults.<sup>4</sup> The infection is more common in men, and the associated mortality rate in these areas is 1 to 15%. During outbreaks of this infection, pregnant women are particularly susceptible and show high mortality and morbidity rates (19 vs. 2.1% in non-pregnant women and 2.8% in men).<sup>67</sup> HEV epidemics are also related to a high risk of prematurity and perinatal mortality.<sup>68</sup>

In contrast, in developed countries, HEV represents a minority of acute hepatitis cases, which are mainly attributed to genotypes 3 and 4. HEV infection in these countries essentially affects persons older than 60 years.<sup>6</sup> The clinical presentation is similar to that seen in endemic areas, but the percentage of icteric hepatitis seems to be higher. Older age and pre-existing chronic liver disease may explain the poorer prognosis in contrast to endemic countries.<sup>69</sup>

Until recently, hepatitis E (as hepatitis A) was considered a self-limiting disease, whose worst outcome was acute liver failure. However, since 2008, an increasing number of chronic infections by HEV have occurred in immunocompromised patients, such as HIV-infected individuals,<sup>24-34</sup> patients with hematologic disorders receiving chemotherapy,<sup>35-37</sup>

and solid organ transplant recipients.<sup>7-23</sup> The importance of these chronic infections is related to the development of hepatic fibrosis and the risk of progression to cirrhosis.<sup>70-72</sup> The mechanisms of acquisition of the infection in these patients are uncertain. Reactivation of a latent infection seems not to play a role.<sup>73,74</sup> In a study performed in solid-organ transplant recipients in France, the only independent factor associated with HEV infection was consumption of game meat,<sup>75</sup> suggesting that zoonotic transmission is the main mode of acquisition.

Infection by HEV genotype 3 in immunosuppressed patients has been related to several extrahepatic manifestations, which are summarized in figure 2. The first neurological manifestation described in relation to HEV infection was an isolated case of Guillain-Barré syndrome.<sup>76</sup> Between 2004 and 2009, up to 5.5% of neurologic complications were reported in a series of 126 patients in two hospitals in the United Kingdom and France.<sup>77</sup> These manifestations included inflammatory polyradiculopathy, Guillain-Barré syndrome, bilateral brachial neuritis, encephalitis, and ataxia/proximal myopathy. A case of Guillain-Barré syndrome associated with severe necrotizing myositis was recently reported in a liver transplant patient with acute HEV infection, who recovered after treatment with ribavirin.<sup>78</sup> As to renal manifestations, membranoproliferative glomerulonephritis and relapses of IgA nephropathy have been related to HEV infection in kidney and liver transplant patients.<sup>18</sup> Most of these patients



**Figure 2.** Extrahepatic manifestations of hepatitis E infection.

Table 1. Chronic hepatitis E cases treated with ribavirin.

Author	Treated cases	Age	Gender	Genotype	Immunosuppression (aetiology)	Ribavirin dosage (cumulative dose)	Days to viral clearance	Ribavirin side effects	Outcome
Mallet, et al. France 2010 <sup>13</sup>	2	40	Male	3f	Kidney and pancreas transplant.	200-800 mg/24 h/84 d (NR)	28 days	Anaemia	VR (3 months)
		57	Female	3c	Idiopathic CD4 T lymphopenia and Ig G primary deficiency.	600 mg/24 h/84 d (50,400 mg)	28 days	None	VR (2 months)
Kamar, et al. France 2010 <sup>15</sup>	6	53	Male	3f	Kidney transplant	200-400/12h/90d (36,000-72,000 mg)	15 days	Anaemia	Relapse (1 month)
		56	Male	3f			60 days	None	VR (6 months)
		48	Male	3f			15 days	None	VR (6 months)
		48	Male	3c			21 days	None	Relapse (2 months)
		55	Female	3f			15 days	None	VR (6 months)
		70	Male	3c			15 days	Anaemia	VR (6 months)
Chaillon, et al. France 2011 <sup>16</sup>	1	57	Male	3c	Heart transplant	Dose not reported for 3 months	30 days	Anaemia	VR (4 months)
Dalton, et al. United Kingdom 2011 <sup>26</sup>	1	48	Male	3a	HIV infection	500 /24 h/42 d 1,000/24 h/42 d (63,000 mg)	NR	Marrow suppression	VR (9 months)*
Alric, et al. France 2011 <sup>36</sup>	1	77	Female	3c	Myelomonocytic leukemia and corticosteroid therapy	600/24 h/90 d (54,000 mg)	14 days	None	VR (> 6 months)
Kamar, et al. France 2012 <sup>18</sup>	9	NR	NR	3	Kidney or liver transplant	600mg/24h/90d (54000 mg)	NR	NR	VR (6 months)
de Niet, et al. The Netherlands 2012 <sup>19</sup>	1	51	Male	3	Kidney transplant and etanercept for Bechterew's disease	800 mg/12 h/90 d (144,000 mg)	60 days	None	VR (6 months)
Koning, et al. The Netherlands 2013 <sup>20</sup>	4	51	Male	3	Heart transplant	200/24 h/90 d (18,000 mg)	NR	NR	VR (NR)
		62	Male			200-400/24 h/240 d (NR)	NR	NR	VR (NR)
		49	Male			200-400/24 h/90 d (NR)	No clearance	NR	Relapse
		38	Male			200-800/24 h/270 d (NR)	NR	NR	VR (NR)
Riezebos-Brilman, et al. The Netherlands 2013 <sup>21</sup>	2	49	Male	3	Lung transplant	400 mg/12 h/120 d (96,000 mg)	42-49 days	None	VR (> 6 months)
		41	Male	3	Lung transplant	400 mg/12 h/120d (96,000 mg)	42-49 days	None	VR (> 6 months)
Pischke, et al. Germany 2013 <sup>22</sup>	11	65	Male	3	Kidney transplant	600-1,000 mg/24 h/150 d (90,000-1500,000 mg)	30 days	1/11 lung graft failure	VR (> 2 years)
		50	Male	3	Kidney transplant		60 days	5/11 anaemia	VR (> 5 months)
		40	Male	3	Kidney transplant		60 days		VR (> 4 months)
		54	Male	3	Kidney transplant		30 days		VR (> 4 months)
		50	Female	3	Heart transplant		60 days		VR (> 2 years)
		66	Male	3	Heart transplant		30 days		VR (> 2 years)
		57	Male	3	Heart transplant		30 days		VR (> 2 years)
		58	Male	3	Heart transplant		No clearance		Relapse and died
		48	Female	3	Lung transplant		60 days		VR (> 2 months)
		56	Female	3	Lung transplant		60 days		VR (> 7 months)
		32	Male	3	Lung transplant		No clearance		Relapse and died
Alric, et al. France 2013 <sup>23</sup>	18	NR	NR	NR	Solid organ recipients or haematological malignancy	600-800 mg/24 h/84 d (50,400-67,200 mg)	NR (at week 8 all patients had negative RNA)	Anaemia (no more data available)	13/18 VR (NR) 5/18 relapse

D: days. HEV: hepatitis E virus. NR: not reported. RNA: ribonucleic acid. VR: virologic response. \*Vey low level of HEV RNA was detected intermittently in both plasma and stool.

had cryoglobulinemia that became negative after spontaneous HEV clearance or after antiviral treatment. Kidney function improved and proteinuria decreased after HEV clearance. As in hepatitis A, hepatitis E infection has been related to hematological disorders, such as severe thrombocytopenia<sup>79</sup> and pure red cell aplasia.<sup>80,81</sup>

### DIAGNOSIS OF HEV: LIMITATIONS

Membership of the four main HEV genotypes in a single serotype has facilitated the development of enzyme-linked immunoassays (ELISA) that are useful for universal diagnosis. These assays can detect specific IgG and IgM antibodies, regardless of HEV genotype. IgM antibodies appear in the acute phase of the infection, early at the end of the incubation period or when jaundice appears. IgM antibodies are an adequate marker for acute infection, as they remain detectable for 4 to 5 months.<sup>82</sup> However, it should be noted that detection of HEV IgM presents specificity (78-98%) and sensitivity (72-98%) problems. HEV-RNA detection is the best marker for diagnosing acute hepatitis E infection,<sup>83</sup> but HEV-RNA is only present during the short duration of the viremia period (around 2 weeks in serum and 4-12 weeks in stools). Thus, a lack of HEV RNA does not exclude the diagnosis of acute hepatitis E. Moreover, the majority of reverse-transcriptase PCR techniques for HEV-RNA determination are neither marketed nor standardized; hence, their interpretation and comparison are difficult.

Laboratory diagnosis of acute hepatitis E is based on the presence of HEV IgM in serum and/or detection of HEV-RNA in serum or stools. In acute infection, HEV IgM and even IgG can be detected, although detection of IgG antibodies alone indicates past infection. Therefore, IgG detection is useful for prevalence studies. Regarding immunosuppressed patients, it is essential to determine HEV-RNA because the antibodies may appear late or even be absent.

### MANAGEMENT OF ACUTE AND CHRONIC HEPATITIS E: WHEN AND HOW TO TREAT?

The majority of acute hepatitis E infections resolve spontaneously and therefore, treatment is not indicated. The factor associated with a risk of developing chronic hepatitis E is immunosuppression. Therefore, the first step in the management of chronic hepatitis E in immunocompromised patients is a reduction in the immunosuppressive therapy regimen, if it is possible.<sup>84</sup> There is no specific recommended therapy for hepatitis E. However, two drugs have been used: pegylated interferon-alpha 2a or 2b and ribavirin. This last seems to be the most useful, or at least the most widely used drug for treating both acute and chronic hepatitis E. However the number of patients treated in this way is still small and very heterogenous. Ribavirin has proven to suppress HEV in different populations: HEV infection after liver or kidney transplantation,<sup>13,15-18</sup> patients with hematologic neoplasms,<sup>36</sup> and those




Figure 3. Therapeutic approach for acute and chronic hepatitis E infection.

Table 2. Acute hepatitis E cases treated with ribavirin.

Author	Treated cases	Age	Gender	Genotype	Acute-on-chronic (aetiology)	Ribavirin dose (cumulative dose)	Days to viral clearance	Ribavirin side effects	Outcome (Follow-up)
Gerolami, et al. France 2011 <sup>85</sup>	1	61	Male	3	No	600 mg/12 h/21 d (25,200 mg)	15 days	None	NR
Péron, et al. France 2011 <sup>86</sup>	2	79	Male	3	Yes (alcoholic)	200 mg/48 h/90 d (9,000 mg)	30 days	None	VR (NR)
		NR	Male	3	Yes (alcoholic)	500 mg/12 h/10 d (10,000 mg)	30 days	Anaemia	VR (NR)
DelBello, et al. France 2012 <sup>78</sup>	1	65	Male	3f	Yes (OLT for alcoholic disease)	400 mg/24 h/90 d (36,000 mg)	15 days	NR	VR (2 months)
Goyal, et al. India 2012 <sup>39</sup>	4	50	Male	1	Yes (HBV + HCC)	300 mg/12 h/49 d (29,400 mg)	28 days	None	VR (6 months)
		16	Male	1	Yes (autoimmune)	200 mg/12 h/21 d (8,400 mg)	21 days	Anaemia	VR (6 months)
		18	Female	1	Yes (autoimmune)	200 mg/24 h/168 d (33,600 mg)	56 days	None	VR (6 months)
		50	Male	1	Yes (NR)	600 mg/24 h/84d (50,400 mg)	28 days	None	NR
Pischke, et al. Germany 2013 <sup>22</sup>	1	42	Female	1e	No	Dose not reported for 6 weeks	NR	NR	VR (NR)

D: days. HEV: hepatitis E virus. NR: not reported. RNA: ribonucleic acid. VR: virologic response.

with lymphocytopenia, both idiopathic and secondary to HIV infection.<sup>13,26</sup> The majority of cases treated with ribavirin achieved viral suppression during treatment (Table 1), with very few relapses after treatment discontinuation occurring in solid organ transplantation.<sup>15,20,22,23</sup> Nonetheless, the available follow-up is relatively short to know the true relapse rate. A recent study showed no difference in the rate of relapses between patients treated with ribavirin or peg-interferon for chronic hepatitis E infection, although only five patients received peg-interferon.<sup>23</sup> Furthermore, the appropriate dose and duration of ribavirin treatment is unknown. Based on the currently available data, a proposal for diagnostic and therapeutic approaches in acute and chronic hepatitis E is suggested in figure 3. The timing of therapy initiation also remains to be defined.

Concerning acute infection in the setting of chronic liver disease, many studies have shown a poor prognosis for acute hepatitis E.<sup>84-86</sup> A study in France<sup>87</sup> reported severe cases of fulminant acute infection, with hepatic encephalopathy and a high mortality rate. Active alcohol abuse and chronic liver disease were more frequent in patients suffering from the severe form. In another series of 28 cases of locally acquired hepatitis E in the United Kingdom,<sup>88</sup> there were two deaths due to liver failure. These two patients were older men with history of alcohol abuse and previously undiagnosed cirrhosis, highlighting the importance of acute-on-chronic liver failure due to HEV. The experience of ribavirin use for acute hepatitis E therapy is shown in table 2.

## HEPATITIS E PREVENTION

Improvements in the sanitary infrastructure and availability of clean drinking water may be the key prevention strategy to avoid hepatitis E transmission in developing areas. In industrialized countries, it seems evident that most cases of native hepatitis E may be related to zoonotic transmission of infection,<sup>53</sup> which can be responsible for hepatic fibrosis and even cirrhosis, mainly in immunocompromised patients. For this reason, it is vital to reinforce the recommendations for systematic cooking of pork (heating at 71° for at least 20 min) and avoiding raw meat if the cooking advice cannot be guaranteed.<sup>89</sup> Another basic issue with regard to immunocompromised patients is blood donation. Several cases of HEV transmission through blood products have been reported worldwide.<sup>48-51</sup> Although approximately 22% of blood

donors from a sample in the United States presented positive serology for anti-HEV antibody, HEV-RNA was not detected and there were no reports of HEV infection.<sup>90</sup> Hence, blood transmission exists, but its prevalence seems to be very low.

It is known that recovery from HEV infection results in protective immunity by neutralizing antibodies that can be detected in serum of individuals exposed to HEV, whatever the human genotype.<sup>91</sup> Protective immunity can also be induced by HEV vaccination. In 2007, a recombinant HEV vaccine developed by GlaxoSmith-Kline was tested in 2000 healthy Nepalese men susceptible to HEV infection included in a Phase II study. After 3 doses, 95.5% of subjects developed anti-HEV antibodies,<sup>92</sup> albeit this vaccine has not been further developed. A phase III study performed in China involving 112,604 participants in Jiangsu Province, where genotypes 3 and 4 are predominant, revealed an efficacy > 99% in preventing clinical hepatitis E among persons who completed the full 3-dose series of HEV 239 compared with placebo.<sup>93</sup> However, it should be noted that this vaccine is based on HEV genotype 1, and its efficacy against genotype 3 is unknown. In addition, the vaccine requires further information in special risk groups, such as patients with chronic liver disease and immunocompromised individuals. Regarding pregnant women, though information on the safety and efficacy of HEV 239 in this group is limited since pregnant women were excluded from the trial, a post hoc analysis of 68 participants whose pregnancies were detected after receiving 1 to 3 doses of vaccine or placebo rules out an increased risk of spontaneous abortion or congenital anomalies.<sup>94</sup> This vaccine has been licensed by the State Food and Drug Administration of China since January 2012, so new data on efficacy may be available in the next few years.

## ABBREVIATIONS

- **HEV:** hepatitis E virus.
- **HIV:** human immunodeficiency virus.
- **ORF:** open reading frames.
- **RNA:** ribonucleic acid.
- **VR:** virologic response.

## CONFLICT OF INTEREST

There is no commercial affiliation or author consultancy that could be construed as a conflict of interest with respect to the submitted data.

## REFERENCES

1. WHO. Viral Hepatitis Statement A62/22. Available from: [http://apps.who.int/gb/ebwha/pdf\\_files/A62/A62\\_22-en.pdf](http://apps.who.int/gb/ebwha/pdf_files/A62/A62_22-en.pdf)
2. Purcell RH, Emerson SU. Hepatitis E: an emerging awareness of an old disease. *J Hepatol* 2008; 48: 494-503.
3. Dalton HR, Bendall R, Ijaz S, Banks M. Hepatitis E: an emerging infection in developed countries. *Lancet Infect Dis* 2008; 8: 698-709.
4. Vishwanathan R. Infectious hepatitis in Delhi (1955-56), a critical study: epidemiology. *Indian J Med Res* 1957; 45: 49-58.
5. Balayan MS, Andjaparidze AG, Savinskaya SS, Ketiladze ES, Braginsky DM, Savinov AP, Poleschuk VF. Evidence for a virus in non-A, non-B hepatitis transmitted via the fecal-oral route. *Intervirology* 1983; 20: 23-31.
6. Bihl F, Negro F. Hepatitis E virus: a zoonosis adapting to humans. *J Antimicrob Chemother* 2010; 65: 817-21.
7. Kamar N, Selves J, Mansuy JM, Ouezzani L, Péron JM, Guittard J, Cointault O, et al. Hepatitis E virus and chronic hepatitis in organ-transplant recipients. *N Engl J Med* 2008; 358: 811-7.
8. Gérolami R, Moal V, Colson P. Chronic hepatitis E with cirrhosis in a kidney-transplant recipient. *N Engl J Med* 2008; 358: 859-60.
9. Gérolami R, Moal V, Picard C, Colson P. Hepatitis E virus as an emerging cause of chronic liver disease in organ transplant recipients. *J Hepatol* 2009; 50: 622-4.
10. Kamar N, Garrouste C, Haagsma EB, Garrigue V, Pischke S, Chauvet C, Dumortier J, et al. Factors associated with chronic hepatitis in patients with hepatitis E virus infection who have received solid organ transplants. *Gastroenterology* 2011; 140: 1481-9.
11. Halac U, Béland K, Lapierre P, Patey N, Ward P, Brassard J, Houde A, et al. Chronic hepatitis E infection in children with liver transplantation. *Gut* 2012; 61: 597-603.
12. Haagsma EB, van den Berg AP, Porte RJ, Benne CA, Venema H, Reimerink JH, Koopmans MP. Chronic hepatitis E virus infection in liver transplant recipients. *Liver Transpl* 2008; 14: 547-53.
13. Mallet V, Nicand E, Sultanik P, Chakvetadze C, Tessé S, Thervet E, Mouthon L, et al. Brief communication: case reports of ribavirin treatment for chronic hepatitis E. *Ann Intern Med* 2010; 153: 85-9.
14. Pischke S, Stiefel P, Franz B, Bremer B, Suneetha PV, Heim A, Ganzenmueller T, et al. Chronic hepatitis e in heart transplant recipients. *Am J Transplant* 2012; 12: 3128-33.
15. Kamar N, Rostaing L, Abravanel F, Garrouste C, Lhomme S, Esposito L, Basse G, et al. Ribavirin therapy inhibits viral replication on patients with chronic hepatitis e virus infection. *Gastroenterology* 2010; 139: 1612-8.
16. Chaillon A, Sirinelli A, De Muret A, Nicand E, d'Alteroche L, Goudeau A. Sustained virologic response with ribavirin in chronic hepatitis E virus infection in heart transplantation. *J Heart Lung Transplant* 2011; 30: 841-3.
17. Wedemeyer H, Pischke S, Manns MP. Pathogenesis and treatment of hepatitis e virus infection. *Gastroenterology* 2012; 142: 1388-97.
18. Kamar N, Weclawiak H, Guilbeau-Frugier C, Legrand-Abravanel F, Cointault O, Ribes D, Esposito L, et al. Hepatitis E virus and the kidney in solid-organ transplant patients. *Transplantation* 2012; 93: 617-23.
19. de Niet A, Zaaier HL, ten Berge I, Weegink CJ, Reesink HW, Beuers U. Chronic hepatitis E after solid organ transplantation. *Neth J Med* 2012; 70: 261-6.

20. Koning L, Pas SD, de Man RA, Balk AH, de Knecht RJ, ten Kate FJ, Osterhaus AD, et al. Clinical implications of chronic hepatitis E virus infection in heart transplant recipients. *J Heart Lung Transplant* 2013; 32: 78-85.
21. Riezebos-Brilman A, Puchhammer-Stöckl E, van der Weide HY, Haagsma EB, Jaksch P, Bejvl I, Niesters HG, et al. Chronic hepatitis E infection in lung transplant recipients. *J Heart Lung Transplant* 2013; 32: 341-6.
22. Pischke S, Hardtke S, Bode U, Birkner S, Chatzikyrkou C, Kauffmann W, Bara CL, et al. Ribavirin treatment of acute and chronic hepatitis E: a single-centre experience. *Liver Int* 2013; 33: 722-6.
23. Alric L, Bonnet D, Palacin A, Thébaud S, Peron JM, Sigur N, Nicot F, et al. Comparison of virological response between ribavirin and Peg-interferon monotherapy during autochthonous chronic hepatitis E virus infection. *J Hepatol* 2013; 58: S499.
24. Colson P, Kaba M, Moreau J, Brouqui P. Hepatitis E in an HIV-infected patient. *J Clin Virol* 2009; 45: 269-71.
25. Dalton HR, Bendall RP, Keane FE, Tedder RS, Ijaz S. Persistent carriage of hepatitis E virus in patients with HIV infection. *N Engl J Med* 2009; 361: 1025-7.
26. Dalton HR, Keane FE, Bendall R, Mathew J, Ijaz S. Treatment of chronic hepatitis E in a patient with HIV infection. *Ann Intern Med* 2011; 155: 479-80.
27. Kenfak-Foguena A, Schöni-Affolter F, Bürgisser P, Witteck A, Darling KE, Kovari H, Kaiser L, et al, and the Data Center of the Swiss HIV Cohort Study, Lausanne, Switzerland. Hepatitis E Virus seroprevalence and chronic infections in patients with HIV, Switzerland. *Emerg Infect Dis* 2011; 17: 1074-8.
28. Crum-Cianflone NF, Curry J, Drobeniuc J, Weintrob A, Landrum M, Ganesan A, Bradley W, et al; Infectious Disease Clinical Research Program HIV Working Group. Hepatitis E virus infection in HIV-infected persons. *Emerg Infect Dis* 2012; 18: 502-6.
29. Curry JA, Adams N, Crum-Cianflone NF. Acute hepatitis E infection in an HIV-infected person in the United States. *Ann Intern Med* 2009; 150: 226-7.
30. Keane F, Gompels M, Bendall R, Drayton R, Jennings L, Black J, Baragwanath G, et al. Hepatitis E virus coinfection in patients with HIV infection. *HIV Med* 2012; 13: 83-8.
31. Renou C, Lefeuvre A, Cadranet JF, Pavio N, Pariente A, Allègre T, Poggi C, et al. Hepatitis E virus in HIV-infected patients. *AIDS* 2010; 24: 1493-9.
32. Madejón A, Vispo E, Bottecchia M, Sánchez-Carrillo M, García-Samaniego J, Soriano V. Lack of hepatitis E virus infection in HIV patients with advanced immunodeficiency or idiopathic liver enzyme elevations. *J Viral Hepat* 2009; 16: 895-6.
33. Jardi R, Crespo M, Homs M, Van den Eynde E, Girones R, Rodríguez-J, Caballero A, et al. HIV, HEV and cirrhosis: evidence of a possible link from eastern Spain. *HIV Med* 2012; 13: 379-83.
34. Péron JM, Mansuy JM, Récher C, Bureau C, Poirson H, Alric L, Izopet J, et al. Prolonged hepatitis E in an immunocompromised patient. *J Gastroenterol Hepatol* 2006; 21: 1223-4.
35. Tamura A, Shimizu YK, Tanaka T, Kuroda K, Arakawa Y, Takahashi K, Mishiro S, et al. Persistent infection of hepatitis E virus transmitted by blood transfusion in a patient with T-cell lymphoma. *Hepatol Res* 2007; 37: 113-20.
36. Alric L, Bonnet D, Beynes-Rauzy O, Izopet J, Kamar N. Definitive clearance of a chronic hepatitis E virus infection with ribavirin treatment. *Am J Gastroenterol* 2011; 106: 1562-3.
37. Ollier L, Tieulie N, Sanderson F, Heudier P, Giordanengo V, Fuzibet JG, Nicand E. Chronic hepatitis after hepatitis E virus infection in a patient with non-Hodgkin lymphoma taking rituximab. *Ann Intern Med* 2009; 150: 430-1.
38. De Silva S, Hassan-Ibrahim MO, Austin M, Newport M, Verma S. Hepatitis E infection is an under recognized cause of acute decompensation in patients with chronic liver disease. *Dig Liver Dis* 2012; 44: 930-4.
39. Goyal R, Kumar A, Panda SK, Paul SB, Acharya SK. Ribavirin therapy for hepatitis E virus-induced acute on chronic liver failure: a preliminary report. *Antivir Ther* 2012; 17: 1091-6.
40. Borkakoti J, Hazam RK, Kar P. Hepatitis E virus: A hidden danger in acute on chronic liver failure. *J Hepatol* 2013; 58: S499-S500.
41. Reyes GR, Purdy MA, Kim JP, Luk KC, Young LM, Fry KE, Bradley DW. Isolation of a cDNA from the virus responsible for enterically transmitted non-A, non-B hepatitis. *Science* 1990; 247: 1335-9.
42. Tam AW, Smith MM, Guerra ME, Huang CC, Bradley DW, Fry KE, Reyes GR. Hepatitis E virus (HEV): molecular cloning and sequencing of the full-length viral genome. *Virology* 1991; 185: 120-31.
43. Xing L, Li TC, Mayakaki N, Simon MN, Wall JS, Moore M, Wang CY, et al. Structure of hepatitis E virion-sized particle reveals an RNA-dependent viral assembly pathway. *J Biol Chem* 2010; 285: 33175-83.
44. Lu L, Li C, Hagedorn CH. Phylogenetic analysis of global hepatitis E virus sequences: genetic diversity, subtypes and zoonosis. *Rev Med Virol* 2006; 16: 5-36.
45. Meng XJ, Purcell RH, Halbur PG, Lehman JR, Webb DM, Tsareva TS, Haynes JS, et al. A novel virus in swine is closely related to the human hepatitis E virus. *Proc Natl Acad Sci USA* 1997; 94: 9860-5.
46. Colson P, Romanet P, Moal V, Borentain P, Purgus R, Benezech A, Motte A, et al. Autochthonous infections with hepatitis E virus genotype 4, France. *Emerg Infect Dis* 2012; 18: 1361-4.
47. Jebblaoui A, Haim-Boukoubza S, Pause S, Mokhtari C, Nicand E, Roque-Afonso AM. Emerging hepatitis E genotype 4 infection in France. *J Hepatol* 2013; 58: S405.
48. Colson P, Coze C, Gallian P, Henry M, De Micco P, Tamalet C. Transfusion-associated hepatitis E, France. *Emerg Infect Dis* 2007; 13: 648-9.
49. Boxall E, Herborn A, Kochethu G, Pratt G, Adams D, Ijaz S, Teo CG. Transfusion-transmitted hepatitis E in a 'nonhyperendemic' country. *Transfus Med* 2006; 16: 79-83.
50. Matsubayashi K, Kang JH, Sakata H, Takahashi K, Shindo M, Kato M, Sato S, et al. A case of transfusion-transmitted hepatitis E caused by blood from a donor infected with hepatitis E virus via zoonotic food-borne route. *Transfusion* 2008; 48: 1368-75.
51. Haim-Boukoubza S, Ferey MP, Vétillard AL, Jebblaoui A, Pélissier E, Pelletier G, Teillet L, et al. Transfusion-transmitted hepatitis E in a misleading context of autoimmunity and drug-induced toxicity. *J Hepatol* 2012; 57: 1374-8.
52. Schlosser B, Stein A, Neuhaus R, Pahl S, Ramez B, Krüger DH, Berg T, et al. Liver transplant from a donor with occult HEV infection induced chronic hepatitis and cirrhosis in the recipient. *J Hepatol* 2012; 56: 500-2.
53. Colson P, Borentain P, Queyriaux B, Kaba M, Moal V, Gallian P, Heyries L, et al. Pig liver sausage as a source of hepatitis E virus transmission to humans. *J Infect Dis* 2010; 202: 825-34.
54. Tei S, Kitajima N, Takahashi K, Mishiro S. Zoonotic transmission of hepatitis E virus from deer to human beings. *Lancet* 2003; 362: 371-3.



55. Izopet J, Dubois M, Bertagnoli S, Lhomme S, Marchandeau S, Boucher S, Kamar N, et al. Hepatitis e virus strains in rabbits and evidence of a closely related strain in humans, France. *Emerg Infect Dis* 2012; 18: 1274-81.
56. Johne R, Heckel G, Plenge-Bönig A, Kindler E, Maresch C, Reetz J, Schielke A, et al. Novel hepatitis E virus genotype in Norway rats, Germany. *Emerg Infect Dis* 2010; 16: 1452-5.
57. Johne R, Dremsek P, Kindler E, Schielke A, Plenge-Bönig A, Gregersen H, Wessels U, et al. Rat hepatitis E virus: geographical clustering within Germany and serological detection in wild Norway rats (*Rattus norvegicus*). *Infect Genet Evol* 2012; 12: 947-56.
58. Lack JB, Volk K, Van Den Bussche RA. Hepatitis e virus genotype 3 in wild rats, United States. *Emerg Infect Dis* 2012; 18: 1268-73.
59. Geng Y, Zhao C, Song A, Wang J, Zhang X, Harrison TJ, Zhou Y, et al. The serological prevalence and genetic diversity of hepatitis E virus in farmed rabbits in China. *Infect Genet Evol* 2011; 11: 476-82.
60. Zhao C, Ma Z, Harrison TJ, Feng R, Zhang C, Qiao Z, Fan J, et al. A novel genotype of hepatitis E virus prevalent among farmed rabbits in China. *J Med Virol* 2009; 81: 1371-9.
61. Takahashi M, Nishizawa T, Sato H, Sato Y, Jirintai, Nagashima S, Okamoto H. Analysis of the full-length genome of a hepatitis E virus isolate obtained from a wild boar in Japan that is classifiable into a novel genotype. *J Gen Virol* 2011; 92: 902-8.
62. Sato Y, Sato H, Naka K, Furuya S, Tsukiji H, Kitagawa K, Sonoda Y, et al. A nationwide survey of hepatitis E virus (HEV) infection in wild boars in Japan: identification of boar HEV strains of genotypes 3 and 4 and unrecognized genotypes. *Arch Virol* 2011; 156: 1345-58.
63. Galiana C, Fernández-Barredo S, García A, Gómez MT, Pérez-Gracia MT. Occupational exposure to hepatitis E virus (HEV) in swine workers. *Am J Trop Med Hyg* 2008; 78: 1012-5.
64. Rodriguez-Manzano J, Miagostovich M, Hundesa A, Clemente-Casares P, Carratala A, Buti M, Jardi R, et al. Analysis of the evolution in the circulation of HAV and HEV in eastern Spain by testing urban sewage samples. *J Water Health* 2010; 8: 346-5.
65. Berto A, Martelli F, Grierson S, Banks M. Hepatitis e virus in pork food chain, United kingdom, 2009-2010. *Emerg Infect Dis* 2012; 18: 1358-60.
66. Di Bartolo I, Diez-Valcarce M, Vasickova P, Kralik P, Hernandez M, Angeloni G, Ostanello F, et al. Hepatitis E virus in pork production chain in Czech Republic, Italy, and Spain, 2010. *Emerg Infect Dis* 2012; 18: 1282-9.
67. Bhatia V, Singhal A, Panda SK, Acharya SK. A 20-year single-center experience with acute liver failure during pregnancy: is the prognosis really worse? *Hepatology* 2008; 48: 1577-85.
68. Khuroo MS, Kamili, Jameel S. Vertical transmission of hepatitis E virus. *Lancet* 1995; 345: 1025-6.
69. Aggarwal R, Naik S. Epidemiology of hepatitis E: Current status. *J Gastroenterol Hepatol* 2009; 24: 1484-93.
70. Kamar N, Mansuy JM, Cointault O, Selves J, Abravanel F, Danjoux M, Otal P, et al. Hepatitis E virus-related cirrhosis in kidney- and kidney-pancreas-transplant recipients. *Am J Transplant* 2008; 8: 1744-8.
71. Hamid S, Atiq M, Shehzad F, Yasmeen A, Nissa T, Salam A, Siddiqui A, et al. Hepatitis E virus superinfection in patients with chronic liver disease. *Hepatology* 2002; 36: 474-8.
72. Gerolami R, Moal V, Picard C, Colson P. Hepatitis E virus as an emerging cause of chronic liver disease in organ transplant recipients. *J Hepatol* 2009; 50: 622-4.
73. Abravanel F, Mansuy JM, Huynh A, Kamar N, Alric L, Peron JM, Récher C, et al. Low risk of hepatitis E virus reactivation after haematopoietic stem cell transplantation. *J Clin Virol* 2012; 54: 152-5.
74. Koenecke C, Pischke S, Heim A, Raggub L, Bremer B, Raupach R, Buchholz S, et al. Chronic hepatitis E in hematopoietic stem cell transplant patients in a low-endemic country? *Transpl Infect Dis* 2012; 14: 103-6.
75. Legrand-Abravanel F, Kamar N, Sandres-Saune K, Garrouste C, Dubois M, Mansuy JM, Muscari F, et al. Characteristics of autochthonous hepatitis E virus infection in solid-organ transplant recipients in France. *J Infect Dis* 2010; 202: 835-44.
76. Sood A, Midha V, Sood N. Guillain-Barré syndrome with acute hepatitis E. *Am J Gastroenterol* 2000; 95: 3667-8.
77. Kamar N, Bendall RP, Peron JM, Cintas P, Prudhomme L, Mansuy JM, Rostaing L, et al. Hepatitis E Virus and neurologic disorders. *Emerg Infect Dis* 2011; 17: 173-9.
78. Del Bello A, Arné-Bes MC, Lavayssière L, Kamar N. Hepatitis E virus-induced severe myositis. *J Hepatol* 2012; 57: 1152-3.
79. Fourquet E, Mansuy JM, Bureau C, Recher C, Vinel JP, Izopet J, Péron JM. Severe thrombocytopenia associated with acute autochthonous hepatitis E. *J Clin Virol* 2010; 48: 73-4.
80. Li C, Wang HF. Hepatitis E virus-related acute liver failure associated with pure red cell aplasia. *Hepatobiliary Pancreat Dis Int* 2011; 10: 557-8.
81. Shah SA, Lal A, Idrees M, Hussain A, Jeet C, Malik FA, Iqbal Z, et al. Hepatitis E virus-associated aplastic anaemia: the first case of its kind. *J Clin Virol* 2012; 54: 96-7.
82. Khudyakov Y, Kamili S. Serological diagnostics of hepatitis E virus infection. *Virus Res* 2011; 161: 84-92.
83. Aggarwal R. Clinical presentation of hepatitis E. *Virus Res* 2011; 161: 15-22.
84. Kamar N, Garrouste C, Haagsma EB, Garrigue V, Pischke S, Chauvert C, Dumortier J, et al. Factors associated with chronic hepatitis in patients with hepatitis E virus infection who have received solid organ transplants. *Gastroenterology* 2011; 140: 1481-9.
85. Gerolami R, Borentain P, Raissouni F, Motte A, Solas C, Colson P. Treatment of severe acute hepatitis E by ribavirin. *J Clin Virol* 2011; 52: 60-2.
86. Péron JM. Acute autochthonous hepatitis E in western patients with underlying chronic liver disease: A role for ribavirin? *J Hepatol* 2011; 54: 1323-4.
87. Péron JM, Bureau C, Poirson H, Mansuy JM, Alric L, Selves J, Dupuis E, et al. Fulminant liver failure from acute autochthonous hepatitis E in France: description of seven patients with acute hepatitis E and encephalopathy. *J Viral Hepatol* 2007; 14: 298-303.
88. Dalton HR, Hazeldine S, Banks M, Ljaz S, Bendall R. Locally acquired hepatitis E in chronic liver disease. *Lancet* 2007; 369: 1260.
89. Barnaud E, Rogée S, Garry P, Rose N, Pavio N. Thermal inactivation of infectious hepatitis E virus in experimentally contaminated food. *Appl Environ Microbiol* 2012; 78: 5153-9.
90. Xu C, Alter HJ. An assessment of HEV prevalence and risk in U.S. blood donors and recipients. Available from: (<http://www3.niddk.nih.gov/fund/other/HepE2012/HepatitisE2012ProgramBook.pdf>)

91. Zhou YH, Purcell RH, Emerson SU. A truncated ORF2 protein contains the most immunogenic site on ORF2: antibody responses to non-vaccine sequences following challenge of vaccinated and non-vaccinated macaques with hepatitis E virus. *Vaccine* 2005; 23: 3157-65.
92. Shrestha MP, Scott RM, Joshi DM, Mammen MP Jr, Thapa GB, Thapa N, Myint KS, et al. Safety and efficacy of a recombinant hepatitis E vaccine. *N Engl J Med* 2007; 356: 895-903.
93. Zhu FC, Zhang J, Zhang XF, Zhou C, Wang ZZ, Huang SJ, Wang H, et al. Efficacy and safety of a recombinant hepatitis E vaccine in healthy adults: a large-scale, randomised, double-blind placebo-controlled, phase 3 trial. *Lancet* 2010; 376: 895-902.
94. Wu T, Zhu FC, Huang SJ, Zhang XF, Wang ZZ, Zhang J, Xia NS. Safety of the hepatitis E vaccine for pregnant women: a preliminary analysis. *Hepatology* 2012; 55: 2038.