Annals of **Hepatology**

CONCISE REVIEW

July-August, Vol. 12 No.4, 2013: 359-363

Occult hepatitis B in patients on hemodialysis: a review

Andrea M.M. Fontenele,* Natalino Salgado Filho,*,** Adalgisa S.P. Ferreira**

*Nephrology Service, University Hospital of Federal University of Maranhão, Brazil. **Center of Clinical Research, University Hospital of Federal University of Maranhão, Brazil.

ABSTRACT

The hepatitis B virus (HBV) is one of the most frequently transmitted agents in dialysis units. Occult hepatitis B is characterized by HBV infection without detectable surface antigen (HBsAg) in the patient's serum, a positive or negative HBV DNA marker result in the serum and a positive result in the liver tissue, which leads to the potential risk of transmission during renal replacement therapy service. There is variation in occult hepatitis B prevalence rates in this population across various studies that may be related to numerous factors. The presence of occult hepatitis B in individuals undergoing renal replacement therapy is important with regard to both the possibility of transmission and the consequences for the patient, especially the development of chronic liver disease and reactivation of the disease after renal transplantation.

Key words. HBV. Renal disease. HBV-DNA. Prevalence. Occult HBV.

INTRODUCTION

The hepatitis B virus (HBV) is one of the most common infections in the world (it is estimated that there are over 360 million carriers of the chronic infection) and is a major cause of hepatic cirrhosis and hepatocellular carcinoma (HCC).¹ HBV transmission by blood products originating from individuals with negative results for the virus surface antigen (HBsAg) has been demonstrated since the 1980s. These individuals are known as occult hepatitis B carriers (OHB).² OHB has been linked to cases of cryptogenic cirrhosis and HCC.^{3,4} As patients with chronic renal failure (CRF) on dialysis are a population at risk for the parenteral transmission of pathogens, HBV infection is a continuing concern despite all of the measures used in its prevention, including mandatory vaccination for all susceptible individuals⁵⁻⁷ OHB can be a significant problem in this population, justifying this review of the topic.

OCCULT HEPATITIS B

The definition of OHB was standardized in 2008 at a meeting of experts in the city of Taormina

Correspondence and reprint request: Andrea MM Fontenele, Pharm, MSc. Tel.: 55 98 2109-1047, 55 98 8142-7743 E-mail: andrea.fontenele@yahoo.com.br

> Manuscript received: April 02, 2013. Manuscript accepted: April 15, 2013.

(Italy).⁸ The main definitions resulting from this meeting were as follows:

- OHB is a clinical entity in which the individual is positive for HBV DNA in liver tissue but has undetectable serum HBsAg by routinely used methods and positive or negative HBV-DNA in the serum, and
- When the HBV-DNA in the serum is positive, it often does not exceed 200 IU/mL.

This last point was added by the meeting participants to distinguish so-called false occult hepatitis B, in which HBsAg is not detected due to infection with the S gene escape mutant of the HBV genome, which prevents the production of HBsAg but has a high viral load that is consistent with loads that are manifested during classic chronic infection phases. There are still differences in the literature regarding the definition of this entity, as many authors consider the latter to be OHB.⁹

It is believed that most cases of OHB follow an acute HBV infection in which viral replication continues indefinitely after the disappearance of HBsAg or occurs after many years of chronic infection in which there was loss of HBsAg.¹⁰⁻¹⁴ These individuals can transmit HBV through transfusions of their blood derivatives or through donation of their organs for transplant and may, if subjected to immunosuppression, show reactivation of hepatitis $B.^{8,14,15}$

© 2019, Fundación Clínica Médica Sur, A.C. Published by Elsevier España S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Most OHB carriers show positivity for other serological markers of HBV (anti-HBc and/or anti-HBs), but 20% of them are seronegative for all markers.¹⁶

Unique characteristics of HBV infection and one that seems to be fundamental in OHB is the persistence of the circular shape of its episomal genomic DNA (cccDNA) in the hepatocyte nucleus, which maintains viral replication.^{17,18} It has been observed that the HBV variant most frequently found in OHB is the wild type strain, with no genetic mutations and with replication stages similar to those occurring in the classical infection,^{19,20} but strong suppression of viral replication and gene expression by antiviral cytotoxic T cells is responsible for the very low or undetectable levels of serum HBV DNA.²⁰ Another event that can suppress replication of HBV is co-infection with other viruses, particularly the hepatitis C virus (HCV).²¹ It has been reported that the antigen core of HCV strongly inhibits the replication of HBV.¹⁷

The exact clinical relevance of OHB is not clear, but the possible role exerted by this infection on the cryptogenic cirrhosis and HCC development has been demonstrated by several studies and recently revised.¹⁷ It has also been noted that the presence of OHB can negatively affect the response to treatment of chronic hepatitis C.²² Another important feature of this entity is the possibility of reactivating hepatitis B under immunosuppressive conditions, such as chemotherapy and organ transplantation.^{15,23,24}

DIAGNOSIS

Standardized tests for the diagnosis of OHB are still not available. The gold standard test currently used is the identification of HBV DNA in liver tissue by very sensitive amplification tests (nested PCR and real-time PCR).^{8,9,16} As tests on liver tissue are not easily available because of the necessity of performing a liver biopsy, in practice, these tests are performed on serum. The Taormina consensus recommended using primers for different highly conserved segments of the viral genome. Usually the S, pre-C/C and X regions are used.²⁵

PREVALENCE

The prevalence of OHB predominantly depends on a region's level of endemicity¹⁹ and the susceptibility of the population studied, which is higher in those populations with higher risk of parenteral infection and immunosuppressed patients,^{16,26,27} HCV carriers²⁸ and those with cirrhosis and HCC of unknown causes.^{3,4}

In addition to these traditional populations, OHB has also been reported in low-risk populations such as blood donors in various parts of the world, including Brazil.^{29,30}

HEPATITIS B IN HEMODIALYSIS

Chronic kidney disease in its most advanced stage (stage V) is a significant problem that is present in almost all countries. The incidence has continued to increase, mainly as a result of the consequences of diabetes mellitus and hypertension.³¹ Renal replacement therapies (RRT) such as hemodialysis are the final treatment for these individuals.

HBV infection can cause increased morbidity in patients in the terminal stage of chronic kidney disease. As one of the main modes of transmission of this virus is the parenteral route, these patients have a high risk of acquiring an infection by this mechanism because hemodialysis requires prolonged vascular access. It has been observed that HBV is the most common virus transmitted by this route in this population.^{7,32,33}

Despite this susceptibility, the prevalence of HBV has decreased in dialysis centers around the world, including those in Brazil.³³⁻³⁴ According to data from the Brazilian Society of Nephrology the prevalence of HBsAg in Brazilian dialysis units was approximately 1.1% in 2010.³⁴ These numbers were as high as 41% for the same serological marker in the 1980s.³⁵ This decrease can be attributed to mandatory vaccination, segregation of infected individuals into different rooms and dedicated hemodialysis machines, and the implementation of standard precautions to prevent pathogen transmission in dialysis centers.^{33,36}

Once the individual on dialysis is contaminated with HBV, the possibility of becoming a chronic carrier is very high, which is not the case with immunocompetent individuals. Although this population is immunocompromised, the natural history of this infection in this population appears no different from that observed in normal individuals.³⁷ Its major clinical importance seems to be after renal transplantation, as it can cause reactivation episodes that may be so severe that they cause the patient's death.³⁸

OCCULT HEPATITIS B IN PATIENTS ON HEMODIALYSIS

HBV infection with HBsAg-negative serum in hemodialysis patients was initially reported by Cabrerizo, *et al.*³⁹ This team investigated 33 pa-

Table 1. Prevalence rates of occult hepatitis B in patients undergoing renal replacement therapy (hemodialysis).

Authors	Year	Country	Population (n)	Prevalence
Cabrerizo ³⁹	1997	Spain	33	19 (58%)
Minuk ⁴¹	2004	Canada	241	9 (3.8%)
Fabrizzi ⁴²	2005	Italy	585	0 (0%)
Teruel ⁴³	2005	Madrid	61	0 (0%)
Siagris ⁴⁴	2006	Greece	49	10 (20.4%)
Goral ⁴⁵	2006	Turkey	50	0 (0%)
Yakaryilmaz ⁴⁶	2006	Turkey	188	5 (2.7%)
Altindis ⁴⁷	2007	Turkey	153	19 (12.4%)
Jardim ⁴⁸	2008	Brazil	34	0 (0%)
Ersoy ⁴⁹	2008	Turkey	80	1 (1.25%)
Gwak ⁵⁰	2008	Korea	83	0 (0%)
Di Stefano ⁵¹	2009	Italy	128	34 (26.6%)
Motta ⁵²	2010	Brazil	100	15 (15%)
Mina ⁵³	2010	Greece	366	3 (0.9%)
Aghakhani ⁵⁴	2010	Iran	289	9 (3.11%)
Ismail ⁵⁵	2010	Egypt	116	2 (3.8%)
Sav ⁵⁶	2010	Turkey	71	12 (16.9%)
Makarem ⁵⁷	2012	Egypt	145	6 (4.1%)
Joukar ⁵⁸	2012	Iran	514	0 (0%)
Albuquerque ⁵⁹	2012	Brazil	752	3 (1.5%)

tients who were HBsAg negative looking for HBV-DNA in the serum and detected the marker in 19 patients (58%). Among these patients, 12 (63%) had a serological marker for HBV. Since that report, many studies have been published on the subject, revealing the presence of a significant problem in almost all countries, as recently reviewed by Hollinger, *et* $al.^{40}$

The prevalence of OHB in hemodialysis is quite variable, ranging from 0-58% (Table 1). This variability, as observed in subjects with normal renal function, must be related to the degree of endemicity of HBV in a region, the sensitivity of the tests used and selection biases.^{39,41-59}

Higher prevalence rates of OHB have been reported in the presence of anti-HBc as an isolated marker, with HBsAg and anti-HBs negative in patients without kidney failure. An Iranian study found the presence of HBV DNA in 50% of patients on hemodialysis with isolated anti-HBc.⁵⁴ In contrast, a study by Fabrizzi, et al.42 involving more than 500 patients on hemodialysis in Italy found no positivity for HBV-DNA among isolated anti-HBc carriers. Similarly, in a recently published Brazilian study, HBV-DNA was positive in only 3 patients out of 135 carriers of the isolated anti-HBc profile.⁵⁹ Thus, the isolated presence of anti-HBc in patients undergoing hemodialysis may have very different meanings depending on the region where it is being studied. Moreover, among hemodialysis patients infected with HCV, OHB has been found in varying degrees from 0-36%.^{44-46,51} In Brazil, where the prevalence of OHB patients on hemodialysis has ranged from 0 to 15%,^{48,52,59} a greater frequency has not been observed among patients with HCV.^{48,52}

The need to identify individuals with OHB in dialysis units is predicated on the fact that these patients may be potential sources of HBV transmission, as the nosocomial transmission of the virus is well documented in this area.^{32,33} Whether all individuals undergoing dialysis should be submitted to HBV-DNA screening or whether this test should be restricted to high-risk populations still has not been ascertained.⁶⁰

CONCLUSION

Although the vast majority of studies on the prevalence of occult hepatitis B in hemodialysis point toward a low prevalence, it seems advisable to search for OHB under certain conditions. Screening for HBV infection should be permanent in dialysis services, and it seems sensible to test for OHB whenever there are co-infections, especially HCV and HIV, in patients with isolated antiHBc, those with liver disease of an indeterminate cause and in renal transplant candidates. This diagnosis can minimize the transmission of the virus under these conditions, as preventive measures can be taken, carriers can be screened for HCC, and serious complications due to reactivation after renal transplantation can be prevented.

REFERENCES

- 1. Lai M, Yun-Fan L. Chronic Hepatitis B: past, present and future. *Clin Liver Dis* 2010; 14: 531-46.
- Brechot C, Degos F, Lugassy C, Thiers V, Zafrani S, Franco D, Bismuth H, et al. Hepatitis B virus DNA in patients with chronic liver disease and negative tests for hepatitis B surface antigen. N Engl J Med 1985; 312: 270-76.
- Chemin L, Zoulin F, Merle P, Arkhis A, Chevallier M, Kay A, Cova L, et al. High incidence of hepatitis B infections among chronic hepatitis cases of unknow aetiology. J Hepatol 2011; 34: 447-54.
- Fang Y, Shang QL, Liu JY, Li D, Xu WZ, Teng X, Zhao HW, et al. Prevalence of occult hepatitis B virus infection among hepatopathy patients and healthy people in China. J Infect 2009; 58: 383-8.
- Kara Y, Yilmaz Me, Suner A, Kadiroglu Ak, Isikoglu B. The evaluation of immune responses that occur after HBV infection and HBV vaccination in hemodialysis patients. *Vaccine* 2004; 22: 3963-67.
- 6. Zanetti Ar, Van Damme P, Shouval D. The global impact of vaccination against hepatitis B: A historical overview. *Vaccine* 2008; 26: 6266-73.
- 7. Edey M, Barraclough K, Johnson DW. Review article: Hepatitis B and dialysis. *Nephrology* 2010; 15: 137-45.
- Raimondo G, Allain J-P, Brunetto M, Buendia MA, Chen D-S, Colombo M, Craxi A, et al. Statements from the Taormina expert meeting on occult hepatitis B virus infection. J Hepatol 2008; 49: 652-7.
- 9. Hollinger FB, Sood G. Occult hepatitis B virus infection: a covert operation. *J Viral Hepat* 2010; 17: 1-15.
- Penna A, Artini M, Cavallii A, Leverero M, Bertoletti A, Pilli M, Chisari FV, et al. Long-lasting memory T-cell responses following self limited acute hepatitis B. J Clin Invest 1996; 98: 1185-94.
- 11. Rehermann B, Ferrari C, Pasquinelli C, Chisari FV. The hepatitis B virus persists for decades after patients' recovery from acute viral hepatitis despite active maintenance of a cytotoxic T-lymph. *Nat Med* 1996; 10: 1104-8.
- Guidotti LG, Rochoford R, Chung J, Shapiro M, Purcell R, Chisari FV. Viral clearance without destruction of infected cells during acute HBV infection. *Science* 1999; 284: 825-9.
- Guidotti LG, Chisari FV. Noncytolytic control of viral infections by the innate and adaptive immune response. *Annu Rev Immunol* 2001; 19: 65-91.
- Lok SFA. Occult hepatitis B virus infection: Diagnosis, implications and management? J Gastroenterol Hepatol 2004; 19: S114-S117.
- Marcellin P, Giostra E, Martinot-Peignoux M, Loriot MA, Jeagle ML, Wolf P, Degott C, et al. Redevelopment of hepatitis B surface antigen after renal transplantation. *Gastroenterology* 1991; 100: 1432-4.
- 16. Toberson M, Thomas DL. Occult hepatitis B. *Lancet Infect Dis* 2002; 2: 479-86.
- 17. Raimondo G, Caccamo G, Filomia R, Pollicino T. Occult HVB infection. *Semin Immunopathol* 2013; 35: 39-52.
- Hollinger FB. Hepatitis B virus infection and transfusion medicine: science and the occult. *Transfusion* 2008; 48: 1001-26.
- Lledó JL, Fernández C, Gutiérrez ML, Ocaña S. Management of occult hepatitis B virus infection: an update for clinician. World J Gastroenterol 2011; 17: 1563-8.
- Bes M, Vargas V, Piron M, Casamitgana N, Esteban JI, Vilanova N, Pinacho A, et al. T cell responses and viral variability in blood donation candidates with occult HBV infection. J Hepatol 2012; 56: 765-74.

- 21. Raimondo G, Pollicino T, Cacciolla I, Squadrito G. Occult hepatitis B virus infection. J Hepatol 2007; 46: 160-70.
- 22. Mrani S, Chemin I, Menouar K, Guillaud O, Pradat P, Borghi G, Trabaud MA, et al. Occult HBV infection may represent a major risk factor of non-response to antiviral therapy of chronic hepatitis C. J Med Virol 2007; 79: 1075-81.
- Kusumoto S, Tanaka Y, Mizokami M, Ueda R. Reactivation of hepatitis B virus following systemic chemotherapy for malignant lymphoma. *Int J Hematol* 2009; 90: 13-23.
- 24. Kalia H, Fabrizi F, Martin P. Hepatitis B virus and renal transplantation. *Transpl Rev* 2011; 25: 102-9.
- Ocana S, Casas ML, Buhigas I, Lledo JL. Diagnostic strategy for occult hepatitis B virus infection. World J Gastroenterol 2011; 17: 1553-7.
- Copolla N, Tonzielo G, Pisauturo M, Messina V, Guastafierro S, Fiori M, Iodice V, et al. Reactivation of overt and occult hepatitis B infection in various immunosuppressive settings. J Med Virol 2011; 83: 1909-16.
- Arababadi MK, Ahmadabadi Bn, Daredor Hy, Kennedy D. Epidemiology of occult hepatitis B infection among thalassemic, hemophilia and hemodialysis patients. *Hepat Mon* 2012; 12: 315-9.
- Carreño V, Bartolomé J, Castillo I, Quiroga A. Occult hepatitis B virus and hepatitis C virus infections. *Rev Med Virol* 2008; 18: 139-57.
- 29. Allain J-P, Mihaljevic I, Gonzalez-Fraile MI, Grubbe K, Holm-Harritshoj L, Garcia JM, Brojer E, et al. Infectivity of blood products from donors with occult hepatitis B virus infection. *Transfusion* 2013. Doi: 10.1111/trf.12096.
- 30. Silva CM, Costi C, Costa C, Michelon C, Oravec R, Ramos AB, Niel C, et al. Low rate of occult hepatitis B virus infection among anti-HBc positive blood donors living in a low prevalence region in Brazil. J Infect 2005; 51: 24-9.
- 31. Rizwan AH. The burden of chronic kidney disease. *BMJ* 2006; 332: 763-74
- Wong PN, Fung TT, Mak SK, Lo KY, Tong GM, Wong Y, Loo CK, et al. Hepatitis B virus infection in dialysis patients. J Gastroenterol Hepatol 2005; 20: 1641-51.
- 33. Fabrizi F, Messa P, Martin P. Hepatitis B virus infection and the dialysis patient. *Semi Dial* 2008; 21: 440-6.
- 34. Sesso RC, Lopes AA, Thomé FS, Lugon JR, Santos DR. Relatório do Censo Brasileiro de diálise de 2010 [2010 Report of the Brazilian Dialysis Census]. J Bras Nefrol 2011; 33: 442-7.
- 35. Cendorologo-Neto M, Draibe SA, Silva AEB, Ferraz ML, Granato C, Pereira CAP, Sesso RC, et al. Incidence of and risk factors for hepatitis B virus and hepatitis C virus Incidence of and risk factors for hepatitis B virus and hepatitis C virus infection among hemodialysis and CAPD patients: evidence for environmental transmission. *Nephrol Dial Transplant* 1995; 10: 240-6.
- 36. Fabrizi F, Marzano A, Messa P, Martin P, Lampertico P. Hepatitis B virus infection in the dialysis population: current perspectives. Int J Artif Organs 2008; 31: 386-94.
- Hanett JD, Parfrey PS, Kennedy M, Zeldis JB, Steinman TI, Guttman RD. The long-term outcome of hepatitis B infection in hemodialysis patients. *Am J Kidney Dis* 1988; 11: 210-3.
- Chen GD, Gu JL, Qiu J, Chen LZ. Outcomes and risk factors for hepatitis B virus (HBV) reactivation after kidney transplantation in occult HBV carriers. *Tranpl Infect Dis* 2013. Doi: 10.1111/tid.12065.
- 39. Cabrerizzo M, Bartolomé J, De Sequeira P, Caramelo C, Carreño V. Hepatitis B virus DNA in serum and blood cells of hepatitis B surface antigen-negative hemodialysis patients and staff. J Am Soc Neprhol 1997; 8: 1443-7.

- Hollinger FB, Habibollahi P, Daneshmand A, Alavian SM. Occult hepatitis B infection in chronic hemodialysis patients: Current concepts and strategy. *Hepat Mon* 2010; 10: 199-204.
- Minuk GY, Sun DF, Greenberg R, Zhanq M, Hawkins K, Uhanova J, Gutkin A, et al. Occult hepatitis B virus infection in a north American adult hemodialysis patient population. *Hepatology* 2004; 40: 1072-7.
- Fabrizzi F, Messa PG, Lunghi G, Aucella F, Bisegna S, Mangano S, Villa M, et al. Occult hepatitis B virus infection in dialysis patients: a multicentre survey. *Aliment Pharma*col Ther 2005; 21: 1341-7.
- Teruel JL, Mateos LA, Fernández L, Ortuño J. Occult hepatitis B virus infection in patients treated with chronic hemodialysis. *Nefrología* 2005; 23: 83-4.
- 44. Siagris D, Christofidou M, Triga K, Pagoni N, Theochris GJ, Goumenos D, Lekkou A, et al. Occult hepatitis B virus infection in hemodialysis patients with chronic HCV infection. J Nephrol 2006; 19: 327-33.
- 45. Goral V, Okzul H, Tekes S, Sit D, Kadiroglu AK. Prevalence of occult HBV infection in haemodialysis patients with chronic HCV. World J Gastroenterol 2006; 7 3420-4.
- 46. Yakaryilmaz F, Gurbuz OA, Guliter S, Mert A, Songur Y, Karakan T, Keles H. Prevalence of occult hepatitis B and hepatitis C virus infections in Turkish hemodialysis patients. *Ren Fail* 2006; 28: 729-35.
- Altindis M, Uslan I, Cetinkaya Z, Yusel S, Cifiçi IH, Demirturk N, Ozdemir M, et al. Investigation of hemodialysis patients in terms of the presence of occult hepatitis B. *Mikrobiyol Bul* 2007; 41: 227-33.
- Jardim RN, Gonçales NS, Pereira JS, Fais VC, Gonçales Junior FL. Occult hepatitis B virus infection in immunocompromised patients. *Braz J Infect Dis* 2008; 12: 300-5.
- 49. Ersoy O, Yilmaz R, Arici M, Turgan C, Bayraktar Y. Prevalence of occult hepatitis B infection in hemodialysis patients. *Dialysis Transplant* 2008; 37: 362-8.
- Gwak GY, Huh W, Lee DH, Min BH, Koh KC, Kim JJ, Oh HY. Occult hepatitis B virus infection in chronic hemodialysis patients in Korea. *Hepatogastroenterology* 2008; 55: 1721-4.
- 51. Di Stefano M, Volpe A, Stallone G, Tartaglia L, Prato R, Martinelli D, Pastore G, et al Occult HBV infection in

hemodialysis setting is marked by presence of isolated antibodies to HBcAg and HCV. *J Nephrol* 2009; 22: 381-6.

- 52. Motta JS, Mello FC, Lago BV, Perez RM, Gomes SA, Figueiredo FF. Occult hepatitis B virus infection and lamivudineresistant mutations in isolates from renal patients undergoing hemodialysis. J Gastroenterol Hepatol 2010; 25: 101-6.
- 53. Mina P, Georgiadou SP, Rizos C, Dalekos GN, Rigopoulou EL. Prevalence of occult hepatitis B virus infection in hemodialysis patients from Central Greece. *World J Gastroenterol* 2010; 16: 225-31.
- 54. Aghakhani A, Banifazl M, Kalantar E, Eslamifar A, Amadi F, Razeqhi E, Atabak S, et al. Occult hepatitis B virus infection in hemodialysis patients with isolated hepatitis B core antibody: a multicenter study. *Ther Apher Dial* 2010; 14: 349-53.
- 55. Ismail H, Soliman M, Ismail N. Occult hepatitis B virus infection in Egyptian hemodialysis patients with or without hepatitis C virus infection. *Pathol Lab Med Int* 2010; 2: 113-20.
- 56. Sav T, Gursoy S, Torun E, Sav NM, Unal A, Oymak O, Utas C. Occult HBV infection in continuous ambulatory peritoneal dialysis and hemodialysis patients. *Ren Fail* 2010; 32: 74-7.
- 57. Abu El Makarem MA, Abdel Hamid M, Abdel Aleem A, Ali A, Shatat M, Sayed D, Deaf A, et al. Prevalence of occult hepatitis B virus infection in hemodialysis patients from Egypt with or without hepatitis C virus infection. *Hepat Mon* 2012; 12: 253-8.
- Joukar F, Mansour-Ghnael F, Beshraati S, Khosh-Sorour M. Occult hepatitis B infection in a hemodialysis population in Guilan province, northern Iran. *Hemodial Int* 2012; 16: 294-7.
- 59. Albuquerque ACC, Coêlho MRCD, Lemos MF, Moreira RC. Occult hepatitis B virus infection in hemodialysis patiens in Recife, Satate of Pernambuco, Brazil. *Rev Soc Bras Med Trop* 2012; 45: 558-62.
- Fabrizzi F, Martin P. Management of hepatitis B and C virus infection before and after renal transplantation. *Curr Opin Organ Transplant* 2006; 11: 583-8.