


Images in hepatology

## Effect of cold storage/reperfusion on rat hepatic collagen network organization

Alejandra B Quintana;<sup>1</sup> Joaquín V Rodríguez;<sup>2</sup> Edgardo E Guibert<sup>3</sup>



<sup>1</sup> Morfología, Departamento de Ciencias Fisiológicas.

<sup>2</sup> Farmacología, Departamento de Ciencias Fisiológicas.

<sup>3</sup> Biología Molecular, Departamento de Ciencias Biológicas, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Argentina.

Address for correspondence:

Dra. Alejandra B. Quintana, Area Morfología, Dto. de Ciencias Fisiológicas, Fac. de Cs. Bioquímicas y Farmacéuticas, UNR.

Suipacha 531 (S2002LRK). Rosario, Argentina. PHONE – FAX: 0054-341-4804601.

E-mail: aquintan@fbioyf.unr.edu.ar

Manuscript received and accepted: 29 August and 27 September 2007

Rat livers were cold stored (0°C-48 Hs) in University of Wisconsin (UW) solution. To reduce cold preservation/reperfusion injuries, three concentrations (50, 100 and 250  $\mu$ M) of S-nitrosoglutathione (GSNO) were studied. GSNO, which releases Nitric Oxide (NO), was added into UW solution during the cold storage period. NO is a vasodilator that acts on hepatic microvascular system protecting the liver from preservation/reperfusion injuries. We compared five groups of livers: a) Normal control (livers neither stored nor reperused); b) cold stored livers for 48 Hs in UW; c) cold stored livers for 48 Hs in UW + 50  $\mu$ M GSNO; d) cold stored livers for 48 Hs in UW + 100  $\mu$ M GSNO; e) cold stored livers for 48 Hs in UW + 250  $\mu$ M GSNO. Groups **b**, **c** and **e** were reperused after cold storage. Pieces of rat livers were histological processed for paraffin embedded after reperfusion. Liver slices were stained with Direct Red 80 – Fast Green and analyzed with polarized microscopy to differentiate collagen type I from type III.

Except livers from group a that showed abundant collagen fibers (collagen type I: red, yellow, white or orange – thin white arrows; collagen type III: green – white empty arrows), all the other groups present less amount of collagen fibers. While collagen type III seem not to be affected by the treatment (cold storage/reperfusion), collagen type I was drastically diminished. Livers from group d conserved some fibers of collagen type I, while the ones of groups **b**, **c** and **e** practically lost them. It seemed that 100  $\mu$ M was the preeminent concentration of GSNO to prevent the complete lost of collagen type I during cold storage in UW solution following by reperfusion.