



Concise reviews

Usefulness of neutrophil-to-lymphocyte ratio (NLR) as a prognostic predictor after treatment of hepatocellular carcinoma. Review article

Mohamad Mouchli^{a,b,c,d,*}, Shravani Reddy^b, Miranda Gerrard^e, Lisa Boardman^c, Marrieth Rubio^{a,b}



^a Virginia Tech Carilion School of Medicine Department of Internal Medicine, Division of Gastroenterology & Hepatology, Roanoke, VA, United States

^b Virginia Tech Carilion School of Medicine Department of Internal Medicine, Roanoke, VA, United States

^c Mayo Clinic, Division of Gastroenterology & Hepatology, Rochester, MN, United States

^d Cleveland Clinic Foundation, Division of Gastroenterology & Hepatology, Cleveland, OH, United States

^e Virginia Tech Carilion School of Medicine, Roanoke, VA, United States

ARTICLE INFO

Article history:

Received 17 May 2020

Accepted 8 August 2020

Available online 5 September 2020

Keywords:

NLR

HCC outcomes

LT

Resection

Chemotherapy and other therapies

ABSTRACT

The neutrophil-to-lymphocyte ratio (NLR) is an inflammatory marker which has been investigated as a prognostic indicator in post-therapeutic recurrence and survival of patients with HCC. Our aim was to review all studies that assessed the prognostic value of pre-treatment NLR in predicting patient survival, cancer recurrence, and graft survival in patients undergoing various therapies for HCC. We searched the database of PubMed and Google Scholar to review all studies that have the word "NLR" and the word "HCC." We included all studies that assessed pre-treatment NLR as a prognostic factor in predicting outcomes in HCC patients. We excluded studies that assessed the correlation between post-treatment NLR or dynamic changes in NLR after treatment and HCC outcomes in an effort to minimize the confounding effect of each treatment on NLR. We reviewed 123 studies that studied the correlation between pre-treatment NLR and patient survival, 72 studies that evaluated the correlation between pre-treatment NLR and tumor recurrence, 21 studies that evaluated the correlation between NLR and tumor behavior, and 4 studies that assessed the correlation between NLR and graft survival. We found a remarkable heterogeneity between the methods of the studies, which is likely responsible for the differences in outcomes. The majority of the studies suggested a correlation between higher levels of pre-treatment NLR and poor outcomes. We concluded that NLR is a reliable and inexpensive biomarker and should be incorporated into other prognostic models to help determine outcomes following HCC treatment.

© 2020 Fundación Clínica Médica Sur, A.C. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

HCC is the fifth most frequently diagnosed malignant tumor and the fourth most common cause of cancer-related deaths worldwide [1]. HCC incidence and death rates are increasing worldwide in comparison to declining death rates of other common cancers (colon, breast, lung and prostate) [1]. In the United States, the mortality rate of HCC increased by 43% between 2006 and 2016 with a 5-year survival of 18% [2,3].

HCC recurrence is associated with morbidity and mortality, and the rates range from 10 to 30% [4]. There are multiple factors associated with HCC recurrence, with the most important being the presence of macrovascular invasion and distant metastasis [2,3,5]. Often assessment of metastatic disease requires invasive interventions while imaging studies are of limited use in some scenarios [6]. Alpha-fetoprotein (AFP) is the only biomarker which was studied extensively and was found to be useful in predicting tumor recur-

Abbreviations: FU, 5-fluorouracil; ALBI, Albumin-bilirubin index; AFP, Alpha-fetoprotein; ALRI, Aspartate amino transferase-lymphocyte ratio index; BCCLC, Barcelona clinic liver cancer; CTP, Child-Turcotte-Pugh; c-TACE, Conventional chemo embolization; DCP, Des-gamma-carboxy prothrombin; DEE, Drug-eluting embolics; GPS, Glasgow Prognostic Scale; HALT-HCC, Hazard Associated with LT for Hepatocellular Carcinoma; HBV-HCC, Hepatitis B-associated hepatocellular carcinoma; HCC, Hepatocellular carcinoma; HIV, Human immune deficiency virus; IFN, Interferon; IL, Interleukin; LT, Liver Transplantation; MiC, Milan Criteria; MELO-NA, Model for End Stage Liver disease-Sodium Levels; MORAL, Model of Recurrence after LT; NLR, Neutrophil-to-lymphocyte ratio; PLR, Platelets-to-lymphocytes ratio; PI, Predictive index; PNI, Prognostic nutritional index; RFA, Radio frequency ablation; SIR, Systemic inflammatory response; TACE, Trans arterial chemo embolization; TAMs, Tumor associated macrophages; VEGF, Vascular endothelial growth factor; Y90, Yttrium-90.

* Corresponding author.

E-mail address: MOUCHLM@ccf.org (M. Mouchli).

rence after liver transplantation (LT) for HCC [7]. This biomarker has its limitations since about 50% of HCC patients do not secrete AFP [8]. A reliable biomarker to better predict outcomes in HCC is needed.

Markers of systemic inflammatory response such as the neutrophil to lymphocyte ratio (NLR) are thought to be useful predictors of survival in patients diagnosed with liver conditions and different cancers such as cholangiocarcinoma, esophageal, pancreatic, gastric, and colorectal cancers [9–19]. Halazun et al. first described the relationship between NLR and HCC, demonstrating that an elevated NLR (>5) was associated with poor overall survival and high recurrence in patients undergoing LT for HCC [20]. Later on, several studies demonstrated that NLR was significantly associated with overall and disease-free-survival after different treatments including transarterial chemoembolization (TACE), radiofrequency ablation (RFA), and resection. These treatments demonstrated hazard ratios ranging between 1.16–4.22 [21–26].

NLR levels fluctuate after treatment, and this fluctuation was reported to have a prognostic value in patients with HCC [24]. Assessing the prognostic value of post-treatment NLR or changes in NLR after treatments would be biased due to the effect of confounding factors related to treatment. This review will focus on studies assessing the utility of baseline NLR as a predictor of post-LT graft survival, cancer recurrence, patient survival, and tumor progression of patients with HCC undergoing therapy.

2. Methods

We conducted a comprehensive search at Virginia Tech Carilion School of Medicine to identify retrospective studies, prospective studies, systematic reviews, meta-analyses, and abstracts which included key terms such as "HCC" and "NLR" between 2008 and 2020. We excluded studies that evaluated other biomarkers such as platelet-to-lymphocyte ratio (PLR). Additional manual searching was also conducted on journals and proceedings of key meetings. We identified 216 eligible studies meeting the search criteria. We narrowed down the results by the treatment received: LT, surgery or hepatectomy, microwave ablation, RFA, TACE, hepatic artery infusion chemotherapy, and chemotherapy. The rest of the studies were excluded since they evaluated the correlation between treatment outcomes and post-treatment NLR or NLR dynamics.

3. Results

We identified 24 studies that evaluated patient survival after LT and 23 studies that evaluated tumor recurrence after LT. Forty-one studies assessed survival after resection, and 24 assessed recurrence after resection. Forty-one studies addressed overall survival following locoregional therapies in addition to 19 studies that addressed recurrence after such therapies. Seventeen studies and 5 studies evaluated overall survival and tumor recurrence, respectively. There was an overlap between studies that assessed survival and studies that assessed tumor recurrence. Only 4 studies evaluated the relationship between graft survival and pre-treatment NLR.

4. Patient survival

4.1. Overall survival following LT

Several Eastern and Western studies have demonstrated that an elevated NLR is associated with poor overall survival in patients who have undergone LT for HCC [20,27–35]. In 2009, Halazun first

described the association between elevated NLR and recurrence-free and overall survival in 150 patients undergoing LT using an NLR cutoff >5 [20]. Several studies thereafter supported this finding using different cutoffs. The first one by Limaye et al. observed the correlation between elevated NLR and poor survival and reported that the five-year cumulative survival of the elevated NLR group was only 38%, compared to a five-year cumulative survival of 68% in the low NLR group [30]. The study also showed a correlation between high NLR and relative lymphopenia suggesting the mechanism of immune response but did not find factors such as tumor size and macrovascular invasion to predict survival. This is likely due to small sample size and unavailability of data on patients who received locoregional therapy. Another study by Ismael et al. reported better LT outcomes in the low NLR group, but the results did not reach statistical significance [36]. In this study, the sample size was small, and there were several confounding factors such as using TACE in the majority of patients and choosing three months as the window for CBC to calculate NLR. Yang et al. reported improved overall and recurrence free survival after LT in hepatitis B-associated hepatocellular carcinoma (HBV-HCC) patients with separate or combined low NLR and low platelet-to-lymphocyte ratio (PLR) values [37].

NLR was also assessed in living donor LT. In a study by Hamamoto et al. that evaluated 216 patients undergoing living donor LT, pre-LT NLR ≥ 2.66 was found to be a risk factor associated with poor disease-free survival, and the 3-year disease-free survival of patients with NLR ≥ 2.66 was 74.4% compared to 92.8% in patients with NLR < 2.66 [35]. In this study, NLR was found to be superior to other inflammatory markers. Another study evaluated the role of NLR in living donor LT and found that preoperative NLR ≥ 2.85 correlated with improved graft and patient survival when evaluated in about 2000 patients undergoing living donor LT [38].

The prognostic accuracy of NLR improves with use in prognostic models combining other factors. Fu et al. reported that combining baseline NLR with baseline plasma fibrinogen increased the prognostic accuracy to predict disease-free and overall survival [39]. Wang et al. incorporated NLR into a model comparable to The Milan Criteria (MiC) to predict overall and transplant-free survival for male recipients selected for LT [33]. Xiao et al. concluded that a scoring system based on elevated NLR and Hangzhou criteria is superior to other criteria in selecting LT candidates for HBV-HCC patients [40]. Halazun et al. designed the Model of Recurrence after LT (MORAL) score that included NLR and was superior to MiC in predicting recurrence after LT for HCC [41]. Na et al. combined NLR and C-reactive protein (CRP) with MiC to select patients with HCC beyond MiC for living donor LT [29]. NLR has been incorporated into measures which have been developed into an entirely novel scoring protocol known as the Hazard Associated with LT for Hepatocellular Carcinoma (HALT-HCC), which shows promise in predicting mortality in LT recipients for HCC [42]. McVey et al. reported that NLR predicted overall survival in patients on the waitlist but lost its power when controlled with Model for End Stage Liver Disease-NA (MELD-NA) [43].

Despite the reported correlation between high NLR and poor outcomes in the studies we reviewed above, a few studies did not show a correlation between elevated NLR and post-LT survival. Sullivan et al. studied 75 patients with median follow-up of 56 weeks and did not find NLR to be predictive of future treatment or to correlate with short-term overall survival [44]. In this study, only Child-Turcotte-Pugh (CTP) score but not MELD score correlated with overall survival [44]. Shindoh et al. reported that NLR had limited prognostic impact and was inferior to AFP or des-gamma-carboxyprothrombin (DCP) in predicting overall survival [45]. Parisi et al. studied 150 patients to evaluate the role of inflammatory markers including NLR in predicting overall and recurrence-free survival in patients within MiC. NLR did not predict such outcomes

in those with LT but predicted response to transarterial embolization in some patients [46].

4.2. Overall survival following resection

The impact of an elevated preoperative NLR ratio on outcomes after resection for HCC was first reported by Gomez et al. in 2008 [23]. Gomez et al. found NLR ≥ 5 to be an independent predictor of poorer disease-free survival [23]. The results of other studies that evaluated the correlation between preoperative NLR and outcomes following tumor resection showed a less clear relationship and variability between the studies. Many studies supported Gomez et al's findings and demonstrated that high preoperative NLR was an independent, significant, and early predictive factor of early mortality after resection for HCC with a hazard ratio (HR) ranging from 1.031 to 4.9 [47–66]. NLR and visceral fat mass were both found to be independent risk factors for overall survival in 215 patients with surgically resected HCC [67]. In a prospective study of 1374 subjects, NLR was found useful in discriminating HCC survivors after resection from non-survivors but was not found to be a risk factor predictive of survival on multivariate analysis [68]. Only age and diabetes mellitus were found to be negative predictors of survival [68].

High NLR was able to predict survival in HCC patients with low AFP levels and patients with associated co-morbid conditions such as portal vein thrombosis. Shiraki found age and NLR > 3.2 to be good markers for survival in 478 HCC patients undergoing resection with normal preoperative AFP levels [69]. Another retrospective study reported that NLR with a cutoff of 2.1 predicted overall survival and discriminated outcomes after surgical resection of small solitary HCC in 222 patients with AFP $< 400 \text{ ng/mL}$ [58]. High NLR (> 2.9) was also found on multivariate analysis to be a good predictor of poor prognosis in 81 patients with HCC and portal/hepatic vein thrombosis undergoing resection. High NLR values correlated with CTP and the maximum diameter of the tumor [70]. Also, in a retrospective study, NLR > 2.3 was found to be a good predictor of survival after two-stage procedures combining reductive surgery and percutaneous isolated hepatic perfusion in 42 Japanese patients with multiple HCCs and with portal vein tumor thrombus [71].

NLR prognostic values were reported to vary based on several factors including ethnicity, degree of fibrosis, tumor stage, and etiology of liver disease. Wong et al. reported that high NLR was associated with disease severity and poor outcomes after resection, but NLR values were confounded by factors such as ethnicity and etiology of chronic liver disease [72]. Wang et al. focused on the effects of three inflammatory markers [NLR, PLR, and prognostic nutritional index (PNI)] in 234 HBV-HCC patients to minimize the effect of ethnicity as a confounding factor and only found NLR to be a significant prognostic factor independent of tumor size [55]. In this study, the impact was more prominent in Ishak stage 0–5 patients compared with Ishak stage 6 patients [55]. Another study assessed 108 Serbian patients and supported the later study by finding NLR to have a prognostic value in noncirrhotic (low fibrosis score) HCC patients, but the role was not confirmed in cirrhotic patients [73]. The role of NLR was compared in a small study between 81 normal and 61 diseased livers from different etiologies (viral hepatitis alone or in combination with fibrosis or cirrhosis) and was found to be an important risk factor to predict survival after HCC resection in diseased livers but not in normal livers [74]. The study suggested two different disease entities between the two groups, but the study was limited by small sample size and confounded by the inclusion of different etiologies and degrees of fibrosis.

A few studies were done to assess the role of NLR in different HCC stages. One found NLR to be a good factor to predict overall survival after hepatectomy in early and intermediate stage HCC but not

advanced stage HCC [57]. Another study reported that high preoperative NLR at a cut-off value of 2.8 predicted overall survival after resection for TNM stage I tumors but not stage II or III [75] and was associated with greater proportion of patients with extrahepatic disease recurrence in stage I tumors [75]. NLR was combined with other inflammatory markers and integrated into predictive models to predict survival after resection of early stage HCC. It was combined with PLR [76] and fibrinogen to predict overall survival in HCC patients with BCLC stage 0–I undergoing curative resection [77]. Ren et al. established a predictive index (PI) model that included NLR to predict 5-year survival for HCC patients after radical resection [78]. In the same study, postoperative TACE treatment was found to be a protective factor for 5-year survival after surgical resection.

Many other studies reported conflicting data and did not find a strong association between preoperative NLR and overall survival. Some did not find a correlation between NLR and predicted survival in large HCCs ($\geq 10 \text{ cm}$) undergoing resection [79]. Chan et al. studied the role of PNI, PLR, and NLR in 324 patients with early stage HCC undergoing resection and did not find elevated NLR (≥ 5) or PLR to be significant prognostic factors for overall survival [80]. A large western study retrospectively assessed 370 patients and found that PLR but not NLR was independently associated with worse overall and recurrence-free survival [81]. NLR score was also compared to other inflammation-based prognostic scores and was found to be inferior to Glasgow Prognostic Scale (GPS) score in predicting overall survival after resection [82,83]. Peng et al. and Dai et al. compared preoperative NLR to postoperative NLR and revealed that increased postoperative NLR but not preoperative NLR was associated with poor overall survival [84,85]. A study evaluated lower NLR values and found that a preoperative NLR value > 2 did not independently predict poorer overall survival in patients with combined HCC and cholangiocarcinoma [86]. This conflicting data illustrates that the role of NLR use in predicting outcomes after resection in some groups (huge HCC, different fibrosis scores, or combined HCC and cholangiocarcinoma) requires further evaluation.

4.3. Overall survival following locoregional therapies

4.3.1. Microwave ablation and cryoablation

Increased NLR was identified as a risk factor for overall survival for patients with HCC treated with microwave ablation [87]. Compared to/Excluding CTP grading and tumor differentiation, pre-treatment NLR prior cryoablation predicted survival in 150 patients receiving cryoablation [88]. NLR with a cut off of 2.14 was studied in 506 recurrent HCC patients treated with thermal ablation and was found to correlate with a statistically significant difference in recurrence-free survival and recurrence rates [89].

4.3.2. Radiofrequency ablation (RFA) and radio embolization

Pre-ablation NLR predicted survival in early stage and locally advanced HCC patients receiving RFA [90–93]. Sukato et al. found elevated NLR along with low albumin levels and high AFP levels to be independent predictors of poor survival [94]. A combination of the NLR and the PLR was superior to the NLR alone in predicting outcomes after RFA [95].

NLR dynamics and postoperative NLR were compared to preoperative NLR in 3 studies. Two studies described that NLR dynamics, but not pre-treatment NLR, was an independent prognostic factor for overall survival [96,97]. Post-treatment NLR, but not pre-treatment NLR was found to be a predictive marker of overall survival in HCC patients with viral hepatitis treated with RFA [98].

4.3.3. Hepatic arterial infusion chemotherapy

In patients with advanced HCC undergoing hepatic arterial infusion with cisplatin plus continuous 5-fluorouracil, NLR < 4 is found to be associated with prolonged survival [99]. Mizukoshi

et al. reported longer overall survival in HCC patients with low pretreatment NLR undergoing arterial infusion using interferon (IFN)/5-fluorouracil (FU) or IFN/FU and cisplatin [100]. Baseline NLR and early DCP change ratio predicted treatment response and progression free survival after hepatic arterial infusion of chemotherapy in advanced HCC [101]. Lastly, NLR was a good marker for predicting disease progression in HCC patients receiving intra-arterial therapies (drug-eluting embolics (DEE), conventional chemoembolization (c-TACE), or Yttrium-90 (Y90) glass microspheres) [102,103].

4.3.4. Transarterial chemoembolization (TACE)

High NLR was a predictor of poor survival in patients with unresectable intermediate- or advanced-stage HCC who received TACE [22,104–108]. The increase in NLR after treatment, not pretreatment NLR indicated favorable outcomes in some studies and poor outcomes in other studies [22,104].

Many studies assessed the prognostic value of NLR after TACE after combining it with other inflammatory factors. Oh et al. found NLR to be an important survival indicator when combined with CRP [109]. Higher pretreatment absolute lymphocyte count and lower pretreatment NLR were associated with favored overall survival in 93 HCC patients receiving TACE with doxorubicin-eluting microspheres [110]. High NLR was reported to predict metastasis after treatment with TACE for recurrent HCC and was inferior to PLR in predicting 1-year overall survival [111]. Baseline NLR (≥ 5) in non-viral HCC patients who underwent TACE was associated with poor survival when combined with diabetes mellitus [112]. A score that included both NLR and PNI was superior to NLR alone in predicting overall survival in patients with unresectable HCC after TACE [113]. He et al. found that neutrophil/platelet-to-lymphocyte ratio score (a score combining NLR and PLR) is an accurate score in predicting survival after TACE therapy [114]. Chon et al. incorporated NLR and other prognostic factors into SNAVORN score to predict mortality in HCC patients after receiving TACE therapy [115].

Zhou et al. demonstrated that the GPS, when compared to other inflammatory scores, is superior to other inflammation-based prognostic scores including NLR for predicting survival for HBV-HCC patients after TACE [116]. Aspartate aminotransferase-lymphocyte ratio, not NLR, was found to be a negative predictive factor for overall survival in 189 HBV-HCC patients following TACE [117]. Changes in NLR, not AFP nor pretreatment NLR, were good markers to predict survival after TACE [118].

4.3.5. Mixed treatments

In patients with advanced HCC who received TACE and Sorafenib, NLR was an independent factor for survival, and the median survival of patients with $\text{NLR} > 3$ was 14 months compared to 26 months for patients with $\text{NLR} < 3$ [119]. It also correlated with poor outcomes in HCC patients who received curative resection with adjuvant Sorafenib [120]. Another study found that poor survival was related to $\text{NLR} \geq 3.09$ in young patients treated with TACE and RFA [121]. $\text{NLR} \geq 1.6$ was a predictive factor of poor survival in patients treated with chemoembolization and sequential resection [122]. Elevated pretreatment $\text{NLR} \geq 2.5$ indicated poor prognosis for patients with unresectable HCC treated with 131I-labeled-metuximab plus TACE [123]. Performing preoperative adjuvant TACE did not impact overall survival, but baseline $\text{NLR} > 4$ did when studied in patients undergoing radical therapies (resection, RFA, or LT) [107]. A study by Xiao et al. identified an association between high NLR and poor overall survival in patients with HCC receiving mixed treatments [26]. Dynamic changes in NLR, but not NLR itself, have a prognostic value in HCC patients with portal vein thrombosis undergoing microwave ablation after TACE [124].

4.4. Overall survival following chemotherapy

Many studies described a correlation between low NLR and good overall survival for patients with intermediate-advanced HCC receiving Sorafenib [125–136]. Diaz-Beveridge et al. identified baseline NLR, CTP score, early-onset diarrhea, and performance status as prognostic factors for overall survival [138]. A study found that peritumoral tissue NLR was higher than that in the intratumoral tissue and correlated negatively with overall survival [139]. In a recent study, NLR of approximately 3.0–4.5 was associated with overall survival in unresectable HCC treated with lenvatinib [140]. Sprinzl et al. found that CRP-based scores were superior for predicting overall survival when compared to cell frequency-based inflammation scores such as NLR [141].

4.5. NLR and tumor characteristics/behavior

Baseline NLR is associated with tumor characteristics, MELD stage, CTP class, Barcelona clinic liver cancer (BCLC) stages, tumor size, tumor grade and differentiation, number of tumors, high level of serum AFP, microvascular invasion, low albumin, and high GGT [26,51,53,72,83,142–148]. Baseline NLR was good predictor of 5-year survival in all HCC cirrhosis patients and overall survival in HCC patients with non-B, non-C cirrhosis [149]. Low NLR was associated with favorable prognostic factors such as lower liver chemistries; AFP; and decreased incidence of metastasis, portal vein thrombosis, and ascites [150]. Elevated NLR correlated with dropout from LT due to patient death, tumor progression, beyond conventional criteria for transplantability, or acute liver failure development after TACE treatment [151]. NLR was also studied as a factor to improve specificity/sensitivity of AFP for diagnosis of HBV–HCC in a Chinese population of 473 patients [152]. Interestingly, preoperative NLR correlated with the development of HCC in HBV cirrhotic patients who underwent splenectomy [153].

The effect of NLR on tumor progression is suggested through several mechanisms. The most common theory suggests that high neutrophil production probably enhances the propensity for tumor progression and vascular invasion by increasing the production of vascular endothelial growth factor (VEGF) [154,155]. Other studies reported high NLR to correlate with CD163-positive tumor-associated macrophages (TAMS), peritumoral CD163 and IL-17-expressing cells, and PD-L1 expression in the tumor center [34,54,55].

4.6. Cancer recurrence and disease progression

4.6.1. Cancer recurrence following LT

Several studies evaluated overall survival after LT for HCC assessed cancer recurrence after LT. Many concluded that NLR may also be valuable in predicting HCC recurrence following LT [28,30,32,33,35,39,43]. As early as 2011, a retrospective study by Bertuzzo et al. concluded that NLR more than or equal to 5 was critical in identifying successful LT candidates based on their risk for recurrence since it predicted overall survival and recurrence-free survival [31]. A large group of LT recipients were evaluated by Agopian et al., which identified NLR as the most powerful tool for HCC recurrence over thirty years of data [156].

Several scoring systems were developed consequential of the powerful prognostic capabilities of NLR to predict recurrence after LT. The MORAL score, which stemmed from a prospective cohort study by Halazun et al., is a popular scoring system [41]. This study found that preoperative $\text{NLR} \geq 5$, $\text{AFP} > 200$, and tumor size > 3 cm are independent risk factors for poor recurrence-free survival [41]. Bodzin established a risk score model that included baseline NLR to predict survival in patients undergoing LT for HCC [157]. Na et al. designed another scoring system using pretreatment NLR and CRP

to predict recurrence in HCC beyond the MiC in patients receiving living donor LT [29]. Wang et al. evaluated a model that included NLR > 4 in 248 male patients receiving LT, and certain ranges were associated with early tumor recurrence [33]. Traditionally utilized MELD scores were also supplemented with NLR as a prognostic tool to predict recurrence-free survival [30].

High NLR was also used in certain groups, including patients with recurrent HCC and patients undergoing living donor LT. A study by Motomura et al. evaluated 158 patients and suggested that NLR augments prognostic abilities in predicting recurrence-free survival in HCC patients undergoing living-donor LT [34]. In this study, the levels of tumor expression of inflammatory markers such as VEGF, interleukin (IL)-8, IL-17, CD68, and CD163 were measured in the high NLR group and the low NLR group and found to be similar in both groups except for peritumoral and circulating IL-17 and peritumoral CD163-positive TAMs [34]. Yoshizumi et al. observed the same findings in three studies evaluating small groups of patients with primary or recurrent HCC undergoing living-donor LT and suggested to reconsider LT in patients with NLR above 4 [158–160]. Harimoto et al. revealed that high baseline NLR predicted more frequent recurrence after LT, and high NLR at recurrence was further associated with mortality of recurrent HCC patients [35]. The study also found elevated DCP and a short interval between last HCC treatment and living donor LT to be risk factors for recurrence [35].

Two studies by Parisi et al. and Lai et al. found that elevated NLR (NLR \geq 5 and NLR \geq 5.4, respectively) was not a significant predictor of HCC recurrence [46,151]. The former indicated that the absence of neoadjuvant therapy and being outside MiC are significant risk factors associated with poor recurrence-free survival [46]. The latter found NLR \geq 5.4 to be a predictor of dropout from the LT list but not a risk factor for recurrence [151]. Another study evaluated 124 patients and found that the impact of NLR in predicting recurrence after living-donor LT is limited but suggested the LT decision to rely on Tokyo criteria in addition to AFP or DCP values [45]. Liese et al. did not find NLR to be a significant risk factor for recurrence in 92 patients undergoing LT when evaluating the role of microRNA (miRNA) microarray analysis in predicting HCC recurrence after LT [161]. However, these negative studies comprise of fairly small sample sizes, which calls into question their overall conclusions when compared to better powered studies such as those discussed above.

4.7. Cancer recurrence following resection

The association between high preoperative NLR and short recurrence-free survival and more frequent recurrence was identified by many research groups, irrespective of treatment choice [23,51,53–56,58,59,65,66,162,163]. Therefore, pretreatment NLR was identified as an independent prognostic factor for recurrence-free survival in the general group [65,162], early or intermediate stage HCC [57], and late-stage or large HCC [70]. Najjar et al. estimated the hazard ratio of this association to range from 1.32 to 2.59 [47]. Interestingly, NLR was an accurate factor to predict extrahepatic recurrence in early stage HCC patients undergoing resection [75].

Combining other inflammatory markers with NLR increased its prognostic values in some studies. Kong et al. reported that NLR combined with fibrinogen predicted recurrence free survival in HCC patients with BCLC stage II–III [77]. NLR predicted recurrence more accurately when combined with PLR [76].

Contrary results are reported from Western patients [81], Eastern patients with early-stage HCC undergoing resection [80], and patients with different etiology of HCC such as HBV-HCC [55,164]. Few series found the dynamic change in NLR or postoperative

NLR, but not preoperative NLR, was an independent risk factor for recurrence-free survival [53,85,165,166].

4.8. Cancer recurrence following locoregional therapies

Monocyte-granulocyte/lymphocyte ratio was found to be a good predictor for early HCC recurrence after TACE [167]. High baseline NLR was found to have a role in early recurrence, response to treatment, and disease progression in patients receiving thermal ablation [89], transarterial infusion [105], RFA [90,93,95,98,168], TACE [101–103,107,109–111], hepatic arterial infusion chemotherapy [105], and other therapies for HCC including combined therapies [107,120,122,169,170].

Other studies did not support the relationship between preoperative NLR and recurrence-free survival or time to progression in HCC patients receiving microwave ablation [87], RFA [92,96–98], or TACE [104] but associated other factors with poor outcomes such as elevated baseline AFP, post-treatment NLR, and NLR dynamics after treatment. Noninvasive Fibrosis Marker such as aspartate aminotransferase-to-platelet ratio index, but not NLR, was found to correlate negatively with tumor recurrence in 98 HCC patients undergoing TACE [171].

4.9. Cancer recurrence after systemic chemotherapy

Several studies addressed the correlation between elevated baseline NLR and short recurrence-free survival or time to progression in HCC patients receiving systemic chemotherapy [127,129,134,136].

4.10. Cancer recurrence in HCV-patients treated with direct-acting antivirals

Casadei et al. illustrated that albumin-bilirubin index (ALBI), platelet count, and aspartate aminotransferase-lymphocyte ratio index (ALRI) correlated with HCC recurrence or occurrence in 416 HCV patients treated with direct-acting antivirals. [172]. Although NLR did not correlate with recurrence, it was shown to increase during treatment with direct-acting antivirals [172].

4.11. Graft dysfunction

Though multiple studies have observed the association of preoperative NLR with outcomes after kidney and heart transplantation [173–176], there are limited studies which have examined elevated NLR and post-transplantation graft dysfunction or survival in patients with HCC. The studies which evaluated the correlation between NLR and graft outcomes included patients undergoing LT of all causes and did not focus on LT for HCC. NLR was high in patients who developed early graft dysfunction after living donor LT [177]. Lin et al. found a correlation between NLR (using a cutoff of 4.6) and graft survival. In this study, the high NLR group developed a greater number of infectious complications compared to the normal NLR group [178]. There has been an associated increase in early graft dysfunction on post-op day 7 as well as 1-year graft failure for those who have preoperative NLR \geq 2.85 after living donor LT [38]. In the same study, NLR was found to be superior to other parameters such as CRP, procalcitonin, PLR, and GPS in predicting graft outcomes.

Though we focus on graft dysfunction and survival after LT, poor post-operative outcomes of small liver grafts have also been linked to elevation in NLR. A study by Hayashi et al. described the association of NLR with the clinical course of patients who underwent living donor LT with different size grafts [179]. Interestingly, the group with small graft size showed higher NLR in post-operative days 3–10 compared to the patients in the group with large graft

size. The results illustrated that total bilirubin and NLR varied most between the two groups between day 3 and day 10 after surgery. However, post-operative complications were not statistically different between the two groups. Small-graft syndrome is mostly associated with endothelial injury related to ischemia/reperfusion which can lead to many inflammatory changes. The relationship to NLR may be helpful in understanding the post-operative course of patients undergoing LT with small grafts.

5. Discussion

Predicting HCC recurrence and survival after treatment would allow us to better select patients and improve utilization of expensive resources. Currently, AFP is the only marker which has been identified, and it is only helpful in about 50 % of cases. Systemic inflammatory response (SIR) is widely considered as a preoperative risk factor for poor outcomes in patients with HCC.

The studies examined in this paper describe the successful use of NLR as an indicator of survival in patients post-liver transplant and post-resection. NLR has also been studied accompanying other treatment methods such as chemotherapy. Overall, studies show a correlation between good patient and graft outcomes and low NLR values as well as between poor patient and graft outcomes and high NLR values. Furthermore, NLR acts as an accurate indicator of disease recurrence, with high values correlating to increased chance of recurrence.

The neutrophil-to-lymphocyte ratio (NLR) is one modality of measuring inflammation that is relatively inexpensive and readily available from a complete blood count with differential. NLR is simple to calculate by dividing the number of neutrophils by the number of lymphocytes from a peripheral blood sample. Our review illustrates that it should be considered as a solo or combined prognostic factor in predicting outcomes in HCC patients undergoing LT from all causes [29,33,39,41]. Its use as a prognostic factor in predicting overall survival and recurrence-free survival was evaluated in several meta-analyses, including the studies we reviewed in our article. In a meta-analysis performed by Sun et al., ten studies demonstrated that LT for HCC patients with elevated NLR was associated with reduced overall survival compared to those in the low NLR group (HR = 2.71, 95 % CI: 1.91–3.83) [180]. It also concluded that there is a correlation between the increase in the cutoff value of NLR and the increase in the hazard ratio for overall survival. This correlation was not altered by the types of LT [180]. Additional meta-analyses concluded that disease-free survival is poorer in patients with higher NLR (Hazard ratios were 3.61 and 3.42, respectively) [32,47,180,181]. The positive correlation between lower relapse-free survival and higher NLR post-LT has a reported hazard ratio ranging from 1.09–67 [47]. These results suggest its use in predicting outcomes in HCC patients undergoing LT, but the cutoff is to be determined.

While different cutoffs were used in different studies for defining elevated NLRs, a cutoff above 4 has shown to reliably predict poor outcomes after LT [31,182]. The difference in outcomes between different cutoffs was compared by Xiao et al., and the greatest difference in outcomes was noticed between patients with NLR 3–4 compared to those with NLR 4–5. The 1-, 3-, and 5-year overall survival rates in patients with NLR between 3–4 versus NLR 4–5 were 86.4%, 57.3%, and 54.3% vs 79.7%, 35.5%, and 31.1%, respectively [182]. Cescon et al. reported that the 5-year overall survival rates ranged between 14–57% in the high preoperative NLR group and 62–84% in the low preoperative NLR group [183]. Different cut-offs are limiting factors in existing studies and have likely interfered with results.

Preoperative NLR has also been studied alone and in combination with other inflammatory markers with variable results and

has a less clear role in comparison to post-LT NLR. Several studies have shown a predictive value in patients undergoing resection, but this relationship is affected by ethnicity, etiology of liver disease, and tumor size and stage. Combining NLR with other inflammatory markers improved its prognostic value [76,77]. Future prospective studies should focus on recruiting more patients and addressing confounding factors to define the role and address unanswered aspects of preoperative NLR on outcomes after resection.

In our review, we found evidence of a good correlation between a low NLR and recurrence-free survival after microwave ablation and cryoablation using a cut off of 2.14 [89]. We also found a correlation after RFA, and it was used in combination with PLR [95]. When using hepatic arterial chemotherapy, NLR and early DCP change ratio predicted treatment response [101]. To assess outcomes after TACE, NLR was combined with CRP and absolute lymphocyte count and incorporated into the SNAVCORN score [109](115). A meta-analysis that included all studies evaluating baseline NLR and dynamic changes in NLR did not find NLR to be predictive of recurrence-free survival after RFA [90]. All these studies are supporting that the prognostic accuracy of NLR improves when used in combination with other markers as part of a panel in patients undergoing locoregional therapies. Further studies are needed to assess the role in such patients since it was not studied extensively when compared to other treatments.

NLR should be considered a good marker to assess survival and recurrence in patients receiving Sorafenib. A recent meta-analysis of 13 studies reported that HCC patients with a lower baseline NLR have better response to Sorafenib [137]. It also concluded that low pretreatment NLR correlated with significantly better overall survival for patients treated with Sorafenib rather than Tivantinib and suggested an NLR of 3 to be the minimum cutoff value for pretreatment NLR to predict overall prognosis [137]. NLR was also predictive of survival after chemotherapy in combination with CTP score, early-onset diarrhea, and performance status [138]. Future studies should focus on NLR in patients receiving newer chemotherapy as it is not studied extensively compared to studies on NLR after LT and resection.

Furthermore, elevated NLR is associated unfavorable tumor characteristics including the presence of multiple tumors, vascular invasion, elevated AFP, cirrhosis, and presence of Hepatitis B surface antigen [184]. NLR is a significant independent predictor of survival in all HCC patients with extrahepatic metastasis [185]. NLR was also found to be a good predictor of survival and recurrence in HCC patients who have chronic conditions such as human immunodeficiency virus (HIV) [186]. This may be explained by high neutrophil production enhancing the propensity for tumor progression and vascular invasion by increasing the production of vascular endothelial growth factor and other cytokines.

The mechanism associating elevated NLR with HCC prognosis involves many factors but is still poorly understood. High NLR reflects relatively depleted lymphocytes in the blood and tumor, and depleted lymphocytes are associated with poor cancer-specific survival by impairing the host immune response against tumor cells [187]. Lymphocytes have a role in cytotoxic cell death and in producing cytokines to inhibit tumor cells [188]. Elevated NLR is associated with high infiltration of TAMs and high production of inflammatory cytokines, such as interleukin (IL)-6, IL-8, and IL-17, which promote systemic neutrophilia [189,190]. It is also considered as a reservoir for VEGF which promotes angiogenesis and tumor growth [191]. This suggests that the negative outcomes in patients with elevated NLR are mediated by an inflammatory microenvironment, rather than a single pathway.

NLR can be affected by many factors such as age, body mass index, weight loss, sepsis, steroid drugs, alcoholic fatty liver, poor nutrition, diabetes, and hemorrhage [192–195]. These factors should be considered when interpreting results. In addition, NLR

can be affected by treatment of chronic conditions such as hepatitis C and hepatitis B infections [196–198]. Controlling confounding factors and performing randomized controlled studies would help increase the specificity of the use of NLR as a prognostic factor. Nevertheless, NLR should still be considered as a promising and inexpensive marker deserving further investigation.

Our review was limited by the heterogeneity between the studies included. Some studies were retrospective and had very small sample sizes. Other studies focused on a specific ethnicity or gender. The use of several cutoffs in different studies for NLR is another important limiting factor. The presence of other confounding factors that could have elevated NLR and the failure to report previous treatments in many studies are other limiting factors that should be addressed in future studies.

In conclusion, NLR is a reliable, relatively inexpensive, and non-invasive biomarker and should be incorporated into prognostic models to help determine outcomes following HCC treatment. Overall, an elevated pre-treatment NLR generally suggests poor survival outcomes in HCC patients. NLR is most useful in conjunction with other markers of inflammation, and the patient's clinical course should be considered. Studies regarding NLR as a predictor of graft success or dysfunction are limited, but existing studies indicate a relationship between high NLR and graft-related complications, revealing a potential focus for future research on the role of NLR in evaluation of HCC patients.

Authors' contributions

M. Mouchli: reviewed studies and data, designed the study and participated in the writing of the manuscript

S. Reddy: reviewed studies and data and participated in the writing of the manuscript

M. Gerrard: reviewed studies and data and participated in the writing of the manuscript

L. Boardman: reviewed studies and data and participated in the writing of the manuscript

M. Rubio: reviewed studies and data, helped design the study and participated in the writing of the manuscript

Conflict of interest

The authors have no conflicts of interest to declare.

References

- [1] Altekruse SF, McGlynn KA, Reichman ME. Hepatocellular carcinoma incidence, mortality, and survival trends in the United States from 1975 to 2005. *J Clin Oncol* 2009;27(9):1485–91.
- [2] Villanueva A. Hepatocellular carcinoma. *N Engl J Med* 2019;380(15):1450–62.
- [3] Jemal A, Ward EM, Johnson CJ, Cronin KA, Ma J, Ryerson B, et al. Annual report to the nation on the status of Cancer, 1975–2014, featuring survival. *J Natl Cancer Inst* 2017;109(9).
- [4] Ma WJ, She WH, Chan ACY, Cheung TT, Fung JYY, Dai WC, et al. Validated model for prediction of recurrent hepatocellular carcinoma after liver transplantation in Asian population. *World J Gastrointest Oncol* 2019;11(4):322–34.
- [5] Cidores MJ, Lucena JL, de la Fuente S, Cuervas-Mons V. Serum biomarkers and risk of hepatocellular carcinoma recurrence after liver transplantation. *World J Hepatol* 2019;11(1):50–64.
- [6] Vibert E, Azoulay D, Hoti E, Iacopinelli S, Samuel D, Salloum C, et al. Progression of Alpha-fetoprotein Before Liver Transplantation for Hepatocellular Carcinoma in Cirrhotic Patients: A Critical Factor. *Am J Transplant* 2010;10(1):129–37.
- [7] Schraiber LD S, de Mattos AA, Zanotelli ML, Cantisani GPC, ABdM Brandão, et al. Alpha-fetoprotein level predicts recurrence after transplantation in hepatocellular carcinoma. *Medicine* 2016;95(3):e2478.
- [8] Bai D-S, Zhang C, Chen P, Jin S-J, Jiang G-Q. The prognostic correlation of AFP level at diagnosis with pathological grade, progression, and survival of patients with hepatocellular carcinoma. *Sci Rep-Uk* 2017;7(1):12870.
- [9] Mouchli MA, Singh S, Loftus Jr EV, Boardman L, Talwalkar J, Rosen CB, et al. Risk Factors and Outcomes of De Novo Cancers (Excluding Nonmelanoma Skin Cancer) After Liver Transplantation for Primary Sclerosing Cholangitis. *Transplantation* 2017;101(8):1859–66.
- [10] Summey R, Mouchli MA, Reddy S, Mir A, Walsh C, Rubio M. Long Term Mortality in Young Patients (<40 y/o) With Alcoholic Liver Cirrhosis After First Hospitalization: Carilion Clinic Experience: 986. *Am J Gastroenterol* 2018;113:S555.
- [11] Mouchli M, Singh S, Loftus EVJ, Talwalkar JA, Hasan B, Hamdeh S, et al. High Neutrophil-to-Lymphocyte Ratio Predicts Outcomes in Patients with Primary Sclerosing Cholangitis Undergoing Liver Transplantation: 834. *Am J Gastroenterol* 2016;111:S367.
- [12] Zhou Y, Wei Q, Fan J, Cheng S, Ding W, Hua Z. Prognostic role of the neutrophil-to-lymphocyte ratio in pancreatic cancer: a meta-analysis containing 8252 patients. *Clin Chim Acta* 2018;479:181–9.
- [13] Kishi Y, Kopetz S, Chun YS, Palavecino M, Abdalla EK, Vauthey JN. Blood neutrophil-to-lymphocyte ratio predicts survival in patients with colorectal liver metastases treated with systemic chemotherapy. *Ann Surg Oncol* 2009;16(3):614–22.
- [14] Sharaiha RZ, Halazun KJ, Mirza F, Port JL, Lee PC, Neugut AI, et al. Elevated pre-operative neutrophil:lymphocyte ratio as a predictor of postoperative disease recurrence in esophageal cancer. *Ann Surg Oncol* 2011;18(12):3362–9.
- [15] Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ. Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol* 2005;91(3):181–4.
- [16] Gomez D, Morris-Stiff G, Toogood GJ, Lodge JP, Prasad KR. Impact of systemic inflammation on outcome following resection for intrahepatic cholangiocarcinoma. *J Surg Oncol* 2008;97(6):513–8.
- [17] Shimada H, Takiguchi N, Kainuma O, Soda H, Ikeda A, Cho A, et al. High preoperative neutrophil-lymphocyte ratio predicts poor survival in patients with gastric cancer. *Gastric Cancer* 2010;13(3):170–6.
- [18] Templeton AJ, McNamara MG, Seruga B, Vera-Badillo FE, Aneja P, et al. Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic review and meta-analysis. *J Natl Cancer Inst* 2014;106(6), dju124.
- [19] Rashtak S, Ruan X, Druliner BR, Liu H, Therneau T, Mouchli M, et al. Peripheral neutrophil to lymphocyte ratio improves prognostication in Colon Cancer. *Clin Colorectal Cancer* 2017;16(2):115–23, e3.
- [20] Halazun KJ, Hardy MA, Rana AA, Woodland DCt, Luyten EJ, Mahadev S, et al. Negative impact of neutrophil-lymphocyte ratio on outcome after liver transplantation for hepatocellular carcinoma. *Ann Surg* 2009;250(1):141–51.
- [21] Zheng J, Cai JY, Li H, Zeng KN, He LY, Fu HY, et al. Neutrophil to lymphocyte ratio and platelet to lymphocyte ratio as prognostic predictors for hepatocellular carcinoma patients with various treatments: a meta-analysis and systematic review. *Cell Physiol Biochem* 2017;44(3):967–81.
- [22] Huang ZL, Luo J, Chen MS, Li JQ, Shi M. Blood neutrophil-to-lymphocyte ratio predicts survival in patients with unresectable hepatocellular carcinoma undergoing transarterial chemoembolization. *J Vasc Interv Radiol* 2011;22(5):702–9.
- [23] Gomez D, Farid S, Malik HZ, Young AL, Toogood GJ, Lodge JP, et al. Preoperative neutrophil-to-lymphocyte ratio as a prognostic predictor after curative resection for hepatocellular carcinoma. *World J Surg* 2008;32(8):1757–62.
- [24] Qi X, Li J, Deng H, Li H, Su C, Guo X. Neutrophil-to-lymphocyte ratio for the prognostic assessment of hepatocellular carcinoma: a systematic review and meta-analysis of observational studies. *Oncotarget* 2016;7(29):45283–301.
- [25] Zhou DS, Zhang YJ, Xu L, Zhou ZG, Huang JT, Chen MS. A monocyte/granulocyte to lymphocyte ratio predicts survival in patients with hepatocellular carcinoma. *Sci Rep-Uk* 2015;5.
- [26] Xiao WK, Chen D, Li SQ, Fu SJ, Peng BG, Liang LJ. Prognostic significance of neutrophil-lymphocyte ratio in hepatocellular carcinoma: a meta-analysis. *BMC Cancer* 2014;14:117.
- [27] Pravisan R, Mocchegiani F, Isola M, Lorenzin D, Adani GL, Cherchi V, et al. Controlling Nutritional Status score does not predict patients' overall survival or hepatocellular carcinoma recurrence after deceased donor liver transplantation. *Clin Transplant* 2020;34(3):e13786.
- [28] Wang GY, Yang Y, Li H, Zhang J, Jiang N, Li MR, et al. A scoring model based on neutrophil to lymphocyte ratio predicts recurrence of HBV-associated hepatocellular carcinoma after liver transplantation. *PLoS One* 2011;6(9):e25295.
- [29] Na GH, Kim DG, Han JH, Kim EY, Lee SH, Hong TH, et al. Inflammatory markers as selection criteria of hepatocellular carcinoma in living-donor liver transplantation. *World J Gastroenterol* 2014;20(21):6594–601.
- [30] Limaye AR, Clark V, Soldevila-Pico C, Morelli G, Suman A, et al. Neutrophil-lymphocyte ratio predicts overall and recurrence-free survival after liver transplantation for hepatocellular carcinoma. *Hepatol Res* 2013;43(7):757–64.
- [31] Bertuzzo VR, Cescon M, Ravaoli M, Grazi GL, Ercolani G, et al. Analysis of factors affecting recurrence of hepatocellular carcinoma after liver transplantation with a special focus on inflammation markers. *Transplantation* 2011;91(11):1279–85.
- [32] Xiao GQ, Liu C, Liu DL, Yang JY, Yan LN. Neutrophil-lymphocyte ratio predicts the prognosis of patients with hepatocellular carcinoma after liver transplantation. *World J Gastroenterol* 2013;19(45):8398–407.
- [33] Wang W, Ye Y, Wang T, Zhang F, Geng L, Yu J, et al. Prognostic prediction of male recipients selected for liver transplantation: with special attention to neutrophil to lymphocyte ratio. *Hepatol Res* 2016;46(9):899–907.
- [34] Motomura T, Shirabe K, Mano Y, Muto J, Toshima T, Umemoto Y, et al. Neutrophil-lymphocyte ratio reflects hepatocellular carcinoma recurrence after liver transplantation via inflammatory microenvironment. *J Hepatol* 2013;58(1):58–64.

- [35] Harimoto N, Yoshizumi T, Shimagaki T, Nagatsu A, Motomura T, Harada N, et al. Inflammation-based prognostic score in patients with living donor liver transplantation for hepatocellular carcinoma. *Anticancer Res* 2016;36(10):5537–42.
- [36] Ismael MN, Forde J, Milla E, Khan W, Cabrera R. Utility of inflammatory markers in predicting hepatocellular carcinoma survival after liver transplantation. *Biomed Res Int* 2019;2019:7284040.
- [37] Yang Z, Zhou L, Wu L, Zheng S. A novel classification of hepatocellular carcinoma based on combined neutrophil and platelet-to-Lymphocyte ratio: implications for prognosis after liver transplantation. *J Am Coll Surgeons* 2015;221(4):S155.
- [38] Kwon H-M, Moon Y-J, Jun K-W, Park Y-S, Jun I-G, Kim S-O, et al. Neutrophil-to-lymphocyte ratio is a predictor of early graft dysfunction following living donor liver transplantation. *Liver Int* 2019;39(8):1545–56.
- [39] Fu SJ, Ji F, Han M, Chen MG, Wang XP, Ju WQ, et al. Prognostic value of combined preoperative fibrinogen and neutrophil-lymphocyte ratio in patients with hepatocellular carcinoma after liver transplantation. *Oncotarget* 2017;8(3):4301–12.
- [40] Xiao GQ, Yang JY, Yan LN. Combined Hangzhou criteria with neutrophil-lymphocyte ratio is superior to other criteria in selecting liver transplantation candidates with HBV-related hepatocellular carcinoma. *HBPD INT* 2015;14(6):588–95.
- [41] Halazun KJ, Najjar M, Abdelmessih RM, Samstein B, Griesemer AD, Guarnera JV, et al. Recurrence after liver transplantation for hepatocellular carcinoma: a new MORAL to the story. *Ann Surg* 2017;265(3):557–64.
- [42] Sasaki K, Firil DJ, Hashimoto K, Fujiki M, Diago-Usó T, Quintini C, et al. Development and validation of the HALT-HCC score to predict mortality in liver transplant recipients with hepatocellular carcinoma: a retrospective cohort analysis. *Lancet Gastroenterol Hepatol* 2017;2(8):595–603.
- [43] McVey JC, Sasaki K, Firil DJ, Fujiki M, Diago-Usó T, Quintini C, et al. Prognostication of inflammatory cells in liver transplantation: Is the waitlist neutrophil-to-lymphocyte ratio really predictive of tumor biology? *Clin Transplant* 2019;33(12):e13743.
- [44] Sullivan KM, Goeschl RT, Turaga KK, Tsai S, Christians KK, White SB, et al. Neutrophil-to-lymphocyte ratio as a predictor of outcomes for patients with hepatocellular carcinoma: a Western perspective. *J Surg Oncol* 2014;109(2):95–7.
- [45] Shindoh J, Sugawara Y, Nagata R, Kaneko J, Tamura S, Aoki T, et al. Evaluation methods for pretransplant oncologic markers and their prognostic impacts in patient undergoing living donor liver transplantation for hepatocellular carcinoma. *Transpl Int* 2014;27(4):391–8.
- [46] Parisi I, Tsochatzis E, Wijewanthana H, Rodriguez-Peralvarez M, De Luca L, et al. Inflammation-based scores do not predict post-transplant recurrence of hepatocellular carcinoma in patients within Milan criteria. *Liver Transpl* 2014;20(11):1327–35.
- [47] Najjar M, Agrawal S, Emond JC, Halazun KJ. Pretreatment neutrophil-lymphocyte ratio: useful prognostic biomarker in hepatocellular carcinoma. *J Hepatocell Carcinoma* 2018;5:17–28.
- [48] Hu XG, Mao W, Park YK, Xu WG, Kim BW, Wang HJ. Blood Neutrophil-to-Lymphocyte Ratio Predicts Tumor Recurrence in Patients with Hepatocellular Carcinoma within Milan Criteria after Hepatectomy. *Yonsei Med J* 2016;57(5):1115–23.
- [49] Wang D, Bai N, Hu X, OuYang XW, Yao L, Tao Y, et al. Preoperative inflammatory markers of NLR and PLR as indicators of poor prognosis in resectable HCC. *PeerJ* 2019;7:e7132.
- [50] Fu YP, Ni XC, Yi Y, Cai XY, He HW, Wang JX, et al. A novel and validated inflammation-based score (IBS) predicts survival in patients with hepatocellular carcinoma following curative surgical resection: a STROBE-Compliant article. *Bull Sch Med Univ Md* 2016;95(7):e2784.
- [51] Hu B, Yang XR, Xu Y, Sun YF, Sun C, Guo W, et al. Systemic immune-inflammation index predicts prognosis of patients after curative resection for hepatocellular carcinoma. *Clin Cancer Res* 2014;20(23):6212–22.
- [52] Uchinaka E, Amisaki M, Morimoto M, Tokuyasu N, Sakamoto T, Honjo S, et al. Utility and limitation of preoperative neutrophil lymphocyte ratio as a prognostic factor in hepatocellular carcinoma. *Yonago Acta Med* 2018;61(4):197–203.
- [53] Hung HC, Lee JC, Cheng CH, Wu TH, Wang YC, Lee JC, et al. Impact of neutrophil to lymphocyte ratio on survival for hepatocellular carcinoma after curative resection. *J Hepatobiliary Pancreat Sci* 2017;24(10):559–69.
- [54] Mano Y, Shirabe K, Yamashita Y, Harimoto N, Tsujita E, Takeishi K, et al. Preoperative neutrophil-to-lymphocyte ratio is a predictor of survival after hepatectomy for hepatocellular carcinoma: a retrospective analysis. *Ann Surg* 2013;258(2):301–5.
- [55] Wang Q, Blank S, Fiel MI, Kadri H, Luan W, Warren L, et al. The severity of liver fibrosis influences the prognostic value of inflammation-based scores in hepatitis B-Associated hepatocellular carcinoma. *Ann Surg Oncol* 2015;22(Suppl 3):S1125–32.
- [56] Ji F, Liang Y, Fu SJ, Guo ZY, Shu M, Shen SL, et al. A novel and accurate predictor of survival for patients with hepatocellular carcinoma after surgical resection: the neutrophil to lymphocyte ratio (NLR) combined with the aspartate aminotransferase/platelet count ratio index (APRI). *BMC Cancer* 2016;16:137.
- [57] Lu SD, Wang YY, Peng NF, Peng YC, Zhong JH, Qin HG, et al. Preoperative Ratio of Neutrophils to Lymphocytes Predicts Postresection Survival in Selected Patients With Early or Intermediate Stage Hepatocellular Carcinoma. *Bull Sch Med Univ Md* 2016;95(5):e2722.
- [58] Liao R, Tang ZW, Li DW, Luo SQ, Huang P, Du CY. Preoperative neutrophil-to-lymphocyte ratio predicts recurrence of patients with single-nodule small hepatocellular carcinoma following curative resection: a retrospective report. *World J Surg Oncol* 2015;13:265.
- [59] Liao W, Zhang J, Zhu Q, Qin L, Yao W, Lei B, et al. Preoperative neutrophil-to-lymphocyte ratio as a new prognostic marker in hepatocellular carcinoma after curative resection. *Transl Oncol* 2014;7(2):248–55.
- [60] Huang GQ, Zhu GQ, Liu YL, Wang LR, Braddock M, Zheng MH, et al. Stratified neutrophil-to-lymphocyte ratio accurately predict mortality risk in hepatocellular carcinoma patients following curative liver resection. *Oncotarget* 2016;7(5):5429–39.
- [61] Huang J, Xu L, Luo Y, He F, Zhang Y, Chen M. The inflammation-based scores to predict prognosis of patients with hepatocellular carcinoma after hepatectomy. *Med Oncol* 2014;31(4):883.
- [62] Spolverato G, Maqsood H, Kim Y, Margonis G, Luo T, Ejaz A, et al. Neutrophil-lymphocyte and platelet-lymphocyte ratio in patients after resection for hepato-pancreatico-biliary malignancies. *J Surg Oncol* 2015;111(7):868–74.
- [63] Yang Hj, Guo Z, Yang YT, Jiang JH, Qi YP, Li JJ, et al. Blood neutrophil-lymphocyte ratio predicts survival after hepatectomy for hepatocellular carcinoma: A propensity score-based analysis. *World J Gastroenterol* 2016;22(21):5088–95.
- [64] Kim WJ, Lim TW, Park PJ, Choi SB, Kim WJ. Prognostic markers affecting the early recurrence of hepatocellular carcinoma with liver cirrhosis after curative resection. *Int J Biol Markers* 2019;34(2):123–31.
- [65] Okamura Y, Ashida R, Ito T, Sugiyama T, Mori K, Uesaka K. Preoperative Neutrophil to Lymphocyte Ratio and Prognostic Nutritional Index Predict Overall Survival After Hepatectomy for Hepatocellular Carcinoma. *World J Surg* 2015;39(6):1501–9.
- [66] Yang Hj, Guo Z, Yang YT, Jiang JH, Qi YP, Li JJ, et al. Blood neutrophil-lymphocyte ratio predicts survival after hepatectomy for hepatocellular carcinoma: A propensity score-based analysis. *World J Gastroentero* 2016;22(21):5088–95.
- [67] Higashi T, Hayashi H, Kaida T, Arima K, Takeyama H, Taki K, et al. Prognostic impact of visceral fat amount and branched-chain amino acids (BCAA) in hepatocellular carcinoma. *Ann Surg Oncol* 2015;22(Suppl 3):S1041–7.
- [68] Sempokuya T, Wong LL. Ten-year survival and recurrence of hepatocellular cancer. *Hepatoma Res* 2019;5.
- [69] Shiraki T, Ishizuka M, Kubota K, Kato M, Matsumoto T, Mori S, et al. An elevated neutrophil-to-lymphocyte ratio predicts a poor postoperative survival in primary hepatocellular carcinoma patients with a normal preoperative serum level of alpha-fetoprotein. *Surg Today* 2019;49(8):661–9.
- [70] Li SH, Wang QX, Yang ZY, Jiang W, Li C, Sun P, et al. Prognostic value of the neutrophil-to-lymphocyte ratio for hepatocellular carcinoma patients with portal/hepatic vein tumor thrombosis. *World J Gastroenterol* 2017;23(17):3122–32.
- [71] Arai K, Fukumoto T, Kido M, Tanaka M, Kuramitsu K, Kinoshita H, et al. Preoperative neutrophil-to-lymphocyte ratio as a predictor of survival after reductive surgery plus percutaneous isolated hepatic perfusion for hepatocellular carcinoma: a retrospective analysis. *Surg Today* 2017;47(3):385–92.
- [72] Wong L, Bozhilov K, Hernandez B, Kwee S, Chan O, Ellis L, et al. Underlying liver disease and advanced stage liver cancer are associated with elevated neutrophil-lymphocyte ratio. *Clin Mol Hepatol* 2019;25(3):305–16.
- [73] Galun D, Bogdanovic A, Kovac JD, Bulajic P, Loncar Z, Zuvela M. Preoperative neutrophil-to-lymphocyte ratio as a prognostic predictor after curative-intent surgery for hepatocellular carcinoma: experience from a developing country. *Cancer Manag Res* 2018;10:977–88.
- [74] Young AL, Adair R, Prasad KR, Toogood GJ, Lodge JPA. Hepatocellular Carcinoma Within a Noncirrhotic, Nonfibrotic, Seronegative Liver: Surgical Approaches and Outcomes. *J Am Coll Surgeons* 2012;214(2):174–83.
- [75] Okamura Y, Sugiyama T, Ito T, Yamamoto Y, Ashida R, Mori K, et al. Neutrophil to lymphocyte ratio as an indicator of the malignant behaviour of hepatocellular carcinoma. *Br J Surg* 2016;103(7):891–8.
- [76] Kabir T, Ye M, Mohd Noor NA, Woon W, Junnarkar SP, et al. Preoperative neutrophil-to-lymphocyte ratio plus platelet-to-lymphocyte ratio predicts the outcomes after curative resection for hepatocellular carcinoma. *Int J Hepatol* 2019;2019, 4239463.
- [77] Kong W, Xu H, Cheng J, Fang Z, Wang H, Zhang J, et al. The Prognostic Role of a Combined Fibrinogen and Neutrophil-to-Lymphocyte Ratio Score in Patients with Resectable Hepatocellular Carcinoma: A Retrospective Study. *Med Sci Monit* 2020;26, e918824-e.
- [78] Ren Z, He S, Fan X, He F, Sang W, Bao Y, et al. Survival prediction model for postoperative hepatocellular carcinoma patients. *Medicine* 2017;96(37), e7902-e.
- [79] Goh BK, Kam JH, Lee SY, Chan CY, Allen JC, Jeyaraj P, et al. Significance of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and prognostic nutrition index as preoperative predictors of early mortality after liver resection for huge (>10 cm) hepatocellular carcinoma. *J Surg Oncol* 2016;113(6):621–7.
- [80] Chan AW, Chan SL, Wong GL, Wong VW, Chong CC, Lai PB, et al. Prognostic nutritional index (PNI) predicts tumor recurrence of very Early/Early stage hepatocellular carcinoma after surgical resection. *Ann Surg Oncol* 2015;22(13):4138–48.
- [81] Zheng J, Seier K, Gonen M, Balachandran VP, Kingham TP, D'Angelica MI, et al. Utility of serum inflammatory markers for predicting microvascular invasion and survival for patients with hepatocellular carcinoma. *Ann Surg Oncol* 2017;24(12):3706–14.

- [82] Ni XC, Yi Y, Fu YP, He HW, Cai XY, Wang JX, et al. Prognostic value of the modified glasgow prognostic score in patients undergoing radical surgery for hepatocellular carcinoma. *Bull Sch Med Univ Md* 2015;94(36):e1486.
- [83] Kinoshita A, Onoda H, Imai N, Iwaku A, Oishi M, Fushiyama N, et al. Comparison of the prognostic value of inflammation-based prognostic scores in patients with hepatocellular carcinoma. *Br J Cancer* 2012;107(6):988–93.
- [84] Dai T, Lin G, Deng M, Zhu S, Li H, Yao J, et al. The prognostic significance of neutrophil-to-lymphocyte ratio at different time points in patients with hepatocellular carcinoma receiving liver resection. *Transl Cancer Res* 2020;9(2):441–57.
- [85] Peng W, Li C, Wen TF, Yan LN, Li B, Wang WT, et al. Neutrophil to lymphocyte ratio changes predict small hepatocellular carcinoma survival. *J Surg Res* 2014;192(2):402–8.
- [86] Cai X, Zhai J, Kaplan DE, Zhang Y, Zhou L, Chen X, et al. Background progenitor activation is associated with recurrence after hepatectomy of combined hepatocellular-cholangiocarcinoma. *Hepatology* 2012;56(5):1804–16.
- [87] Dou JP, Yu J, Yang XH, Cheng ZG, Han ZY, Liu FY, et al. Outcomes of microwave ablation for hepatocellular carcinoma adjacent to large vessels: a propensity score analysis. *Oncotarget* 2017;8(17):28758–68.
- [88] Chang JX, Zeng Z, Lu YY, Qu HJ, Xu LG, Gao X, et al. 01-2-5 - Nlr of Pro-Cryoablation Can Predict the Prognosis of Patients with Advanced Hepatocellular Carcinoma. *Ann Oncol* 2014;25:v45.
- [89] Li X, Han Z, Cheng Z, Yu J, Liu S, Yu X, et al. Preoperative neutrophil-to-lymphocyte ratio is a predictor of recurrence following thermal ablation for recurrent hepatocellular carcinoma: a retrospective analysis. *PLoS One* 2014;9(10):e110546.
- [90] Casadei Gardini A, Marisi G, Canale M, Foschi FG, et al. Radiofrequency ablation of hepatocellular carcinoma: a meta-analysis of overall survival and recurrence-free survival. *Oncotargets Ther* 2018;11:6555–67.
- [91] Tan W, Sun WK, Li X, Zhao L, Wang C, Zang AH, et al. Preablation neutrophil-to-lymphocyte ratio as an independent prognostic factor in locally advanced hepatocellular carcinoma patients following radiofrequency ablation. *J Cancer Res Ther* 2018;14(1):84–9.
- [92] Chu MO, Shen CH, Chang TS, Xu HW, Yen CW, Lu SN, et al. Pretreatment inflammation-based markers predict survival outcomes in patients with early stage hepatocellular carcinoma after radiofrequency ablation. *Sci Rep-Uk* 2018;8.
- [93] Kim CG, Lee HW, Choi HJ, Lee JI, Lee HW, Kim SU, et al. Development and validation of a prognostic model for patients with hepatocellular carcinoma undergoing radiofrequency ablation. *Cancer Med* 2019;8(11):5023–32.
- [94] Sukato DC, Tohme S, Chalhoub D, Han K, Zajko A, Amesur N, et al. The Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Patients with Unresectable Hepatocellular Carcinoma Treated with Radioembolization. *J Vasc Interv Radiol* 2015;26(6):816–24.
- [95] Chen K, Zhan MX, Hu BS, Li Y, He X, Fu SR, et al. Combination of the neutrophil to lymphocyte ratio and the platelet to lymphocyte ratio as a useful predictor for recurrence following radiofrequency ablation of hepatocellular carcinoma. *Oncol Lett* 2018;15(1):315–23.
- [96] Chen TM, Lin CC, Huang PT, Wen CF. Neutrophil-to-lymphocyte ratio associated with mortality in early hepatocellular carcinoma patients after radiofrequency ablation. *J Gastroenterol Hepatol* 2012;27(3):553–61.
- [97] Dan J, Zhang Y, Peng Z, Huang J, Gao H, Xu L, et al. Postoperative neutrophil-to-lymphocyte ratio change predicts survival of patients with small hepatocellular carcinoma undergoing radiofrequency ablation. *PLoS One* 2013;8(3):e58184.
- [98] Tajiri K, Baba H, Kawai K, Minemura M, Yasumura S, Takahara T, et al. Neutrophil-to-lymphocyte ratio predicts recurrence after radiofrequency ablation in hepatitis B virus infection. *J Gastroenterol Hepatol* 2016;31(7):1291–9.
- [99] Tajiri K, Kawai K, Minemura M, Yasumura S, Hosokawa A, Kawabe H, et al. Neutrophil/lymphocyte ratio as a prognostic indicator of hepatic arterial infusion chemotherapy with arterial cisplatin plus continuous 5-fluorouracil. *Hepatol Res* 2015;45(7):755–63.
- [100] Mizukoshi E, Yamashita T, Arai K, Terashima T, Kitahara M, Nakagawa H, et al. Myeloid-derived suppressor cells correlate with patient outcomes in hepatic arterial infusion chemotherapy for hepatocellular carcinoma. *Cancer Immunol Immunother* 2016;65(6):715–25.
- [101] Tsunematsu S, Suda G, Yamasaki K, Kimura M, Takaaki I, Umemura M, et al. Combination of neutrophil-to-lymphocyte ratio and early des-gamma-carboxyprothrombin change ratio as a useful predictor of treatment response for hepatic arterial infusion chemotherapy against advanced hepatocellular carcinoma. *Hepatol Res* 2017;47(6):533–41.
- [102] Taussig MD, Irene Koran ME, Mouli SK, Ahmad A, Geevarghese S, Baker JC, et al. Neutrophil to lymphocyte ratio predicts disease progression following intra-arterial therapy of hepatocellular carcinoma. *HPB (Oxford)* 2017;19(5):458–64.
- [103] Cruz JC, Watchmaker JM, Albin MM, Wang L, Wu G, Baker JC, et al. Neutrophil/Lymphocyte ratio predicts increased risk of immediate progressive disease following chemoembolization of hepatocellular carcinoma. *J Vasc Interv Radiol* 2019;30(12):1887–92.
- [104] McNally ME, Martinez A, Khabiri H, Guy G, Michaels AJ, Hanje J, et al. Inflammatory Markers are Associated with Outcome in Patients with Unresectable Hepatocellular Carcinoma Undergoing Transarterial Chemoembolization. *Ann Surg Oncol* 2013;20(3):923–8.
- [105] Terashima T, Yamashita T, Iida N, Yamashita T, Nakagawa H, Arai K, et al. Blood neutrophil to lymphocyte ratio as a predictor in patients with advanced hepatocellular carcinoma treated with hepatic arterial infusion chemotherapy. *Hepatol Res* 2015;45(9):949–59.
- [106] Xu X, Chen W, Zhang L, Miao R, Zhou Y, Wan Y, et al. Prognostic significance of neutrophil to lymphocyte ratio in patients with hepatocellular carcinoma after transcatheter arterial chemoembolization. *Chin Med J (Engl)* 2014;127(24):4204–9.
- [107] Jianyong L, Jinjing Z, Lunan Y, Jingqiang Z, Wentao W, Yong Z, et al. Preoperative adjuvant transarterial chemoembolization cannot improve the long term outcome of radical therapies for hepatocellular carcinoma. *Sci Rep* 2017;7:41624.
- [108] Rebonato A, Graziosi L, Maiettini D, Marino E, De Angelis V, Brunese L, et al. Inflammatory Markers as Prognostic Factors of Survival in Patients Affected by Hepatocellular Carcinoma Undergoing Transarterial Chemoembolization. *Gastroenter Res Pract* 2017.
- [109] Oh BS, Jang JW, Kwon JH, You CR, Chung KW, Kay CS, et al. Prognostic value of C-reactive protein and neutrophil-to-lymphocyte ratio in patients with hepatocellular carcinoma. *BMC Cancer* 2013;13.
- [110] Sandow TA, Arndt SE, Albar AA, DeVun DA, Kirsch DS, Gimenez JM, et al. Assessment of Response to Transcatheter Arterial Chemoembolization with Doxorubicin-eluting Microspheres: Tumor Biology and Hepatocellular Carcinoma Recurrence in a 5-year Transplant Cohort. *Radiology* 2018;286(3):1072–83.
- [111] Fan WZ, Zhang YQ, Wang Y, Yao XH, Yang JY, Li JP. Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios as Predictors of Survival and Metastasis for Recurrent Hepatocellular Carcinoma after Transarterial Chemoembolization. *PLoS One* 2015;10(3).
- [112] Zhang J, Gong F, Li L, Zhao M, Song J. Diabetes mellitus and the neutrophil to lymphocyte ratio predict overall survival in non-viral hepatocellular carcinoma treated with transarterial chemoembolization. *Oncol Lett* 2014;7(5):1704–10.
- [113] Liu C, Li L, Lu WS, Du H, Yan LN, Yang JY, et al. Neutrophil-lymphocyte Ratio Prognostic Nutritional Index Predicts the Outcomes of Patients with Unresectable Hepatocellular Carcinoma After Transarterial Chemoembolization. *Sci Rep* 2017;7(1):13873.
- [114] He C, Zhang Y, Cai Z, Lin X. The prognostic and predictive value of the Combination of the neutrophil-to-lymphocyte ratio and the platelet-to-lymphocyte ratio in patients with hepatocellular carcinoma who receive transarterial chemoembolization therapy. *Cancer Manag Res* 2019;11:1391–400.
- [115] Chon YE, Park H, Hyun HK, Ha Y, Kim MN, Kim BK, et al. Development of a New Nomogram Including Neutrophil-to-Lymphocyte Ratio to Predict Survival in Patients with Hepatocellular Carcinoma Undergoing Transarterial Chemoembolization. *Cancers (Basel)* 2019;11(4).
- [116] Zhou DS, Xu L, Luo YL, He FY, Huang JT, Zhang YJ, et al. Inflammation scores predict survival for hepatitis B virus-related hepatocellular carcinoma patients after transarterial chemoembolization. *World J Gastroentero* 2015;21(18):5582–90.
- [117] Yang ZG, Zhang JL, Lu YF, Xu QNA, Tang BZ, Wang Q, et al. Aspartate aminotransferase-lymphocyte ratio index and systemic immune-inflammation index predict overall survival in HBV-related hepatocellular carcinoma patients after transcatheter arterial chemoembolization. *Oncotarget* 2015;6(40):43090–8.
- [118] Pinato DJ, Sharma R. An inflammation-based prognostic index predicts survival advantage after transarterial chemoembolization in hepatocellular carcinoma. *Transl Res* 2012;160(2):146–52.
- [119] Wei K, Wang M, Zhang W, Mu H, Song TQ. Neutrophil-lymphocyte ratio as a predictor of outcomes for patients with hepatocellular carcinoma undergoing TAE combined with Sorafenib. *Med Oncol* 2014;31(6):969.
- [120] Zhang W, Zhao G, Wei K, Zhang QX, Ma WW, Wu Q, et al. Adjuvant sorafenib therapy in patients with resected hepatocellular carcinoma: evaluation of predictive factors. *Med Oncol* 2015;32(4).
- [121] Sun S, Wang X, Chen J. Using pre-treatment neutrophil-to-Lymphocyte ratio to predict the prognosis of young patients with hepatocellular carcinoma implemented minimally invasive treatment. *J Adolesc Young Adult Oncol* 2020;9(1):85–9.
- [122] Hong YM, Cho M, Yoon KT, Ryu JH, Yang KH, Jeon UB, et al. Neutrophil-lymphocyte ratio predicts the therapeutic benefit of neoadjuvant transarterial chemoembolization in patients with resectable hepatocellular carcinoma. *Eur J Gastroenterol Hepatol* 2019.
- [123] Liu C, Zeng G-J, Jia B-S, Yang J-Y, Jia B-S, Du H, et al. Neutrophil and platelet-to-lymphocyte ratio predicts outcomes in patients with unresectable hepatocellular carcinoma undergoing 131I-labeled-metuximab plus transarterial chemoembolization. *Oncotarget* 2018.
- [124] Long J, Zheng JS, Sun B, Lu N. Microwave ablation of hepatocellular carcinoma with portal vein tumor thrombosis after transarterial chemoembolization: a prospective study. *Hepatol Int* 2016;10(1):175–84.
- [125] Shiozawa K, Watanabe M, Ikehara T, Matsukiyo Y, Kogame M, Shinohara M, et al. [Plasma Biomarkers as Predictive Factors for Advanced Hepatocellular Carcinoma with Sorafenib]. *Gan To Kagaku Ryoho* 2016;43(7):863–7.
- [126] Lué A, Bustamante FJ, Iñarrairaegui M, Arenas JL, Serrano MT, Testillano M, et al. P979 neutrophil-to-lymphocyte ratio is a predictor of one-year survival in patients with hepatocellular carcinoma receiving sorafenib. *J Hepatol* 2014;60(1):S401.
- [127] Afshar M, Clarke H, Jackson-Wilding A, Ahmed A, Ma YT, Punia P. P0352 : Neutrophil lymphocyte ratio (NLR) at diagnosis is a predictor for survival in patients receiving sorafenib for advanced hepatocellular carcinoma (HCC): a large UK cohort. *J Hepatol* 2015;62:S442.

- [128] da Fonseca LG, Barroso-Sousa R, Bento Ada S, Blanco BP, Valente GL, Pfiffer TE, et al. Pre-treatment neutrophil-to-lymphocyte ratio affects survival in patients with advanced hepatocellular carcinoma treated with sorafenib. *Med Oncol* 2014;31(11):264.
- [129] Zheng YB, Zhao W, Liu B, Lu LG, He X, Huang JW, et al. The blood neutrophil-to-lymphocyte ratio predicts survival in patients with advanced hepatocellular carcinoma receiving sorafenib. *Asian Pac J Cancer Prev* 2013;14(9):5527–31.
- [130] Personeni N, Giordano L, Abbadessa G, Porta C, Borbath I, Daniele B, et al. Prognostic value of the neutrophil-to-lymphocyte ratio in the ARQ 197-215 second-line study for advanced hepatocellular carcinoma. *Oncotarget* 2017;8(9):14408–15.
- [131] Lue A, Serrano MT, Bustamante FJ, Inarraegui M, Arenas JJ, Testillano M, et al. Neutrophil-to-lymphocyte ratio predicts survival in European patients with hepatocellular carcinoma administered sorafenib. *Oncotarget* 2017;8(61):103077–86.
- [132] Howell J, Pinato DJ, Ramaswami R, Arizumi T, Ferrari C, Gibbin A, et al. Integration of the cancer-related inflammatory response as a stratifying biomarker of survival in hepatocellular carcinoma treated with sorafenib. *Oncotarget* 2017;8(22):36161–70.
- [133] Bruix J, Cheng AL, Meinhardt G, Nakajima K, De Sanctis Y, et al. Prognostic factors and predictors of sorafenib benefit in patients with hepatocellular carcinoma: Analysis of two phase III studies. *J Hepatol* 2017;67(5):999–1008.
- [134] Casadei Gardini A, Scarpi E, Faloppi L, Scartozzi M, et al. Immune inflammation indicators and implication for immune modulation strategies in advanced hepatocellular carcinoma patients receiving sorafenib. *Oncotarget* 2016;7(41):67142–9.
- [135] Afshar M, Fletcher P, Bardoli AD, Ma YT, Punia P. Non-secretion of AFP and neutrophil lymphocyte ratio as predictors for survival in hepatocellular carcinoma patients treated with sorafenib: a large UK cohort. *Oncotarget* 2018;9(24):16988–95.
- [136] Zheng YB, Zhao W, Liu B, Lu LG, He X, Huang JW, et al. The Blood Neutrophil-to-lymphocyte Ratio Predicts Survival in Patients with Advanced Hepatocellular Carcinoma Receiving Sorafenib. *Asian Pac J Cancer Prev* 2013;14(9):5527–31.
- [137] Liu L, Gong Y, Zhang Q, Cai P, Feng L. Prognostic Roles of Blood Inflammatory Markers in Hepatocellular Carcinoma Patients Taking Sorafenib. A Systematic Review and Meta-Analysis. *Front Oncol* 2019;9:1557.
- [138] Diaz-Beveridge R, Bruxola G, Lorente D, Caballero J, Rodrigo E, Segura A, et al. An internally validated new clinical and inflammation-based prognostic score for patients with advanced hepatocellular carcinoma treated with sorafenib. *Clin Transl Oncol* 2018;20(3):322–9.
- [139] He G, Zhang H, Zhou J, Wang B, Chen Y, Kong Y, et al. Peritumoural neutrophils negatively regulate adaptive immunity via the PD-L1/PD-1 signalling pathway in hepatocellular carcinoma. *J Exp Clin Cancer Res* 2015;34:141.
- [140] Tada T, Kumada T, Hiraoka A, Michitaka K, Atsukawa M, Hirooka M, et al. Neutrophil-to-lymphocyte ratio is associated with survival in patients with unresectable hepatocellular carcinoma treated with lenvatinib. *Liver Int* 2020;40(4):968–76.
- [141] Sprinzl MF, Kirstein MM, Koch S, Seib ML, Weinmann-Menke J, Lang H, et al. Improved Prediction of Survival by a Risk Factor-Integrating Inflammatory Score in Sorafenib-Treated Hepatocellular Carcinoma. *Liver Cancer* 2019;8(5):387–402.
- [142] Koga Y, Beppu T, Miyata T, Kitano Y, Tsuji A, Nakagawa S, et al. Predicting poorly differentiated hepatocellular carcinoma that meets the milan criteria. *Anticancer Res* 2018;38(7):4093–9.
- [143] Gao F, Li X, Geng M, Ye X, Liu H, Liu Y, et al. Pretreatment neutrophil-lymphocyte ratio: an independent predictor of survival in patients with hepatocellular carcinoma. *Bull Sch Med Univ Md* 2015;94(11):e639.
- [144] Pinato DJ, Stebbing J, Ishizuka M, Khan SA, Wasan HS, North BV, et al. A novel and validated prognostic index in hepatocellular carcinoma: the inflammation based index (IBI). *J Hepatol* 2012;57(5):1013–20.
- [145] Hu Z, Chen H, Chen S, Huang Z, Qin S, Zhong J, et al. The value of neutrophil to lymphocyte ratio and gamma-glutamyl transpeptidase to platelet ratio in patients with hepatocellular carcinoma. *Bull Sch Med Univ Md* 2019;98(9):e14749.
- [146] Shen J, Tang L, Zhang X, Peng W, Wen T, Li C, et al. A Novel Index in Hepatocellular Carcinoma Patients After Curative Hepatectomy: Albumin to Gamma-Glutamyltransferase Ratio (AGR). *Front Oncol* 2019;9(817).
- [147] Zeng F, Chen B, Zeng J, Wang Z, Xiao L, Deng G. Preoperative neutrophil-lymphocyte ratio predicts the risk of microvascular invasion in hepatocellular carcinoma: a meta-analysis. *Int J Biol Markers* 2019;34(3):213–20.
- [148] Deng G, Yao L, Zeng F, Xiao L, Wang Z. Nomogram for preoperative prediction of microvascular invasion risk in hepatocellular carcinoma. *Cancer Manag Res* 2019;11:9037–45.
- [149] Hatanaka T, Naganuma A, Uehara R, Saito N, Nakano S, Kato M, et al. Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratio predict overall survival in hepatocellular carcinoma patients with non-B, non-C cirrhosis. *Clin Nutr* 2018;37:S204.
- [150] Li X, Chen ZH, Ma XK, Chen J, Wu DH, Lin Q, et al. Neutrophil-to-lymphocyte ratio acts as a prognostic factor for patients with advanced hepatocellular carcinoma. *Tumour Biol* 2014;35(11):11057–63.
- [151] Lai Q, Castro Santa E, Rico Juri JM, Pinheiro RS, Lerut J. Neutrophil and platelet-to-lymphocyte ratio as new predictors of dropout and recurrence after liver transplantation for hepatocellular cancer. *Transpl Int* 2014;27(1):32–41.
- [152] Ding Y, Liu K, Xu Y, Zhao Q, Lou S, Xiang X, et al. Combination of inflammatory score/liver function and AFP improves the diagnostic accuracy of HBV-related hepatocellular carcinoma. *Cancer Med* 2020.
- [153] Du Z, Dong J, Bi J, Bai R, Zhang J, Wu Z, et al. Predictive value of the preoperative neutrophil-to-lymphocyte ratio for the development of hepatocellular carcinoma in HBV-associated cirrhotic patients after splenectomy. *PLoS One* 2018;13(4):e0195336.
- [154] Kusumanto YH, Dam WA, Hospers GA, Meijer C, Mulder NH. Platelets and granulocytes, in particular the neutrophils, form important compartments for circulating vascular endothelial growth factor. *Angiogenesis* 2003;6(4):283–7.
- [155] Tanigawa N, Amaya H, Matsumura M, Shimomatsu T. Correlation between expression of vascular endothelial growth factor and tumor vascularity, and patient outcome in human gastric carcinoma. *J Clin Oncol* 1997;15(2):826–32.
- [156] Agopian VG, Harlander-Locke M, Zarrinpar A, Kaldas FM, Farmer DG, et al. A novel prognostic nomogram accurately predicts hepatocellular carcinoma recurrence after liver transplantation: analysis of 865 consecutive liver transplant recipients. *J Am Coll Surg* 2015;220(4):416–27.
- [157] Bodzin AS, Lunsford KE, Markovic D, Harlander-Locke MP, Busuttil RW, et al. Predicting mortality in patients developing recurrent hepatocellular carcinoma after liver transplantation: impact of treatment modality and recurrence characteristics. *Ann Surg* 2017;266(1):118–25.
- [158] Yoshizumi T, Ikegami T, Toshima T, Harimoto N, Uchiyama H, Soejima Y, et al. Two-step selection criteria for living donor liver transplantation in patients with hepatocellular carcinoma. *Transpl P* 2013;45(9):3310–3.
- [159] Yoshizumi T, Harimoto N, Itoh S, Okabe H, Kimura K, Uchiyama H, et al. Living donor liver transplantation for hepatocellular carcinoma within milan criteria in the present era. *Anticancer Res* 2016;36(1):439–45.
- [160] Yoshizumi T, Ikegami T, Yoshiya S, Motomura T, Mano Y, Muto J, et al. Impact of tumor size, number of tumors and neutrophil-to-lymphocyte ratio in liver transplantation for recurrent hepatocellular carcinoma. *Hepatol Res* 2013;43(7):709–16.
- [161] Liese J, Peveling-Oberhag J, Doering C, Schnitzbauer AA, Herrmann E, Zangos S, et al. A possible role of microRNAs as predictive markers for the recurrence of hepatocellular carcinoma after liver transplantation. *Transpl Int* 2016;29(3):369–80.
- [162] Yamamura K, Sugimoto H, Kanda M, Yamada S, Nomoto S, Nakayama G, et al. Comparison of inflammation-based prognostic scores as predictors of tumor recurrence in patients with hepatocellular carcinoma after curative resection. *J Hepatobiliary Pancreat Sci* 2014;21(9):682–8.
- [163] Shimoda M, Tago K, Shiraki T, Mori S, Kato M, Aoki T, et al. Risk factors for early recurrence of single lesion hepatocellular carcinoma after curative resection. *World J Surg* 2016;40(10):2466–71.
- [164] Fu SJ, Shen SL, Li SQ, Hua YP, Hu WJ, Liang LJ, et al. Prognostic value of preoperative peripheral neutrophil-to-lymphocyte ratio in patients with HBV-associated hepatocellular carcinoma after radical hepatectomy. *Med Oncol* 2013;30(4):721.
- [165] Ruan DY, Lin ZX, Li Y, Jiang N, Li X, Wu DH, et al. Poor oncologic outcomes of hepatocellular carcinoma patients with intra-abdominal infection after hepatectomy. *World J Gastroenterol* 2015;21(18):5598–606.
- [166] Li C, Wen TF, Yan LN, Li B, Wang WT, Yang JY, et al. Postoperative neutrophil-to-lymphocyte ratio plus platelet-to-lymphocyte ratio predicts the outcomes of hepatocellular carcinoma. *J Surg Res* 2015;198(1):73–9.
- [167] Elalfy H, Besheer T, El-Maksoud MA, Farid K, Elegezy M, et al. Monocyte/granulocyte to lymphocyte ratio and the MELD score as predictors for early recurrence of hepatocellular carcinoma after trans-arterial chemoembolization. *Br J Biomed Sci* 2018;75(4):187–91.
- [168] Rahman EMA, El-Senosy FM, Alsebaey A, Hussien SA. Neutrophil to lymphocyte ratio as predictive marker of recurrence of HCC after radiofrequency ablation. *Am J Clin Med Res* 2019;7(1):1–8.
- [169] DiNorcia J, Florman Ss, Haydel B, Tabrizian P, Ruiz Rm, Klintmalm Gb, et al. Pathologic Response to Pretransplant Locoregional Therapy is Predictive of Patient Outcome After Liver Transplantation for Hepatocellular Carcinoma: Analysis From the US Multicenter HCC Transplant Consortium. *Ann Surg* 2020;271(4):616–24.
- [170] Sabry M, El-Nakeeb S, Moaty ASA, Abdelsattar HA. Neutrophil, lymphocyte, and platelet-containing ratios as predictors for hepatocellular carcinoma recurrence in Egyptian patients. *Egypt Liver J* 2018;8(3):98–104.
- [171] Chung HA, Kim J-H, Hwang Y, Choi HS, Ko SY, Choe WH, et al. Noninvasive fibrosis marker can predict recurrence of hepatocellular carcinoma after radiofrequency ablation. *Saudi J Gastroenterol* 2016;22(1):57–63.
- [172] Casadei Gardini A, Foschi FG, Conti F, Petracci E, et al. Immune inflammation indicators and ALBI score to predict liver cancer in HCV-patients treated with direct-acting antivirals. *Dig Liver Dis* 2019;51(5):681–8.
- [173] Halazun KJ, Marangoni G, Hakeem A, Fraser SM, Farid SG, Ahmad N. Elevated preoperative recipient neutrophil-lymphocyte ratio is associated with delayed graft function following kidney transplantation. *Transplant Proc* 2013;45(9):3254–7.
- [174] Ahmed N, Kim Y, Gandhi H, Saeed O, Patel S, Murthy S, et al. Pre-transplant neutrophil to lymphocyte ratio predicts 30-day readmission for patients undergoing cardiac transplantation. *J Heart Lung Transplant* 2018;37(4):S415–6.
- [175] Ahmed N, Gandhi H, Rahgozar K, Guo S, Sun E, Saeed O, et al. Elevated Pre-Transplant Neutrophil to Lymphocyte Ratio is Associated with Increased Vasoplegia Syndrome in Cardiac Transplantation. *J Heart Lung Transplant* 2019;38(4, Supplement):S211.
- [176] Narango M, Agrawal A, Goyal A, Rangaswami J. Neutrophil-to-Lymphocyte ratio and platelet-to-Lymphocyte ratio predict acute cellular rejection in the kidney allograft. *Ann Transplant* 2018;23:467–74.

- [177] Chae MS, Kim J-W, Chung HS, Park CS, Lee J, Choi JH, et al. The impact of serum cytokines in the development of early allograft dysfunction in living donor liver transplantation. *Medicine* 2018;97(16).
- [178] Lin BY, Zhou L, Geng L, Zheng ZY, Jia JJ, Zhang J, et al. High neutrophil-lymphocyte ratio indicates poor prognosis for acute-on-chronic liver failure after liver transplantation. *World J Gastroenterol* 2015;21(11):3317–24.
- [179] Hayashi H, Takamura H, Ohbatake Y, Nakanuma S, Makino I, Nakagawara H, et al. Postoperative neutrophil-to-lymphocyte ratio of living-donor liver transplant: association with graft size. *Asian J Surg* 2016;39(2):103–8.
- [180] Sun XD, Shi XJ, Chen YG, Wang CL, Ma Q, Lv GY. Elevated preoperative neutrophil-lymphocyte ratio is associated with poor prognosis in hepatocellular carcinoma patients treated with liver transplantation: a meta-analysis. *Gastroenterol Res Pract* 2016;2016:4743808.
- [181] Xu ZG, Ye CJ, Liu LX, Wu G, Zhao ZX, Wang YZ, et al. The pretransplant neutrophil-lymphocyte ratio as a new prognostic predictor after liver transplantation for hepatocellular cancer: a systematic review and meta-analysis. *Biomark Med* 2018;12(2):189–99.
- [182] Xiao GQ, Liu C, Liu DL, Yang JY, Yan LN. Neutrophil-lymphocyte ratio predicts the prognosis of patients with hepatocellular carcinoma after liver transplantation. *World J Gastroentero* 2013;19(45):8398–407.
- [183] Cescon M, Bertuzzo VR, Ercolani G, Ravaioli M, Odaldi F, Pinna AD. Liver transplantation for hepatocellular carcinoma: role of inflammatory and immunological state on recurrence and prognosis. *World J Gastroenterol* 2013;19(48):9174–82.
- [184] Min GT, Li YM, Yao N, Wang J, Wang HP, Chen W. The pretreatment neutrophil-lymphocyte ratio may predict prognosis of patients with liver cancer: a systematic review and meta-analysis. *Clin Transplant* 2018;32(1).
- [185] Aino H, Sumie S, Niizeki T, Kuromatsu R, Tajiri N, Nakano M, et al. Clinical characteristics and prognostic factors for advanced hepatocellular carcinoma with extrahepatic metastasis. *Mol Clin Oncol* 2014;2(3):393–8.
- [186] Quirós-Roldán E, Properzi M, Amadası S, Raffetti E, Ferraresi A, et al. Prognostic role of inflammatory biomarkers in HIV-infected patients with a first diagnosis of hepatocellular carcinoma: a single-center study. *J Med Virol* 2019;91(2):241–8.
- [187] Ohtani H. Focus on TILs: prognostic significance of tumor infiltrating lymphocytes in human colorectal cancer. *Cancer Immun* 2007;7:4.
- [188] Ding PR, An X, Zhang RX, Fang YJ, Li LR, Chen G, et al. Elevated preoperative neutrophil to lymphocyte ratio predicts risk of recurrence following curative resection for stage II A colon cancer. *Int J Colorectal Dis* 2010;25(12):1427–33.
- [189] Kuang D-M, Zhao Q, Wu Y, Peng C, Wang J, Xu Z, et al. Peritumoral neutrophils link inflammatory response to disease progression by fostering angiogenesis in hepatocellular carcinoma. *J Hepatol* 2011;54(5):948–55.
- [190] Waugh DJJ, Wilson C. The Interleukin-8 pathway in Cancer. *Clin Cancer Res* 2008;14(21):6735–41.
- [191] Motomura T, Shirabe K, Mano Y, Muto J, Toshima T, Umemoto Y, et al. Neutrophil-lymphocyte ratio reflects hepatocellular carcinoma recurrence after liver transplantation via inflammatory microenvironment. *J Hepatol* 2013;58(1):58–64.
- [192] Bakshi SS. Neutrophil-to-Lymphocyte ratio in patients with cirrhosis. *Clin Gastroenterol Hepatol* 2019;17(4):799.
- [193] Alkhouri N, Morris-Stiff G, Campbell C, Lopez R, Tamimi TA, Yerian L, et al. Neutrophil to lymphocyte ratio: a new marker for predicting steatohepatitis and fibrosis in patients with nonalcoholic fatty liver disease. *Liver Int* 2012;32(2):297–302.
- [194] Liu S, Zheng H, Zhu X, Mao F, Zhang S, Shi H, et al. Neutrophil-to-lymphocyte ratio is associated with diabetic peripheral neuropathy in type 2 diabetes patients. *Diabetes Res Clin Pract* 2017;130:90–7.
- [195] Sato Y, Gonda K, Harada M, Tanisaka Y, Arai S, Mashimo Y, et al. Increased neutrophil-to-lymphocyte ratio is a novel marker for nutrition, inflammation and chemotherapy outcome in patients with locally advanced and metastatic esophageal squamous cell carcinoma. *Biomed Rep* 2017;7(1):79–84.
- [196] Kuo MT, Hu TH, Lu SN, Hung CH, Wang JH, Chen CH, et al. Neutrophil-to-lymphocyte ratio as a predictor of response to peginterferon plus N. ribavirin therapy for chronic hepatitis C. *Dis Markers* 2014;2014:462958.
- [197] Wróblewska A, Lorenc B, Cheba M, Bielawski KP, Sikorska K. Neutrocyte-to-lymphocyte ratio predicts the presence of a replicative hepatitis C virus strand after therapy with direct-acting antivirals. *Clin Exp Med* 2019;19(3):401–6.
- [198] Le P-H, Liang K-H, Chang M-L, Hsu C-W, Chen Y-C, Lin C-L, et al. Clinical Predictors for Neutrophil-to-Lymphocyte Ratio ChangM.-L.s in Patients with Chronic Hepatitis B Receiving Peginterferon Treatment. In Vivo (Brooklyn) 2017;31(4):723–9.