



Contents lists available at ScienceDirect

## Annals of Hepatology

journal homepage: [www.elsevier.es/annalsofhepatology](http://www.elsevier.es/annalsofhepatology)

Original article

## Efficacy and safety of glecaprevir/pibrentasvir in treatment-naïve adults with chronic hepatitis C virus genotypes 1–6 in Brazil



Mario Peribañez-Gonzalez<sup>a,\*</sup>, Hugo Cheinquer<sup>b</sup>, Lino Rodrigues<sup>c</sup>, Maria Patelli Lima<sup>d</sup>, Mário Reis Álvares-da-Silva<sup>e</sup>, José Madruga<sup>f</sup>, Edison Roberto Parise<sup>g</sup>, Mário Guimarães Pessoa<sup>h</sup>, Juvencio Furtado<sup>i</sup>, Marcia Villanova<sup>j</sup>, Adalgisa Ferreira<sup>k</sup>, Felipe Mazzoleni<sup>l</sup>, Ecio Nascimento<sup>m</sup>, Giovanni Faria Silva<sup>n</sup>, Linda Fredrick<sup>c</sup>, Preethi Krishnan<sup>c</sup>, Margaret Burroughs<sup>c</sup>, Tania Reuter<sup>o</sup>

<sup>a</sup> Hospital Dia, Av Dr Arnaldo, 165, Segundo andar, São Paulo, Brazil<sup>b</sup> Ramiro Barcelos 2350, CPC Sala, 21216, Porto Alegre, Brazil<sup>c</sup> AbbVie Inc., North Chicago, IL, USA<sup>d</sup> Instituto de Infectologia Campinas, Rua Dr. Quirino, 524, Sala 72, Centro, Campinas, Brazil<sup>e</sup> Hospital de Clínicas de Porto Alegre, GI/Liver Division, School of Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil<sup>f</sup> Rua Santa Cruz, 81, Vila Mariana, São Paulo, Brazil<sup>g</sup> Rua Diogo de Faria, 816 - Vila Clementino, São Paulo, Brazil<sup>h</sup> Division of Gastroenterology and Hepatology University of São Paulo School of Medicine, Avenida Enéas de Carvalho Aguiar, 255 - Bloco B - 4<sup>o</sup> andar, São Paulo, Brazil<sup>i</sup> Rua Cônego Xavier, 276 - Amb de Infectologia, São Paulo, Brazil<sup>j</sup> Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo, Campus Universitário s/n, Monte Alegre, Bloco G Subsolo 2, Unidade de Pesquisa Clínica, Ribeirão Preto, Brazil<sup>k</sup> Centro de Pesquisa Clínica Hospital Universitário da Universidade Federal do Maranhão, Rua Almirante Tamandaré, 01, Centro, São Luís, Brazil<sup>l</sup> Hospital Ernesto Dornelles, Av. Ipiranga 1801, 7<sup>o</sup> Andar, CpdA-Pesquisa, Porto Alegre, Brazil<sup>m</sup> Av. Mandacaru, 1590, Maringá, Brazil<sup>n</sup> Unesp Campus de Botucatu, Rua Prof Dr Armanda Alves, s/n<sup>o</sup> Bairro: Botucatu, Brazil<sup>o</sup> Marechal Campos Av. 1355, Outclinic number 5, Vitoria, Brazil

## ARTICLE INFO

## Article history:

Received 30 June 2020

Accepted 4 September 2020

Available online 17 September 2020

## Keywords:

Antiviral agents

Brazil

Hepatitis C

Liver cirrhosis

Glecaprevir and pibrentasvir

## ABSTRACT

**Introduction and objectives:** Glecaprevir/pibrentasvir is a highly effective and well tolerated treatment for hepatitis C infection. Brazilian patients were not included in the original development studies for glecaprevir/pibrentasvir. This study aimed to assess safety and efficacy of glecaprevir/pibrentasvir in treatment-naïve Brazilian adults without cirrhosis or with compensated cirrhosis.

**Patients and methods:** EXPEDITION-3 was a Phase 3, open-label, multicenter study in treatment-naïve Brazilian adults with hepatitis C infection genotype 1–6. Patients without cirrhosis (F2 or F3) or with compensated cirrhosis (F4) received 8 or 12 weeks of glecaprevir/pibrentasvir, respectively. The primary efficacy endpoint was the rate of sustained virologic response at post-treatment Week 12. Secondary endpoints were on-treatment virologic failure and relapse rates. Baseline polymorphisms were assessed in NS3 and NS5A. Adverse events and laboratory abnormalities were monitored.

**Results:** 100 patients were enrolled, 75 received 8 weeks of treatment and 25 received 12 weeks; all patients completed treatment. Overall sustained virologic response at post-treatment Week 12 rate was high (98.0%; 98/100; 95% confidence interval: 93.0–99.4) and remained high regardless of baseline viral or host factors, including demographics, hepatitis C virus RNA levels, polymorphisms in NS3 and/or NS5A,

\* Corresponding author:

E-mail addresses: [mpgonzalez@terra.com.br](mailto:mpgonzalez@terra.com.br) (M. Peribañez-Gonzalez), [drhugo@terra.com.br](mailto:drhugo@terra.com.br) (H. Cheinquer), [lino.rodrigues@abbvie.com](mailto:lino.rodrigues@abbvie.com) (L. Rodrigues), [mariapatelli@uol.com.br](mailto:mariapatelli@uol.com.br) (M.P. Lima), [marioreis@live.com](mailto:marioreis@live.com) (M.R. Álvares-da-Silva), [valdezmr@uol.com.br](mailto:valdezmr@uol.com.br) (J. Madruga), [parise@sbhepatologia.org.br](mailto:parise@sbhepatologia.org.br) (E.R. Parise), [mgpessoa@uol.com.br](mailto:mgpessoa@uol.com.br) (M.G. Pessoa), [juvencio.furtado@terra.com.br](mailto:juvencio.furtado@terra.com.br) (J. Furtado), [marciavillanova@gmail.com](mailto:marciavillanova@gmail.com) (M. Villanova), [adalgisaf@terra.com.br](mailto:adalgisaf@terra.com.br) (A. Ferreira), [fimazzoleni@hotmail.com](mailto:fimazzoleni@hotmail.com) (F. Mazzoleni), [eanascimento@uem.br](mailto:eanascimento@uem.br) (E. Nascimento), [Giovanni.f.silva@unesp.br](mailto:Giovanni.f.silva@unesp.br) (G.F. Silva), [linda.m.fredrick@abbvie.com](mailto:linda.m.fredrick@abbvie.com) (L. Fredrick), [Preethi.krishnan@abbvie.com](mailto:Preethi.krishnan@abbvie.com) (P. Krishnan), [Margaret.burroughs@abbvie.com](mailto:Margaret.burroughs@abbvie.com) (M. Burroughs), [tania.reuter@gmail.com](mailto:tania.reuter@gmail.com) (T. Reuter).

<https://doi.org/10.1016/j.aohep.2020.09.002>1665-2681/© 2020 Published by Elsevier España, S.L.U. on behalf of Fundación Clínica Médica Sur, A.C. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

genotype, and relevant comorbidities. 55% of patients reported  $\geq 1$  adverse event, the most common being headache (18.0%). Four patients reported serious adverse events; none were considered drug related or led to study drug discontinuation. No hepatic decompensations were observed.

**Conclusions:** Glecaprevir/pibrentasvir was effective and well tolerated in treatment-naïve Brazilian patients with hepatitis C infection without cirrhosis and with compensated cirrhosis.

**Trial Registration:** [ClinicalTrials.gov](https://clinicaltrials.gov) NCT03219216.

© 2020 Published by Elsevier España, S.L.U. on behalf of Fundación Clínica Médica Sur, A.C. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

In 2015, the World Health Organization (WHO) estimated that 71 million people were living with hepatitis C virus (HCV) [1]. In Brazil, it is estimated that 1.4–1.7 million people are chronically infected with HCV [2]. With such a high prevalence, at the 2017 World Hepatitis Summit in Sao Paulo, the viral hepatitis community requested that viral hepatitis be given the same priority as that given to human immunodeficiency virus, tuberculosis, and malaria, in order to end the epidemic in Brazil [3].

Brazil's HCV treatment program is growing, with more than 41,000 people having received treatment in 2016, compared with 7500 in 2015 [4]. Furthermore, treatment protocols were also updated in August 2017 to expand treatment eligibility to include patients with F2-stage fibrosis regardless of time of diagnosis, increasing the number of patients eligible for treatment in Brazil to 80,000 [4]. Achieving the WHO's 2030 elimination targets is considered feasible in Brazil; however, it would require substantial upscaling in prevention activities, screening, and treatment of HCV [4,5]. In particular, increasing the proportion of diagnosed patients and those eligible for treatment is necessary in order to reduce HCV burden in Brazil [6].


Glecaprevir/pibrentasvir, an all oral, pangenotypic, interferon-free, ribavirin-free direct-acting antiviral regimen [7], was granted marketing authorization on the 16th of April 2018 in Brazil [8]. The approved treatment durations for treatment-naïve (TN) patients in Brazil are 8 weeks in those without cirrhosis, and 12 weeks in those with compensated cirrhosis (CC) [9]. Brazilian patients were not included in the original development studies for glecaprevir/pibrentasvir; therefore, this study was conducted to evaluate the efficacy and safety of glecaprevir/pibrentasvir in this specific population. After the initiation of this study, the approved treatment duration was reduced to 8 weeks in all TN patients with CC in the European Union (EU) and United States [7,10]. This change was based on the results of the EXPEDITION-8 study, which was performed to investigate 8 weeks of glecaprevir/pibrentasvir in TN patients with CC [11]. At the time of study design for EXPEDITION-3, the data from EXPEDITION-8 were not yet available.

## 2. Material and methods

### 2.1. Study design and patients

EXPEDITION-3 was a Phase 3, open-label, multicenter study in TN Brazilian adults with chronic HCV GT1–6, without cirrhosis or with CC (Fig. 1). A minimum of approximately 35 GT1 and 25 GT3 patients were planned for inclusion across 14 sites (approximately 80 F2–3 patients and approximately 20 F4 patients).

Patients without cirrhosis (F2–F3) were identified by (1) a liver biopsy within 24 months before or during screening that demonstrated the absence of cirrhosis; (2) a FibroScan® (Echosens,



**Fig. 1.** Study schematics.<sup>†</sup>

<sup>†</sup>The study enrollment was monitored to meet the following enrollment criteria: 1) a minimum of approximately 35 GT1 and 25 GT3 patients and 2) approximately 80 F2–3 and a maximum of approximately 20 F4 patients. G/P, glecaprevir/pibrentasvir; PTW12, post-treatment Week 12; QD, once daily.

Waltham, MA) score of 8.8 to  $<12.5$  kPa within 6 months prior to screening or during screening; or (3) a screening FibroTest (Bio-Predictive, Paris, France) score of 0.49–0.72 (inclusive). Patients without cirrhosis were enrolled into Arm A and received 8 weeks of treatment. Patients with CC (F4) were identified by (1) a previous histologic diagnosis of cirrhosis on liver biopsy; (2) a FibroScan® score of  $\geq 12.5$  kPa within 6 months prior to screening or during screening; or (3) a screening FibroTest score of  $\geq 0.73$ . Patients with CC were enrolled into Arm B and received 12 weeks of treatment. For enrollment purposes, the result of the liver biopsy superseded those of the FibroScan® and FibroTest, and the results of the FibroScan® superseded that of the FibroTest. Patients with cirrhosis were required to be compensated (Child–Pugh Score of  $\leq 6$ ) at screening. All patients were monitored for 12 weeks after their last dose (post-treatment Week 12 [PTW12]).

#### 2.1.1. Key inclusion criteria

Enrolled patients met the following criteria: male or female adults (aged  $\geq 18$  years at the time of screening) with chronic HCV GT1–6 infection (mixed and indeterminate GTs were acceptable); positive plasma HCV antibody and HCV RNA viral load  $\geq 1000$  IU/mL at screening; documented as without cirrhosis (with a METAVIR equivalent fibrosis stage of F2–F3), or with CC (with a METAVIR equivalent fibrosis stage of F4 and a Child–Pugh score of  $\leq 6$ ). Patients with CC also needed to demonstrate absence of hepatocellular carcinoma (HCC) by a negative ultrasound, computed tomography scan or magnetic resonance imaging within 3 months prior to screening or a negative ultrasound at screening.

#### 2.1.2. Key exclusion criteria

Patients were excluded from the study if they met any of the following criteria: current hepatitis B virus (HBV) infection (defined as a positive HBV surface antigen or HBV DNA higher than the lower limit of quantification [LLOQ] in patients with isolated positive anti-HBV core antibody); current or past clinical evidence of liver decompensation, including ascites on physical examination, hepatic encephalopathy, or variceal bleeding; history of solid organ transplantation (unless the transplanted organ had been removed, or was non-functional and the patient was clinically stable off immunosuppressive medication for  $\geq 6$  months prior to screen-

ing); alanine aminotransferase (ALT) or aspartate aminotransferase (AST) >10 times the upper limit of normal (ULN); total bilirubin >3.0 mg/dL; albumin < lower limit of normal (patients without cirrhosis) or <2.8 mg/dL (patients with CC); platelets <90,000  $10^3/\mu\text{L}$  (patients without cirrhosis) or <60,000  $10^3/\mu\text{L}$  (patients with CC); and receipt of any investigational or commercially available anti-HCV agents.

### 2.1.3. Ethics

The study was conducted in accordance with the protocol, International Council for Harmonisation guidelines, applicable regulations and guidelines governing clinical study conduct, and the ethical principles that have their origin in the Declaration of Helsinki. The Independent Ethics Committee/Institutional Review Board reviewed the study prior to initiation and approved the protocol, informed consent and patient information. Informed consent was provided by patients prior to any study-related screening procedures.

### 2.1.4. Procedures

Patients were treated with 300 mg/120 mg glecaprevir/pibrentasvir once daily for 8 weeks in Arm A and for 12 weeks in Arm B. Plasma samples were collected at screening and GTs were assessed using Versant<sup>®</sup> HCV Genotype Inno LiPA Assay, Version 2.0 or higher (LiPA; Siemens Healthcare Diagnostics, Tarrytown, NY), or by Sanger sequencing of the nonstructural protein (NS) 5B gene region by the central laboratory if the LiPA assay was unable to determine the sample GT. Plasma HCV RNA levels were determined by COBAS<sup>®</sup> AmpliPrep/COBAS<sup>®</sup> TaqMan HCV Quantitative Test v2.0 (Roche Molecular Systems, Pleasanton, CA). Study visits took place at screening, Day 1, Week 4, Week 8 (Arm B only), end of treatment (EOT), post-treatment Week 4 (PTW4), and PTW12.

## 2.2. Efficacy endpoints

The primary efficacy endpoint was the percentage of patients who achieved sustained virologic response at PTW12 (SVR12), defined as HCV RNA < LLOQ 12 weeks after the last actual dose of study drug, across all GTs. The secondary efficacy endpoints were the percentage of patients with HCV on-treatment virologic failure (OTVF), defined as (1) a confirmed increase of >1  $\log_{10}$  IU/mL above nadir during treatment or HCV RNA  $\geq 100$  IU/mL after HCV RNA < LLOQ during treatment, or (2) HCV RNA  $\geq$  LLOQ at EOT with  $\geq 6$  weeks of treatment; and the percentage of patients with HCV virologic relapse, defined as HCV RNA  $\geq$  LLOQ between EOT and 12 weeks after the last dose of glecaprevir/pibrentasvir among patients who completed treatment as planned with HCV RNA < LLOQ at EOT, excluding those who were shown to be reinfected. Completion of treatment for efficacy analyses was defined as any patient with treatment duration  $\geq 52$  days and  $\geq 77$  days for 8 and 12 weeks, respectively.

### 2.2.1. Resistance

Regions encoding full-length NS3/4A or NS5A were sequenced by next generation sequencing (NGS) from available baseline (BL) samples from all patients. HCV GTs and subtypes were subsequently confirmed via phylogenetic analysis of available NS3/4A and/or NS5A sequences. For resistance analyses, BL polymorphisms (defined as a BL amino acid difference relative to the appropriate subtype-specific reference sequence), and substitutions at the time of virologic failure at amino acid positions of interest for the NS3/4A protease (positions 155, 156, 168) and NS5A inhibitor class (positions 24, 28, 30, 31, 58, 92, 93), relative to subtype-specific reference sequences (detection threshold 15%) were determined by NGS.

## 2.3. Safety assessments

Safety of glecaprevir/pibrentasvir was assessed by monitoring adverse events (AEs) from Day 1 until 30 days after treatment completion (treatment-emergent period) and laboratory abnormalities during treatment. AEs were coded using the Medical Dictionary for Regulatory Activities (Version 21.1, MedDRA, McLean, VA) and the severity was graded according to National Cancer Institute Common Terminology Criteria for Adverse Events (Version 4.0, Bethesda, MD). AE severity and relationship to glecaprevir/pibrentasvir were determined by the investigator. Clinical laboratory samples were analyzed by a central laboratory. Hepatic laboratory abnormalities were assessed, including ALT values  $> 5 \times$  ULN, total bilirubin  $> 3 \times$  ULN, and post-nadir ALT  $> 3 \times$  ULN with total bilirubin  $\leq 2 \times$  ULN. Safety and laboratory abnormalities were summarized descriptively.

## 2.4. Statistical analysis

Analyses of BL characteristics and efficacy data were performed for the set of all patients in the intention-to-treat (ITT) population, defined as all enrolled patients who received  $\geq 1$  dose of the study drug; BL characteristics were also summarized by treatment arm. Sensitivity analyses of the primary efficacy endpoint were performed for the set of all patients in the modified ITT (mITT) population, defined as the ITT population modified to exclude patients who did not achieve SVR12 for reasons other than virologic failure. A backward imputation method was used to impute missing responses for SVR12. For the primary and secondary endpoints, two-sided 95% confidence intervals (CIs) using the Wilson's score method were provided. Subgroup analyses of the primary endpoint were also performed, with CIs provided for subgroups including  $\geq 10$  patients.

Safety analyses were performed for all patients who received  $\geq 1$  dose of glecaprevir/pibrentasvir. Treatment-emergent AEs (TEAEs) and laboratory abnormalities were summarized. Statistical Analysis System (SAS Institute, Inc., Cary, NC) Version 9.4 was used for all analyses.

## 3. Results

### 3.1. Patient characteristics

A total of 100 patients were enrolled and received  $\geq 1$  dose of glecaprevir/pibrentasvir. Seventy-five patients received 8 weeks of treatment (Arm A), and 25 received 12 weeks (Arm B). BL characteristics are summarized in Table 1. All patients completed treatment. All but one patient completed the study; one patient in Arm B was lost to follow-up. Compliance data were available for 94 patients, all of whom were compliant with the treatment.

### 3.2. Efficacy

SVR12 was achieved in 98.0% (98/100; 95% CI: 93.0–99.4) of all patients in the ITT population (Table 2). Two patients did not achieve SVR12: one (1.0%; 95% CI: 0.2–5.4) GT3a patient relapsed at PTW4, and one (1.0%) GT3a patient had a non-virologic failure (missing SVR12 data). No patients experienced OTVF (0%; 95% CI: 0.0–3.7); the patient who relapsed was a TN 61-year-old Caucasian female with HCV GT3a infection, a BL fibrosis score of F2, and a BL HCV RNA of 7.44  $\log_{10}$  IU/mL. In the mITT population, the SVR12 rate was 99.0% (98/99; 95% CI: 94.5–99.8). The SVR12 rate was  $\geq 94.9\%$  and  $\geq 97.4\%$  for all GT and fibrosis stage subgroups analyzed in the ITT and mITT populations, respectively (Fig. 2).

**Table 1**  
Baseline demographics and clinical characteristics (ITT population).

Characteristic	8 weeks Arm A N = 75	12 weeks Arm B N = 25	All patients N = 100
Male	44 (58.7)	20 (80.0)	64 (64.0)
Age, years, median (range)	55 (29–79)	58 (28–70)	56 (28–79)
Race			
White	46 (61.3)	17 (68.0)	63 (63.0)
Black or African American	22 (29.3)	4 (16.0)	26 (26.0)
Asian	1 (1.3)	1 (4.0)	2 (2.0)
Multi-race	6 (8.0)	3 (12.0)	9 (9.0)
BMI, kg/m <sup>2</sup> , median (range)	26.05 (18.12–40.08)	27.74 (23.63–45.42)	26.25 (18.12–45.42)
HCV GT <sup>a</sup>			
GT1	44 (58.7)	12 (48.0)	56 (56.0)
GT2	5 (6.7)	0	5 (5.0)
GT3	26 (34.7)	13 (52.0)	39 (39.0)
GT4/5/6	0 / 0 / 0	0 / 0 / 0	0 / 0 / 0
HCV RNA, log <sub>10</sub> IU/mL, median (range)	6.14 (3.83–7.44)	6.05 (4.73–7.35)	6.13 (3.83–7.44)
Fibrosis stage			
F2	44 (58.7)	0	44 (44.0)
F3	30 (40.0)	0	30 (30.0)
F4	1 (1.3)	25 (100)	26 (26.0)
Child–Pugh score <sup>b,c</sup>			
5	1 (1.3)	23 (92.0)	24 (24.0)
6	0	2 (8.0)	2 (2.0)
FIB-4, median (range)	1.62 (0.51–4.57)	3.32 (0.81–10.64)	1.83 (0.51–10.64)
APRI, median (range)	0.58 (0.22–3.03)	1.26 (0.41–6.93)	0.74 (0.22–6.93)
Baseline polymorphisms, n/N (%) <sup>d</sup>			
Any NS3 <sup>e</sup>	–	–	1/96 (1.0)
Any NS5A <sup>e</sup>	–	–	26/100 (26.0)
NS3 or NS5A <sup>f</sup>	–	–	27/100 (27.0)
NS3 + NS5A <sup>g</sup>	–	–	0/96

Data are n (%) unless stated otherwise.

APRI, aspartate aminotransferase to platelet ratio index; BMI, body mass index; FIB-4, fibrosis-4 index for liver fibrosis; GT, genotype; HCV, hepatitis C virus; ITT, intention-to-treat; NS, nonstructural protein; RNA, ribonucleic acid.

<sup>a</sup> Based on phylogenetic analysis, if available, otherwise from the central laboratory.

<sup>b</sup> Percentages are based on the overall patient population (N) rather than the sub-population of patients with compensated cirrhosis, hence, the sum of percentages across the different categories do not add up to 100%.

<sup>c</sup> Cirrhosis status was based on electronic data capture entries, one patient was enrolled as non-cirrhotic based on a METAVIR score of F2–3, derived from an invalid method for determining cirrhosis. The patient's valid FibroTest (BioPredictive, Paris, France) result of F4 was used in analysis and a Child–Pugh score was calculated. Therefore, the number of patients reported as cirrhotic does not match the number of patients with a Child–Pugh score result or the number of patients with fibrosis stage F4.

<sup>d</sup> Included are baseline polymorphisms in available samples at amino acid positions 155, 156, and 168 in NS3 and amino acid positions 24, 28, 30, 31, 58, 92, and 93 in NS5A at a detection level of 15%.

<sup>e</sup> “Any” indicates the total number of patients with any polymorphism within the indicated target. Total number of available sequences may vary by target.

<sup>f</sup> Patients with baseline polymorphisms in NS3 or NS5A, and includes all patients with available sequences.

<sup>g</sup> Patients with baseline polymorphisms in NS3 as well as NS5A, and only includes the patients for whom both NS3/4A and NS5A sequences are available.

**Table 2**  
SVR12 rates, virologic and non-virologic failure.

	ITT population N = 100	mITT population N = 99
SVR12,	98 (98.0)	98 (99.0)
95% CI <sup>a</sup>	93.0–99.4	94.5–99.8
Non-response	2 (2.0)	1 (1.0)
Reason for non-response		
Virologic failure	1 (1.0)	0
OTVF	0	0
Relapse	1 (1.0)	1 (1.0)
Non-virologic failure		
Missing SVR12 data	1 (1.0)	N/A

Data are n (%) unless stated otherwise.

CI, confidence interval; ITT, intention-to-treat; mITT, modified ITT; N/A, not applicable; OTVF, on-treatment virologic failure; SVR12, sustained virologic response at post-treatment Week 12.

<sup>a</sup> 2-sided 95% CI based on the Wilson score method.

### 3.3. Resistance


In this study, 56, 5, and 39 patients were infected with GT1, 2, and 3a, respectively. BL polymorphisms in NS3 were rare (1.0%, 1/96), while those in NS5A were detected in 26.0% (26/100) of the patients. BL polymorphisms in NS3 (at positions 155, 156, or 168) were not detected in GT1 and GT2-infected patients and were

detected in 2.7% (1/37) of the GT3a-infected patients (Table 3). BL polymorphisms in NS5A (at positions 24, 28, 30, 31, 58, 92, or 93) were detected in 17.9% (10/56), 100% (5/5), and 28.2% (11/39) of the GT1-, GT2-, and GT3-infected patients, respectively (Table 3). Among GT3a-infected patients, three had NS5A-A30K and one had NS5A-Y93H at BL. One GT3a-infected patient experienced virologic failure; at baseline, this patient did not have polymorphisms in NS3 and had A30-K in NS5A. At the time of failure, the patient had treatment-emergent Y93H in NS5A, while NS3 sequence was not available for analysis. BL polymorphisms in NS3 and/or NS5A had no apparent impact on the efficacy of treatment for GT1–3 infected patients.

### 3.4. Safety

Fifty-five (55.0%) patients experienced one or more TEAEs, most with a maximum severity of Grade 1 (40/55) (Table 4). The most common and only TEAE ≥10% was headache (18.0%). Four patients experienced six serious TEAEs, none of which were considered to be related to glecaprevir/pibrentasvir or led to treatment discontinuation: one patient had a Grade 3 iron deficiency anemia; one patient had a Grade 3 gastric ulcer and a Grade 3 transfusion reaction; one patient had a Grade 4 infectious diarrhea and a Grade 4 acute kidney injury; and one patient had a Grade 1 abdominal





**Fig. 2.** SVR12 rates by subgroups of interest: A. ITT population, B. mITT population. Numbers represent the number of patients with SVR12 (underlined) and the total number of patients in each group. GT, genotype; HCV, hepatitis C virus; ITT, intention-to-treat; mITT, modified ITT; SVR12, sustained virologic response at post-treatment Week 12. a) SVR12 by subgroups ITT. b) SVR12 by subgroups mITT-VF.

pain. No patients experienced a drug-related TEAE of Grade  $\geq 3$ , discontinued treatment due to an AE, or died during the study. No AEs of special interest (hepatic decompensation/hepatic failure, post-BL events of HCC) were identified. No patients experienced on-treatment hepatic laboratory abnormalities of interest (Table 4).

#### 4. Discussion

In this study of TN Brazilian patients with HCV GT1–6 infection, glecaprevir/pibrentasvir achieved high SVR12 rates of 98.0% in the ITT population and 99.0% in the mITT population. No patients experienced OTVF and one patient relapsed at PTW4. High SVR12 rates were also observed regardless of viral or host factors, including BL polymorphisms in NS3 and/or NS5A, GT, HCV RNA levels and fibrosis stage. Treatment was well tolerated with no new safety signals identified, no serious AEs were considered to be related to glecaprevir/pibrentasvir, and no AEs led to treatment discontinuation. One patient experienced virologic failure, at the time of failure, the patient had treatment-emergent Y93H in NS5A, while NS3 sequence was not available for analysis. Though NS3 sequence at the time of failure was not available for this patient, it should be noted that a pooled resistance analysis of 17 GT3a-infected virologic failures in phase 2 and 3 studies with glecaprevir/pibrentasvir regimen indicated that treatment-emergent substitutions in NS3, specifically Y56H, Q80R, A156G, or Q168(L/R), were detectable in 9 of 17 patients [12].

**Table 3**  
Baseline polymorphisms in NS3 and/or NS5A.

HCV subtype	Target	BL polymorphisms <sup>a</sup>	% (n/N)	
GT1a	NS3	Any <sup>b</sup>	(0/22)	
		NS5A	Any <sup>b</sup>	4.5 (1/22)
		L31M	4.5 (1/22)	
GT1b	NS3	Any <sup>b</sup>	(0/34)	
		NS5A	Any <sup>b</sup>	26.5 (9/34)
		L28M	2.9 (1/34)	
		R30Q	8.8 (3/34)	
		P58S	5.9 (2/34)	
		Y93H	5.9 (2/34)	
GT2a	NS3	Any <sup>b</sup>	(0/1)	
		NS5A	Any <sup>b</sup>	100 (1/1)
GT2b	NS3	L31M	100 (1/1)	
		NS5A	Any <sup>b</sup>	0 (0/2)
		Any <sup>b</sup>	100 (4/4)	
GT3a	NS3	M31L	100 (4/4)	
		Any <sup>b</sup>	2.7 (1/37)	
		R155K	2.7 (1/37)	
		NS5A	Any <sup>b</sup>	28.2 (11/39)
		S24T	2.6 (1/39)	
		M28V	5.1 (2/39)	
		A30K/S/V	17.9 (7/39)	
		P58T/S	5.1 (2/39)	
		Y93H	2.6 (1/39)	

BL, baseline; GT, genotype; HCV, hepatitis C virus; ITT, intention-to-treat; NS3, non-structural viral protein 3; NS5A, nonstructural viral protein 5A.

<sup>a</sup> Included are baseline polymorphisms in available samples at amino acid positions 155, 156, and 168 in NS3 and amino acid positions 24, 28, 30, 31, 58, 92, and 93 in NS5A at a detection threshold of 15%.

<sup>b</sup> “Any” indicates the total number of patients with any polymorphism within the indicated target. Total number of available sequences may vary by target.

**Table 4**  
Summary of adverse events.

Event	All patients N = 100
Any TEAE	55 (55.0)
Any TEAE $\geq$ Grade 3	3 (3.0)
Any serious TEAE, n (%)	4 (4.0)
Any TEAE possibly related to DAAs (glecaprevir/pibrentasvir) <sup>a</sup>	29 (29.0)
Any DAA-related TEAE with $\geq$ Grade 3	0
Any DAA-related serious TEAE	0
Any TEAE leading to discontinuation of study drug	0
Any fatal AE	0
TEAE reported in $\geq 5\%$ of patients	
Headache	18 (18.0)
Pruritus	7 (7.0)
Nausea	6 (6.0)
Fatigue	5 (5.0)
Laboratory abnormalities of interest	
ALT $> 5 \times$ ULN	0
Total bilirubin $> 3 \times$ ULN	0
Post-nadir ALT $> 3 \times$ ULN with total bilirubin $\leq 2 \times$ ULN	0

Data are n (%).

AE, adverse event; ALT, alanine aminotransferase; DAA, direct-acting antivirals; TEAE, treatment-emergent AE; ULN, upper limit of normal.

<sup>a</sup> As assessed by investigator.

The high SVR12 rates observed in this Brazilian population are similar to those observed in previous glecaprevir/pibrentasvir clinical studies [13–15]. Available real-world evidence on the use of direct-acting antivirals in Brazil supports their effectiveness in a non-clinical trial setting, and their transferability into real-world populations [16,17]. It should be noted, however, that while this study was conducted using the 12-week regimen for patients with CC, a recent study, EXPEDITION-8, has demonstrated the high efficacy and tolerability of 8 weeks of glecaprevir/pibrentasvir in TN patients with CC for all GTs [11]. These findings have led to the approval of the 8-week regimen for all TN patients in the EU and United States [7,10]. The data from EXPEDITION-8 have also been

submitted to the Brazilian Minister of Health (ANVISA) for review. Shorter treatment duration in TN patients has the potential to improve treatment adherence, reduce healthcare-associated costs [11], and may be more convenient for patients [18].

Limitations of the study include the non-comparative, open-label design of the study. No GT5 or GT6 patients were enrolled in the study, reflecting the low prevalence of GT5 and GT6 in Brazil: 0.1% and 0%, respectively [19]. Nonetheless, a similar study in non-Brazilian TN patients (ENDURANCE-5,6) investigated the efficacy and tolerability of 8-week glecaprevir/pibrentasvir in TN patients without cirrhosis and 12 weeks in TN patients with CC and demonstrated 95.7% SVR12 for GT5 (22/23) and 98.4% SVR12 for GT6 (60/61) [15]. Therefore, high SVR12 rates can be expected for GT5 and GT6 patients in the Brazilian population. Furthermore, high SVR12 rates in GT3-infected patients were observed in this study, a generally more challenging GT to treat [20].

## 5. Conclusion

In this study of Brazilian TN patients without cirrhosis or with CC infected with HCV GT1–6, glecaprevir/pibrentasvir demonstrated high efficacy and was well tolerated, with no new safety signals observed. A shorter treatment duration for TN patients with CC (8 weeks), if approved in Brazil, may further support the efforts for HCV treatment in Brazil, ultimately working toward WHO's HCV elimination target by 2030 [1].

### Abbreviations

WHO	World Health Organization
HCV	hepatitis C virus
TN	treatment-naïve
CC	compensated cirrhosis
EU	European Union
GT	genotype
PTW12	post-treatment Week 12
RNA	ribonucleic acid
HCC	hepatocellular carcinoma
HBV	hepatitis B virus
LLOQ	lower limit of quantification
ALT	alanine aminotransferase
AST	aspartate aminotransferase
ULN	upper limit of normal
NS	nonstructural protein
EOT	end of treatment
PTW4	post-treatment Week 4
SVR12	sustained virologic response at post-treatment Week 12
OTVF	on-treatment virologic failure
NGS	next-generation sequencing
BL	baseline
AE	adverse event
MedDRA	Medical Dictionary for Regulatory Activities
ITT	intention-to-treat
mITT	modified ITT
CI	confidence interval
TEAE	treatment-emergent adverse event

### Patient consent

Prior to any study-related screening procedures being performed on the patient, the informed consent statement was reviewed and signed and dated by the patient, the person who administered the informed consent and any other signatories according to local requirements.

### Author contributions

Each author either made substantial contributions to the conception or design of the work, or was involved in the acquisition, analysis, or interpretation of data for the work; and drafted the manuscript or revised it critically for important intellectual content; and provided final approval of the version to be published.

### Ethics approval

The Independent Ethics Committee (IEC)/Institutional Review Board (IRB) reviewed the ethical, scientific, and medical appropriateness of the study before it was conducted. IEC/IRB approval of the protocol, informed consent, and subject information and/or advertising, as relevant, was obtained prior to the authorization of drug shipment to a study site.

### Financial support

The study was designed and conducted by the investigators and the sponsor, AbbVie. The study analysis and financial support for the study were provided by AbbVie. AbbVie participated in the interpretation of data and in the review and approval of the content. All authors had access to all relevant data and participated in writing, review, and approval of the manuscript for publication; the corresponding author had full access to all data and the final responsibility to submit for publication.

### Conflict of interest

AbbVie sponsored the study (NCT03219216); contributed to its design; and participated in the collection, analysis, and interpretation of the data and in the writing, reviewing, and approval of the publication.

**Mario Peribañez-Gonzalez:** principle investigator of this study; invited speaker in medical events. **Hugo Cheinquer:** principle investigator of this study. **Lino Rodrigues:** employee of AbbVie and may hold stock or share options. **Maria Patelli Lima:** principle investigator of this study. **Mário Reis Álvares-da-Silva:** principle investigator of this study; advisory board, research and/or speaker for AbbVie, Bayer, Gilead, Novartis. **José Madruga:** principle investigator of this study. **Edison Roberto Parise:** principle investigator of this study. **Mário Guimarães Pessoa:** principle investigator of this study; scientific advisor, speaker; employee of University São Paulo (state University). **Juvenio Furtado:** principle investigator of this study. **Marcia Villanova:** principle investigator of this study. **Adalgisa Ferreira:** principle investigator of this study. **Felipe Mazzoleni:** principle investigator of this study. **Ecio Nascimento:** principle investigator of this study. **Giovanni Faria Silva:** principle investigator of this study; advisory board, research and/or speaker for AbbVie, Gilead, Bayer, Bristol Myers Squibb. **Linda Fredrick:** employee of AbbVie and may hold stock or share options. **Preethi Krishnan:** employee of AbbVie and may hold stock or share options. **Margaret Burroughs:** employee of AbbVie and may hold stock or share options. **Tania Reuter:** principle investigator of this study.

### Acknowledgments

AbbVie sponsored the trial (NCT03219216). We thank the patients, trial investigators, coordinators, and trial staff who made this trial possible. Glecaprevir was identified by AbbVie and Enanta. Medical writing support was provided by Annie Massa, MBiolSci, and Jenny Mayes, PhD, of Fishawack Communications Ltd.; funded by AbbVie.

## References

- [1] World Health Organization. In: Global Hepatitis Report; 2017. Available at: <http://apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf>. [Accessed December 2019].
- [2] Mesquita F, Santos ME, Benzaken A, Correa RG, Cattapan E, Sereno LS, et al. The Brazilian comprehensive response to hepatitis C: from strategic thinking to access to interferon-free therapy. *BMC Public Health* 2016;16:1132, <http://dx.doi.org/10.1186/s12889-016-3784-4>.
- [3] World Hepatitis Alliance. Sao Paulo Community Declaration on Hepatitis World Health Summit. Available at: [https://www.who.int/hepatitis/news-events/sao\\_paulo\\_community\\_declaration\\_on\\_hepatitis.pdf?ua=1](https://www.who.int/hepatitis/news-events/sao_paulo_community_declaration_on_hepatitis.pdf?ua=1). [Accessed December 2019].
- [4] World Health Organization. Progress Report on Access to Hepatitis C Treatment. Available at: <https://apps.who.int/iris/bitstream/handle/10665/260445/WHO-CDS-HIV-18.4-eng.pdf;jsessionid=A752ADD1C6C1BEC9D586B19EEF4912B1?sequence=1>. [Accessed December 2019].
- [5] Benzaken AS, Girade R, Cattapan E, Pereira GFM, de Almeida EC, Vivaldini S, et al. Hepatitis C disease burden and strategies for elimination by 2030 in Brazil. A mathematical modeling approach. *Braz J Infect Dis* 2019;23:182–90, <http://dx.doi.org/10.1016/j.bjid.2019.04.010>.
- [6] Ferreira PR, Brandao-Mello CE, Estes C, Goncalves Jr FL, Coelho HSM, Razavi H, et al. Disease burden of chronic hepatitis C in Brazil. *Braz J Infect Dis* 2015;19:363–8, <http://dx.doi.org/10.1016/j.bjid.2015.04.004>.
- [7] AbbVie. Mavyret US. Available at: [https://www.rxabbvie.com/pdf/mavyret\\_pi.pdf](https://www.rxabbvie.com/pdf/mavyret_pi.pdf). [Accessed December 2019].
- [8] AbbVie. Maviret (Glecaprevir/Pibrentasvir) 2019 application for inclusion on WHO model list of essential medicines; 2019.
- [9] AbbVie. Maviret Brazil. Available at: <https://consultas.anvisa.gov.br/#/medicamentos/25351467396201779/?nomeProduto=Maviret>. [Accessed January 2020].
- [10] European Medicines Agency. Maviret EU. Available at: [https://www.ema.europa.eu/en/documents/product-information/maviret-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/maviret-epar-product-information_en.pdf). [Accessed March 2020].
- [11] Brown Jr RS, Buti M, Rodrigues L, Chulanov V, Chuang WL, Aguilar H, et al. Glecaprevir/pibrentasvir for 8 weeks in treatment-naïve patients with chronic HCV genotypes 1–6 and compensated cirrhosis: the EXPEDITION-8 trial. *J Hepatol* 2020;72:441–9, <http://dx.doi.org/10.1016/j.jhep.2019.10.020>.
- [12] Krishnan P, Pilot-Matias T, Schnell G, Tripathi R, Ng TI, Reisch T, et al. Pooled resistance analysis in patients with hepatitis C virus genotype 1 to 6 infection treated with glecaprevir/pibrentasvir in phase 2 and 3 clinical trials. *Antimicrob Agents Chemother* 2018;62, <http://dx.doi.org/10.1128/AAC.01249-18>.
- [13] Fornis X, Lee SS, Valdes J, Lens S, Ghalib R, Aguilar H, et al. Glecaprevir plus pibrentasvir for chronic hepatitis C virus genotype 1, 2, 4, 5, or 6 infection in adults with compensated cirrhosis (EXPEDITION-1): a single-arm, open-label, multicentre phase 3 trial. *Lancet Infect Dis* 2017;17:1062–8, [http://dx.doi.org/10.1016/S1473-3099\(17\)30496-6](http://dx.doi.org/10.1016/S1473-3099(17)30496-6).
- [14] Asselah T, Kowdley KV, Zadeikis N, Wang S, Hassanein T, Horsmans Y, et al. Efficacy of glecaprevir/pibrentasvir for 8 or 12 weeks in patients with hepatitis C virus genotype 2, 4, 5, or 6 infection without cirrhosis. *Clin Gastroenterol Hepatol* 2018;16:417–26, <http://dx.doi.org/10.1016/j.cgh.2017.09.027>.
- [15] Asselah T, Lee SS, Yao BB, Nguyen T, Wong F, Mahomed A, et al. Efficacy and safety of glecaprevir/pibrentasvir in patients with chronic hepatitis C virus genotype 5 or 6 infection (ENDURANCE-5,6): an open-label, multicentre, phase 3b trial. *Lancet Gastroenterol Hepatol* 2019;4:45–51, [http://dx.doi.org/10.1016/S2468-1253\(18\)30341-8](http://dx.doi.org/10.1016/S2468-1253(18)30341-8).
- [16] Holzmann I, Tovo CV, Minme R, Leal M, Kliemann M, Ubirajara C, et al. Effectiveness of chronic hepatitis C treatment with direct-acting antivirals in the Public Health System in Brazil. *Braz J Infect Dis* 2018;22:317–22, <http://dx.doi.org/10.1016/j.bjid.2018.06.004>.
- [17] Ferreira VL, Borba HHL, Wiens A, Pedroso MLA, de Camargo Radunz VF, Ivantes CAP, et al. Effectiveness and tolerability of direct-acting antivirals for chronic hepatitis C patients in a Southern state of Brazil. *Braz J Infect Dis* 2018;22:186–92, <http://dx.doi.org/10.1016/j.bjid.2018.04.003>.
- [18] Welzel TM, Yang M, Sajeev G, Chen YJ, Pinsky B, Bao Y, et al. Assessing patient preferences for treatment decisions for new direct acting antiviral (DAA) therapies for chronic hepatitis C virus infections. *Adv Ther* 2019;36:2475–86, <http://dx.doi.org/10.1007/s12325-019-01012-6>.
- [19] Nutini MFR, Hunter J, Giron L, Pires AFNPC, Kohiyama IM, Camargo M, et al. HCV genotype profile in Brazil of mono-infected and HIV co-infected individuals: a survey representative of an entire country. *PLoS One* 2020;15:e0227082, <http://dx.doi.org/10.1371/journal.pone.0227082>.
- [20] Fathi H, Clark A, Hill NR, Dusheiko G. Effectiveness of current and future regimens for treating genotype 3 hepatitis C virus infection: a large-scale systematic review. *BMC Infect Dis* 2017;17:722, <http://dx.doi.org/10.1186/s12879-017-2820-z>.