

metabolic changes, proved to be effective in controlling this pathology.

Gender	33 H (%)		27 M (%)
	Minimum	Maximum	Average
Age (Years)	26	63	47
Weight	59	142	88
Blood glucose	88	246	96
Cholesterol (LDL)	71	226	129
Cholesterol (HDL)	37	82	41
Triglycerides	99	504	193
TGO	17	99	39
TGP	23	152	69
YGT	11	103	47
Ferritin	64	736	380
Insulin	16	41	16,21
Hidroxi-vit-D	16	49	23

<https://doi.org/10.1016/j.aohep.2021.100483>

P-124 POST PARACENTESIS COMPLICATIONS IN PATIENTS WITH DIAGNOSIS OF LIVER CIRRHOSIS

Kevin Molina

Department of Internal Medicine, Roosevelt Hospital, Guatemala, Guatemala

Introduction: Post paracentesis complications are correlated to a high morbidity and mortality rate in patients with a diagnosis of liver cirrhosis, among whom a high incidence of them has been observed after performing this procedure.

Objectives: To identify post paracentesis complications in patients diagnosed with liver cirrhosis in the Department of Internal Medicine of the Roosevelt Hospital from January 1 to December 31, 2018, Guatemala.

Population and Methods: Cross-sectional descriptive study carried out in patients with a diagnosis of liver cirrhosis who had undergone decompressive / diagnostic paracentesis.

Results: The majority of patients were male (70%) with child pugh C liver cirrhosis (71%) aged between 40 to 49 years of age (44%), with less than 1 year of diagnosis of liver cirrhosis (64%). Persistent leakage of ascites fluid from the puncture site was the most frequent complication (35%), followed by secondary bacterial peritonitis and hematoma of the abdominal wall at the puncture site (13% and 12% respectively). A third of the patients did not present any complications after the procedure (31%). Alteration in liver function tests (0.0001), decreased platelets and prolonged clotting times (0.001) presented a statistically significant relationship of greater probability of presenting some complication after the procedure, the bilirubin level did not present a statistically significant relationship for complications occur. (0.3). A third of the patients were indicated decompressive paracentesis (48%), of which a higher rate of complications was observed after the procedure (67%).

Conclusions: The most frequent complication was the persistent leakage of ascites fluid. Hypoalbuminemia, coagulopathy, and platelet alteration correlate with a higher risk of complications.

Key Words: Complications, paracentesis, liver cirrhosis

<https://doi.org/10.1016/j.aohep.2021.100484>

P-125 REX SHUNT IN A DEVELOPING COUNTRY – IS IT POSSIBLE?

Caroline Montagner Dias, Melina Utz Melere, Carolina Soares da Silva, Luiza Salgado Nader,

Cristine Suzana Trein, Alex Finger Horbe, Cristina Targa Ferreira, Riccardo Superina

¹ Hospital da Criança Santo Antônio – Porto Alegre / Brazil

² The Ann & Robert H. Lurie Children's Hospital of Chicago / USA

Introduction: Extrahepatic portal vein obstruction (EHPVO) is a frequent cause of noncirrhotic portal hypertension in children.

Objective: Describe the experience in the surgical treatment of EHPVO in children, in a developing country.

Methods: Retrospective case series study, with medical records review of patients with EHPVO, who underwent surgical treatment, by an experienced surgeon, between July 2016 and May 2019. Patient profile, laboratory test, images, liver histology, surgery performed, postoperative complications and shunt patency were analysed.

Results: 12 patients, median age of 4 years, umbilical catheterization was present in 8 patients (66,6%). Ten patients performed portography, and 60% had type A by Baveno VI criteria. Despite normal liver tests, liver biopsy revealed ductular proliferation in 83,3% of patients and mild portal fibrosis in 66,7%. Splenomegaly was present in 91,7% and thrombocytopenia in 83,3%. All patients had oesophageal varices and gastrointestinal bleeding occurred in 83,3%. Among the coagulation tests, the deficiency of C and S proteins is noteworthy in most patients, with 72,3% and 63,6% respectively.

It was possible to perform meso-Rex bypass in 10 patients (83,3%); in the other 2 distal splenorenal shunt was performed. Early post-operative complications occurred in 58,3% of patients, the most common was ascites in 4 (33,3%), which resolved in less than 1 month. One patient developed shunt thrombosis in the first 7 days after surgery, not resolved with thrombectomy. In outpatient follow-up one patient developed thrombosis in the Rex shunt and another 4 had stenosis. All of them underwent to interventional radiology. Currently 8 of 10 meso-Rex patients (80%) have patent shunt.

Conclusion: Rex shunt is possible in developing countries with an experienced surgeon and multidisciplinary team.

<https://doi.org/10.1016/j.aohep.2021.100485>

P-126 INTERPHASE HEPATITIS IN PRIMARY BILIARY CHOLANGITIS, SEVERITY FACTOR AND NO RESPONSE TO URSODEOXYCHOLIC ACID?

Felipe Ramírez¹, Alvaro Urzúa¹, Máximo Cattaneo¹, Pablo Roblero Juan¹, Jaime Poniachik¹

¹ Sección de Gastroenterología, Departamento de Medicina Interna, Hospital Clínico Universidad de Chile, Santiago, Chile

Introduction: Primary biliary cholangitis (PBC) is a chronic cholestatic disease that can progress to cirrhosis. The presence of fibrosis represents a predictive factor of progression and failure of response to ursodeoxycholic acid (UDCA). Currently, liver biopsy is not required for its diagnosis, however the finding of interface hepatitis (IH) in the histology could have a prognostic role.

Objectives: To compare in patients with biopsied PBC the presence of fibrosis and response to UDCA (Barcelona, Mayo II and Paris II criteria) according to the presence or absence of IH.

Methods: Histological findings and clinical characteristics of patients with biopsied PBC were retrospectively analyzed, at the stage when it was necessary for the diagnosis or in case of subsequent diagnostic doubt, between 2013-2019. Patients meeting the Paris criteria for PBC/HAI overlap were excluded.

Results: 36 patients were identified: 94% women, mean age 53 years (32-68), ANA (+) in 77%, elevated IgG in 58%. 11/36 with IH