

³ Department of Medicine, San Marcos National University, Lima, Perú

Introduction and Objectives: Acute on chronic liver failure is characterized by acute decompensation of chronic liver disease, associated with different organ failure and, therefore, with high mortality. Management is based on supportive treatment and liver transplantation. Successful liver transplantation in Peru began on March 24, 2000. The ACLF consensus dates back to 2009; the first patient with ACLF transplanted in Peru was performed in January 2015; she was a 61 years old woman with cryptogenic liver cirrhosis with three organ failures, ACLF - 3, with CLIF - C ACLF score of 55 points. This study aimed to stratify the different organ failures involved in acute on chronic liver failure in patients undergoing liver transplantation as treatment.

Materials and Methods: Retrospective, a descriptive study from January 2015 to April 2022, included 72 adult liver transplant patients at the "Guillermo Almenara" Hospital. Patients with Hepatopulmonary Syndrome, Liver retransplant, Combined liver-kidney transplant, Hepatorenal polycystosis, SPLIT and Domino Technique, and Pediatric patients were excluded.

Results: Of the 72 liver transplant patients, 40.3% (29 patients) had ACLF, 12 (41.4%) type 1 patients, 5 (17.2%) type 2 patients, and 12 (41.4%) type 3 patients. Average CLIF C - ACLF 50 points. The most frequent organ failure after hepatic was cerebral with encephalopathy 2 in 12 (41.4%) patients; the next failure was coagulation with INR 2 - <2.5 in 9 (31%) patients.

Conclusions: Liver transplantation represents the optimal and definitive treatment. In our casuistry, 40.3% of cirrhotic patients with ACLF were transplanted, with improvement in organ failure and survival at 28 and 90 days of 100%. The average CLIF C - ACLF score of these patients was 50.4 points, with a maximum of 70 points.

VARIABLE	ACLF 1	ACLF 2	ACLF 3
Sex			
Male, n (%)	7 (58.3)	4 (80)	5 (41.7)
Female n (%)	5 (41.7)	1 (20)	7 (58.3)
Age			
18 - 40 years	2	1	2
41 - 64 years	8	3	10
>equal 65 years	2	1	-
MELD, average, interval	25 (15 - 34)	32 (26 - 38)	32 (21 - 40)
Etiology of chronic liver disease			
NASH	4	-	4
Overlap syndrome	3	2	2
Autoimmune hepatitis	3	-	3
NASH - ASH	-	1	1
Others	2	2	2
Number organ failure, average CLIF C - ACLF			
2-Jan	12 (43)	5 (50)	-
4-Mar	-	-	8 (55)
6-May	-	-	4 (65)

<https://doi.org/10.1016/j.aohep.2023.101003>

P-120 ANTHROPOMETRIC AND METABOLIC PROFILE IN NON-ALCOHOLIC FATTY LIVER DISEASE

Lara Ramos de Prado,
Mariana Sophia Santos Almeida,
Ana Ester Amorim de Paula,
Maria Auxiliadora Nogueira Saad,
Rosa Leonora Salerno Soares, Priscila Pollo Flores,
Débora Vieira Soares

Hepatic Department, Federal University of Fluminense, Fluminense, Brazil

Introduction and Objectives: Non-alcoholic fatty liver disease (NAFLD) is the most frequent cause of liver disease, with a worldwide

prevalence of 25%. This disease is characterized by the accumulation of fat in the hepatocyte in the absence of secondary causes such as excessive alcohol consumption, drugs, or hereditary causes and can progress to steatohepatitis with or without fibrosis, cirrhosis and even hepatocellular carcinoma. The association between NAFLD and obesity, type 2 diabetes mellitus and metabolic syndrome is well established. It is estimated that approximately 76% of individuals with obesity, mainly visceral obesity, have NAFLD. In addition, previous studies have shown that simple anthropometric measures of body fat assessment, such as body mass index (BMI), neck circumference (NC), waist circumference (WC) and waist-hip ratio (WHR), are predictors of NAFLD. This study aimed to assess the prevalence of NAFLD in obese individuals and the role of anthropometric measurements that estimate visceral fat as a predictor of NAFLD.

Materials and Methods: Cross-sectional study. The study sample is a convenience sample: adults over 18 years of age, followed up at the outpatient clinics of Internal Medicine and Endocrinology of the Hospital University Antonio Pedro and at risk of NAFLD (pre-diabetes, type 2 diabetes mellitus, metabolic syndrome and/or obesity). To participate in the study, it was necessary to sign an informed consent form and clinical and anthropometric assessment, metabolic profile and liver ultrasound, elastography and electrical bioimpedance tests were performed.

Results: The evaluation was performed on 95 patients. There is a predominance of females in relation to males (81% vs. 18.9%, respectively) and a higher prevalence of alcoholism and diabetes in males (50% and 66.6%) when compared to females (18.1% and 48%). Furthermore, there is a high prevalence of physical inactivity, smoking, hypertension and dyslipidemia in both sexes. The prevalence of hepatic steatosis in 91.30% of women and 63.6% of men who underwent abdominal ultrasounds is another important observation. Anthropometric measurements such as NC, WC, and WHR are high in both sexes. Circumferences, in cm, of the neck and waist were greater in males (medians 42 cm and 106.9 cm) compared to females (medians 36.1 cm and 105 cm).

Conclusions: To date, a high prevalence of patients with visceral obesity, hepatic steatosis and metabolic diseases has been observed. Regarding the anthropometric measures of visceral obesity, they are high in both sexes, proving to be an important risk factor for NAFLD. The study is ongoing and further statistical analyzes will be performed to identify the association of hepatic steatosis with cardiometabolic diseases.

<https://doi.org/10.1016/j.aohep.2023.101004>

P-121 CORRELATION BETWEEN HEPATOPULMONARY SYNDROME AND OXYGEN SATURATION PULSE OXIMETRY IN CIRRHOTIC PATIENTS

Dayana Christo, Verônica Nicoli, Perla Schulz,
Andrea Vieira, Roberto da Silva Junior

Department of Medicine, Santa Casa de São Paulo
School of Medical Sciences, São Paulo, Brazil

Introduction and Objectives: Hepatopulmonary Syndrome (HPS) is a chronic and irreversible disease caused by systemic changes associated with portal hypertension, which greatly compromises patients' expectations and quality of life. It is associated with an increase in morbidity and mortality regardless of the degree of liver dysfunction. Data on the accuracy of the diagnosis of HPS in cirrhosis is limited. This study aimed to analyze the prevalence of HPS in cirrhotic patients at our service and to correlate it with oxygen saturation (SatO₂) using a pulse oximeter to evaluate if this is useful as a screening test for HPS.