



Original article

Hepatitis C virus NS5A and core protein induce fibrosis-related genes regulation on Huh7 cells through activation of LX2 cells

Tania G. Heredia-Torres^a, Veronica Alvarado-Martínez^a, Ana R. Rincón-Sánchez^b,
Sonia A. Lozano-Sepúlveda^a, Kame A. Galán-Huerta^a, Daniel Arellanos-Soto^a,
Ana M. Rivas-Estilla^{a,*}

^a Department of Biochemistry and Molecular Medicine, CIViM, School of Medicine, Autonomous University of Nuevo Leon, Monterrey, Nuevo León, Mexico

^b IBMMTG, Departamento de Biología Molecular y Genómica, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Jalisco, Mexico

ARTICLE INFO

Article History:

Received 8 December 2023

Accepted 26 April 2024

Available online 7 June 2024

Keywords:

Viral hepatitis

Fibrogenesis

Hepatic stellate cells activation

Co-culture

ABSTRACT

Introduction and Objectives: Liver fibrosis remains a complication derived from a chronic Hepatitis C Virus (HCV) infection even when it is resolved, and no liver antifibrotic drug has been approved. Molecular mechanisms on hepatocytes and activation of hepatic stellate cells (HSCs) play a central role in liver fibrogenesis. To elucidate molecular mechanisms, it is important to analyze pathway regulation during HSC activation and HCV infection.

Materials and Methods: We evaluate the fibrosis-associated molecular mechanisms during a co-culture of human HSCs (LX2), with human hepatocytes (Huh7) that express HCV NS5A or Core protein. We evaluated LX2 activation induced by HCV NS5A or Core expression in Huh7 cells during co-culture. We determined a fibrosis-associated gene expression profile in Huh7 that expresses NS5A or Core proteins during the co-culture with LX2.

Results: We observed that NS5A induced 8.3-, 6.7- and 4-fold changes and that Core induced 6.5-, 1.8-, and 6.2-fold changes in the collagen1, TGF β 1, and timp1 gene expression, respectively, in LX2 co-cultured with transfected Huh7. In addition, NS5A induced the expression of 30 genes while Core induced 41 genes and reduced the expression of 30 genes related to fibrosis in Huh7 cells during the co-culture with LX2, compared to control. The molecular pathways enriched from the gene expression profile were involved in TGF β signaling and the organization of extracellular matrix.

Conclusions: We demonstrated that HCV NS5A and Core protein expression regulate LX2 activation. NS5A and Core-induced LX2 activation, in turn, regulates diverse fibrosis-related gene expression at different levels in Huh7, which can be further analyzed as potential antifibrotic targets during HCV infection.

© 2024 Fundación Clínica Médica Sur, A.C. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

Hepatocellular carcinoma (HCC) is considered the main form of liver cancer and the second cause of cancer-related death worldwide, with approximately 140,000 cases attributable to chronic hepatitis C virus (HCV) infection [1]. HCV is a hepatotropic positive-sense RNA virus that encodes a polyprotein that generates different structural and non-structural proteins [2]. Chronic liver damage causes unregulated tissue regeneration, leading to progressive fibrosis. If it is not reverted, this can lead to HCC

Abbreviations: DEG, differentially expressed genes; ECM, Extracellular matrix; GO, Gene ontology; HCC, Hepatocellular carcinoma; HCV, Hepatitis C virus; HSC, Hepatic stellate cells; MMP, Metalloproteinase; RLU, relative light unit; TIMP, tissue inhibitors of metalloproteinase

* Corresponding author.

E-mail address: amrivas1@yahoo.ca (A.M. Rivas-Estilla).

<https://doi.org/10.1016/j.aohep.2024.101517>

1665-2681/© 2024 Fundación Clínica Médica Sur, A.C. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

development [3]. Unfortunately, there are no approved drugs for the treatment of liver fibrosis yet. Liver fibrosis has been characterized by excessive production of the extracellular matrix (ECM) components, including Collagen 1, and/or regulatory molecules of ECM composition, such as some metalloproteinases (MPPs) and tissue inhibitors of metalloproteinase (TIMPs) [4]. Diverse cellular processes during liver damage, including hepatic stellate cells (HSC) activation, can induce liver fibrosis.

On the other hand, HSCs are considered the most important hepatic cells in fibrosis development. HSCs can respond to liver damage through transdifferentiation and activation processes to regenerate damaged tissue through ECM components production. When the damage is chronic, HSCs continue in an activated state in conjunction with the deregulated ECM production [5]. In addition, TGF- β 1 has been shown to be one of the major HSCs activation factors, but some

HCV proteins such as NS3/4A and Core have been reported to activate HSCs through different pathways [6–8].

It is important to consider how intercellular communication between HSCs and hepatocytes during viral proteins expression could regulate the molecular mechanisms associated with fibrosis development to have a better approximation of the molecular pathogenesis in HCV infection. This can help us to find control points as potential therapeutic targets. In this study, we evaluate the fibrosis-associated molecular mechanisms regulation during a co-culture of human hepatic stellate cells (LX2), with human hepatocytes (Huh7) that express HCV NS5A protein.

2. Materials and methods

2.1. Cell culture

The human hepatoma cell line Huh-7 (donated by Dr. Koromilas; McGill University, Montreal, Canada) was cultured in Advanced Dulbecco's Modified Eagle Medium supplemented with 2 % fetal bovine serum, 2 mM L-glutamine, 100 U/mL penicillin, 100 µg/mL streptomycin and 1 % nonessential amino acids (All reagent from Gibco, USA). Human hepatic stellate cell line LX-2 (donated by Dr. María Luz Martínez Chantar; Metabolomics laboratory, CIC bioGUNE, Spain) were cultured in Dulbecco's Modified Eagle Medium, supplemented with 2 % fetal bovine serum and 100 U/mL penicillin, and 100 µg/mL streptomycin (All reagent from Gibco, USA). Both cell lines were maintained at 37 °C and 5 % CO₂.

2.2. Plasmids transfection

The pNluc-NS5A/HCV was constructed in Mammalian Gene Expression Vector with *Nluc* as a reporter (Vector Builder, VB200914-1750qfm). The pCore/HCV was constructed in the pcDNA 3.1 (+) (GenScript, USA) expression vector with the sequence of the Core protein of HCV (GenBank: AJ238799.1) (Project ID: U6951CJ230-2). For plasmids transfection, 2×10^5 Huh7 cells were cultured in transwell inserts of 6-well plates. After 24 h, the transfection was performed using Lipofectamine 3000 (Invitrogen; Thermo Fisher Scientific, USA). For each well, we used 3.75 µL of lipofectamine, 1 µg of each plasmid, and 250 µL of Opti-MEM (Gibco, USA). Since the expression of pCore/HCV is regulated by the T7 promoter, infection of Huh7 cells with *Vaccinia virus* (VR-2153, strain vTF7-3 [Wr], ATCC, USA) that encodes the T7 RNA polymerase is necessary one hour before the transfection with lipofectamine 3000 as mentioned above.

2.3. Transwell co-culture of transfected Huh7 cells with LX-2 cells

Huh7 cells (2×10^5) were plated into the 6-well format cell culture insert with 3 µm pore size (Thermo Fisher Scientific, USA). After 24 h Huh7 were transfected with 1 µg of each plasmid as mentioned above. LX-2 cells (2×10^4) were seeded in 6-well plates in DMEM without FBS 24 h before co-culture with transfected Huh7 cells. Control Huh7 or Huh7 transfected cells were co-cultured with LX-2 for 48 h and 72 h, after which LX-2 and Huh7 cells were harvested for analysis.




Fig. 1. NS5A effect on the regulation of HSC activation biomarkers in LX2 cells. Huh7 cells were transfected with pNluc-NS5A/HCV for 24 h and were co-cultured with LX2 for 48 and 72 h, then HSC activation biomarkers expression was evaluated in LX2 cells. (A) Transfection efficiency by quantitation of Copy number of pNluc-NS5A/HCV mRNA transcript by RT-qPCR in co-cultured Huh7 cells and (B) Protein level of pNluc-NS5A/HCV determined by the nanoluc activity in co-cultured Huh7 cells. (C) mRNA relative expression of *col1a1*, *acta2*, *tgfb1* and *timp1* normalized to *gapdh* and *actb* evaluated by RT-qPCR in co-cultured LX2 cells. (D) Protein levels of Collagen1, α-SMA, TGFβ1 and GAPDH by Western blot and densitometry analysis in co-cultured LX2 cells. Graphical results are presented as mean ± SEM of three independent experiments. T-test, ****p < 0.0001, ***p < 0.001, **p < 0.01, *p < 0.05.




Fig. 2. Core effect on the regulation of HSC activation biomarkers in LX2 cells. Huh7 cells were transfected with pCore/HCV for 24 h and were co-cultured with LX2 for 48 and 72 h, then HSC activation biomarkers expression was evaluated in LX2 cells. (A) Transfection efficiency by quantitation of Copy number of pCore/HCV mRNA transcript by RT-qPCR and (B) protein level of pCore/HCV determined by Western Blot in co-cultured Huh7 cells. (C) mRNA relative expression of *col1a1*, *acta2*, *tgfb1* and *timp1* normalized to *gapdh* and *actb* evaluated by RT-qPCR in co-cultured LX2 cells. (D) Protein levels of Collagen1, α -SMA, TGF β 1 and GAPDH by Western blot and densitometry analysis in co-cultured LX2 cells. Graphical results are presented as mean \pm SEM of three independent experiments. T-test, **** p < 0.0001, *** p < 0.001, ** p < 0.01, * p < 0.05.

2.4. RT-qPCR

Cells were harvested and RNA was isolated with TRIzol reagent (Invitrogen, Thermo Fisher Scientific, USA). We lysed the cells with 400 μ L of TRIzol and separated phases with 80 μ L of chloroform. RNA was precipitated with 500 μ L of isopropanol, washed with ethanol, and resuspended on RNase-free water. Reverse transcription (RT) was performed using 2 μ g of total RNA, 250 ng of random primers, 500 μ M of dNTP's, Dithiothreitol (DTT) at 10 mM, 40 Units of RNase OUT and the specific buffer of the M-MLV reverse transcriptase (Thermo Fisher Scientific, USA). Followed by this, qPCR was carried out in a StepOnePlus™ (96-well) PCR System (Applied Biosystems, USA) to quantify mRNA relative levels for *tgfb1*, *acta2*, *col1*, *timp1*, and *gapdh*, 18 s, and *actb* as endogenous controls. SYBR-Green PCR Master Mix 2 \times (Applied Biosystems, USA) was used for qPCR reactions, 200–400 nm of each primer and 50 ng of cDNA. Thermal cycling conditions were 95 $^{\circ}$ C for 10 min, followed by 40 cycles of 95 $^{\circ}$ C for 15 s and 60 $^{\circ}$ C for 60 s.

2.5. Western blot analysis

LX-2 cells were harvested and lysed in PKR buffer (0.01 M Tris-HCl pH 7.5, 0.02 mM [DTT], 0.2 M MgCl₂, 0.05 M KCl, 1 % Triton X-100, and 0.1742 μ g Phenylmethanesulfonyl fluoride [PMSF] and a

protease inhibitor cocktail) for 20 min at 4 $^{\circ}$ C (Roche, Germany). The supernatant fractions were transferred to a fresh tube after centrifugation at 13,000 rpm for one minute at 4 $^{\circ}$ C. Protein concentrations were measured using Protein Assay Dye (Bio-Rad Laboratories, USA). An equal amount of protein from each sample was boiled in standard protein sample buffer and separated via sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene difluoride (PVDF) membrane, incubated with different primary antibodies (Col1, TGF- β 1, α -SMA or GAPDH) for 16 h at 4 $^{\circ}$ C. After being washed with TBST buffer (20 mmol/L Tris-HCl pH 7.5, 150 mmol/L NaCl, 0.1 % Tween-20), they were incubated with a secondary antibody (Anti-mouse or Anti-rabbit) for 2 h at room temperature. Membranes were subsequently washed with TBST. Chemiluminescence on the membrane was detected using the SuperSignal West Pico PLUS Chemiluminescent Substrate Kit (Thermo Fisher Scientific, USA) and ChemiDoc XRS system (Bio-Rad Laboratories, USA). Densitometric analyses of protein band intensities were quantified using ImageJ software (version 1.53a; National Institutes of Health, USA).

2.6. Nanoluciferase reporter assay

Total protein insolation was used to evaluate the transfection efficiency by nanoluciferase activity as a reporter for NS5A/HCV




Fig. 3. Fibrosis-related genes expression regulation in Huh7 [pNluc-NS5A/HCV] during co-cultured with LX2 cells. Huh7 cells were transfected with pNluc-NS5A/HCV for 24 h and were co-cultured with LX2 for 48 and 72 h, then 84 fibrosis-related genes expression were evaluated in Huh7 cells. (A) Heat map of the relative expression of fibrosis-related genes evaluated by the RT2 Profiler human fibrosis PCR Array. (B) mRNA relative expression of differentially expressed fibrosis-related genes. The dotted line represents the relative mRNA expression of control group. Graphical results are presented as mean ± SEM of three independent experiments. Student T-test, ***p* < 0.01, **p* < 0.05.

expression. The nano luciferase activity in each lysate was measured using a NanoLuc assay system (Promega, USA) according to the manufacturer's instructions, an equal amount of protein samples and the mixture of buffer with the substrate (50:1) were placed into a white flat bottom 96-well plate. Relative light units were normalized and reported as RLU per μg of total protein.

2.7. Gene expression by profiler qPCR array

Huh7 cells co-cultured with LX2 cells were harvested and RNA was isolated with RNeasy Mini Kit according to the manufacturer's specifications (QIAGEN, Germany). Total RNA (1 μg) was reverse transcribed using the RT² First Strand Kit (Qiagen, Germany), and qPCR was performed using the human fibrosis PCR array (RT² Profiler PCR Array PAHS-120Z, Qiagen) according to the manufacturer's instructions. Five housekeeping genes to normalize array data (ACTB, B2M, GAPDH, HPRT1, and RPLP0), and RT and qPCR controls were included in each run. qPCR array data were analyzed using the web-based software RT² Profiler PCR Array Data Analysis, available at the manufacturer's website. The differentially expressed genes (DEGs) were analyzed by the Enrichr database online tool to obtain Gene Ontology (GO) and Signalling Pathway Enrichment analysis [9].

2.8. Statistical analysis

The statistical analysis was executed with Prism software (v9.01, Prism GraphPad, USA). Unpaired Student's T test and one-way ANOVA were applied for comparison between groups. All data in this

study were obtained from three individual experiments and presented as mean with SEM and *p* < 0.05 was considered to indicate a statistically significant difference.

2.9. Ethical statement

This was a non-interventional study where ethical approval was not required. We performed *in vitro* analysis using the hepatocarcinoma cell line (Huh7) and human hepatic stellate cells cell line (LX2). Universidad Autónoma de Nuevo León ethics committee did not require ethical approval for cell lines and non-interventional studies.

3. Results

3.1. HCV NS5A protein expression in Huh7 during co-culture regulates LX2 activation

The NS5A-Huh7 cells' effect on the activation of LX2 was evaluated. Huh7 cells were transfected with pNluc-NS5A/HCV for 24 h and then co-cultured with SFB-free LX2 cells for 48 and 72 h. We determined the transfection efficiency by the increase in NS5A mRNA copy number by qPCR and nanoluciferase activity of pNluc-NS5A/HCV in co-cultured Huh7 cells at different times (Fig. 1A and B). Therefore, we evaluated the relative expression of HSC activation biomarkers in LX2 co-cultured cells. We detected an increase in *col1*, *acta2*, and *tgfb1* mRNA relative expression in LX2 at 48 and 72 h, whereas *timp1* was increased at 48 h compared to the control (Fig. 1C). Furthermore, we observed an increase of collagen1 and TGFβ1 protein level at 48




Fig. 4. Fibrosis-related genes expression regulation in Huh7 [pCore/HCV] during co-cultured with LX2 cells. Huh7 cells were transfected with pCore/HCV for 24 h and were co-cultured with LX2 for 48 and 72 h, then 84 fibrosis-related genes expression were evaluated in Huh7 cells. (A) Heat map of the relative expression of fibrosis-related genes evaluated by the RT2 Profiler human fibrosis PCR Array. (B) mRNA relative expression of differentially expressed fibrosis-related genes. The dotted line represents the relative mRNA expression of control group. Graphical results are presented as mean ± SEM of three independent experiments. Student T-test, ##*p* < 0.0001, #*p* < 0.001, **p* < 0.01, ***p* < 0.05.

Table 1
Differentially expressed fibrosis-related genes in NS5A-Huh7 cells co-cultured with LX2.

Refseq	Gene	Description	Fold Change	p-Value
48 h				
NM_001901	CCN2	Cellular communication network factor 2	3.10	0.010
NM_000212	ITGB3	Integrin, beta 3	2.74	0.013
NM_005904	SMAD7	SMAD family member 7	2.61	0.015
NM_000362	TIMP3	TIMP metalloproteinase inhibitor 3	2.48	0.039
NM_003467	CXCR4	Chemokine (C-X-C motif) receptor 4	2.41	0.026
NM_001920	DCN	Decorin	2.39	0.035
NM_000090	COL3A1	Collagen, type III, alpha 1	2.35	0.025
NM_003238	TGFB2	Transforming growth factor, beta 2	2.31	0.025
NM_004612	TGFBR1	Transforming growth factor, beta receptor 1	2.27	0.032
NM_002424	MMP8	Matrix metalloproteinase 8	2.26	0.027
NM_000572	IL10	Interleukin 10	2.11	0.047
NM_000575	IL1A1	Interleukin 1, alpha	2.11	0.046
NM_003244	TGIF1	TGFB-induced factor homeobox 1	2.11	0.002
NM_000619	IFNG	Interferon, gamma	1.96	0.043
NM_000029	AGT	Angiotensinogen	1.19	0.040
72 h				
NM_000594	TNF	Tumor necrosis factor	2.63	0.014
NM_031479	INHBE	inhibin, beta E	2.60	0.009
NM_013372	GREM1	Gremlin 1	2.39	0.017
NM_000602	SERPINE1	Serpin peptidase inhibitor, clade E, member 1	2.18	0.011
NM_000627	LTBP1	Latent TGFB binding protein 1	1.96	0.025
NM_000660	TGFB1	Transforming growth factor, beta 1	1.96	0.008
NM_181501	ITGA1	Integrin, alpha 1	1.93	0.008

(continued)

Table 1 (Continued)

Refseq	Gene	Description	Fold Change	p-Value
NM_002317	LOX	Lysyl oxidase	1.75	0.041
NM_003246	THBS1	Thrombospondin 1	1.72	0.015
NM_002203	ITGA2	Integrin, alpha 2	1.71	0.044
NM_000029	AGT	Angiotensinogen	1.43	0.023
NM_007315	STAT1	Signal transducer activator of transcription 1	1.37	0.026
NM_003244	TGIF1	TGFB-induced factor homeobox 1	1.32	0.010
NM_004517	ILK	Integrin-linked kinase	1.26	0.022
NM_000295	SERPINA1	Serpin peptidase inhibitor, clade A, member 1	1.23	0.020

and 72 h respectively in LX2 co-cultured with NS5A-Huh7 cells (Fig. 1D). Therefore, we analyzed whether the co-culture time of Huh7 with LX2 influenced the expression of activation biomarkers in LX2 cells. Relative mRNA expression revealed that *acta2* only increases in control cells upon 72 h of co-culture. On the other hand, *col1a1* increases and *timp1* decreases at 72 h of co-culture, in LX2 co-cultured with in NS5A-Huh7 cells (Fig. S1A). Similarly, the protein level of collagen1 increases while α -SMA decreases at 72 h of co-culture, in LX2 co-cultured with NS5A-Huh7 cells (Fig. S1B).

3.2. HCV core protein expression in Huh7 during co-culture regulates LX2 activation

Likewise, Huh7 cells were transfected with pCore/HCV for 24 h and co-cultured with SFB-free LX2 cells for 48 and 72 h. We determined the transfection efficiency by measuring HCV-Core mRNA copy number by qPCR and protein expression in co-cultured Core-Huh7 cells at different times (Fig. 2A and B). Therefore, when we evaluated activation biomarkers in LX2 co-cultured cells, we detected an increase in *col1*, *tgfb1* and *timp1*




Fig. 5. Functional Enrichment and protein-protein interaction networks analysis of DEGs induced by HCV NS5A expression in Huh7 co-cultured with LX2 cells. The fibrosis-related genes evaluated by the RT2 Profiler human fibrosis PCR Array were analyzed with Enrichr and STRING bioinformatic platforms. (A) Signaling pathway enrichment and Gene Ontology analysis of the DEGs genes at 48 h. (B) Interaction networks of proteins from DEGs at 48 h of co-culture associated with different cellular functions from STRING analysis. (C) Signaling pathway enrichment and Gene Ontology analysis of the DEGs genes at 72 h. (D) Interaction networks of proteins from DEGs at 72 h of co-culture associated with different cellular functions from STRING analysis.

mRNA relative expression at 48 h of co-culture with Core-Huh7 cells compared to control (Fig. 2C). However, we did not observe difference of collagen1 and α -SMA protein, while TGF β -1 was downregulated at 72 h in co-cultured LX2 (Fig. 2D). We analyzed how the co-culture time influenced the expression of activation biomarkers in LX2 cells. Relative mRNA expression revealed that *acta2* only increases in control cells at 72 h of co-culture. On the other hand, *col1a1*, *tgfb1*, and *timp1* increase at 72 h of co-culture, in both, control and Core-Huh7 groups. (Fig. S2A). In addition, protein levels of collagen1, α -SMA, and TGF β 1 have no difference at 72 h of co-culture (Fig. S2B).

3.3. Fibrosis-related genes are regulated by HCV NS5A expression in Huh7 co-cultured with LX2

The intercellular communication between hepatocytes and HSCs during NS5A expression may impact or regulate diverse signaling pathways. Therefore, we analyzed the mRNA profiling of 84 fibrosis-related genes in NS5A-Huh7 cells after 48 and 72 h of co-culture with LX2 cells (Fig. 3A). We found overexpression in 15 genes at 48 h of co-culture, including some signal transduction TGF β superfamily

members (*DCN*, *SMAD7*, *TGFβ2*, *TGFβ1*, and *TGIF1*) and some ECM and cell adhesion molecules (*COL3A1*, *ITGB3*, *MMP8*, and *TIMP3*) (Fig. 3A). Also, we found 15 overexpressed genes at 72 h of co-culture including ECM and cell adhesion molecules (*ITGA1*, *ITGA2*, *LOX*, *SERPINA1* and *SERPINE1*), and signal transduction TGF β superfamily members (*GREM1*, *INHBE*, *LTBP1*, *TGFβ1*, *TGIF1* and *THBS1*) (Fig. 3B, Table 1).

3.4. Fibrosis-related genes are regulated by HCV core expression in Huh7 co-cultured with LX2

Likewise, we analyzed the mRNA profiling of 84 fibrosis-related genes in Core-Huh7 cells after 48 and 72 h of co-culture with LX2 cells (Fig. 4A). We found the overexpression of 29 genes at 48 h of co-culture, including some signal transduction TGF β superfamily members (*CAV1*, *SMAD2*, *TGFβ1*, *TGFβ1* and *TGFβ2*), some epithelial-mesenchymal transition molecules (*AKT*, *ILK*, *SERPINE1*, *ITGB1*) and two genes subexpressed (*CCN2* and *VEGFA*) at the same time of co-culture (Fig. 4B, Table 2). Also, we found 11 overexpressed genes like *MYC*, *JUN*, *IL4*, and *CCR2*, among others at 72 h of co-culture. In addition, we detected 29 subexpressed genes at 72 h of co-culture

including diverse ECM and cell adhesion molecules (*LOX*, *MMP14*, *PLG*, *SERPINE1*, *TIMP1*, *TIMP3*, among others), and signal transduction *TGFβ* superfamily members (*CAV1*, *INHBE*, *SMAD2*, *TGFB1*, *TGFB1*, among others) (Fig 4B, Table 3).

3.5. Molecular signaling pathways induced by HCV NS5A expression in Huh7 co-cultured with LX2

The DEGs obtained in the qPCR Array were analyzed by Enrichr (maayanlab.cloud/Enrichr/). Significantly enriched signaling pathways analysis reveals that NS5A-Huh7 cells are most associated with *TGFβ* receptor complex and ECM organization pathways at 48 and 72 h of co-culture with LX2, respectively. Furthermore, GO analysis revealed *TGFβ* receptor binding and receptor ligand activity as a molecular function enrichment at 48 and 72 h of co-culture, respectively. The Collagen-containing ECM and ECM organization were the cellular component and biological process respectively more associated in both, 48 and 72 h of co-culture (Fig. 5A, C). Additionally, we uploaded our DEGs into the STRING database (string-db.org) to identify relationships and protein-protein interaction networks among our DEGs. Different clusters were obtained depending on the

Table 2
Differentially expressed fibrosis-related genes in Core-Huh7 cells co-cultured 48 h with LX2.

Refseq	Gene	Description	Fold Change	p-Value
NM_003246	THBS1	Thrombospondin 1	1.71	0.001
NM_004995	MMP14	Matrix metalloproteinase 14	1.66	0.014
NM_005902	SMAD3	SMAD family member 3	1.64	0.019
NM_031479	INHBE	Inhibin, beta E	1.55	0.001
NM_000602	SERPINE1	Serpin peptidase inhibitor, clade E	1.55	0.001
NM_002658	PLAU	Plasminogen activator, urokinase	1.53	0.039
NM_005194	CEBPB	CCAAT/enhancer binding protein beta	1.48	0.045
NM_005904	SMAD7	SMAD family member 7	1.46	0.002
NM_002203	ITGA2	Integrin, alpha 2	1.41	0.001
NM_000660	TGFB1	Transforming growth factor, beta 1	1.37	0.001
NM_002607	PDGFA	Platelet-derived growth factor alpha polypeptide	1.36	0.001
NM_002228	JUN	Jun proto-oncogene	1.35	0.009
NM_007315	STAT1	Signal transducer and activator of transcription	1.31	0.008
NM_002213	ITGB5	Integrin, beta 5	1.26	0.013
NM_003998	NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B cells	1.24	0.002
NM_005163	AKT1	V-akt murine thymoma viral oncogene	1.22	0.016
NM_004517	ILK	Integrin-linked kinase	1.21	0.001
NM_005901	SMAD2	SMAD family member 2	1.21	0.002
NM_005359	SMAD4	SMAD family member 4	1.20	0.030
NM_001963	EGF	Epidermal growth factor	1.17	0.012
NM_000627	LTBP1	Latent TGFβ binding protein	1.16	0.017
NM_128473	SP1	Sp1 transcription factor	1.16	0.007
NM_001753	CAV1	Caveolin 1	1.15	0.035
NM_002211	ITGB1	Integrin, beta 1	1.15	0.007
NM_000029	AGT	Angiotensinogen	1.13	0.008
NM_004612	TGFB1	TGFβ receptor 1	1.11	0.047
NM_000295	SERPINA1	Serpin peptidase inhibitor, clade A	1.10	0.021
NM_000301	PLG	Plasminogen	1.07	0.019
NM_003242	TGFB2	TGFβ receptor 2	1.07	0.011
NM_001901	CCN2	Cellular communication network factor 2	0.88	0.009
NM_003376	VEGFA	Vascular endothelial growth factor A	0.75	0.018

Table 3
Differentially expressed fibrosis-related genes in Core-Huh7 cells co-cultured 72 h with LX2.

RefSeq	Gene	Description	Fold Change	p-Value
NM_001955	EDN1	Endothelin1	4.28	0.001
NM_001123	CCR2	Chemokine	4.23	0.003
NM_000589	IL-4	Interleukin 4	4.07	0.010
NM_000594	TNF	Tumor necrosis factor	3.96	0.009
NM_000089	COL1A2	Collagen, type 1, alpha 2	3.32	0.001
NM_002607	PDGFA	Platelet-derived growth factor alpha	2.20	0.003
NM_002228	JUN	Jun proto-oncogene	2.11	0.007
NM_001963	EGF	Epidermal growth factor	1.98	0.001
NM_000362	TIMP3	TIMP metalloproteinase inhibitor 3	1.51	0.004
NM_003376	VEGFA	Vascular endothelial growth factor A	1.40	0.032
NM_003244	TGIF1	TGFβ-induced factor homeobox 1	1.35	0.009
NM_002467	MYC	V-myc myelocytomatosis viral oncogene	1.19	0.027
NM_000295	SERPINA1	Serpin peptidase inhibitor, clade A, 1	0.89	0.011
NM_001235	SERPINH1	Serpin peptidase inhibitor, clade H, 1	0.79	0.018
NM_001753	CAV1	Caveolin 1	0.77	0.009
NM_005904	SMAD7	SMAD family member 7	0.77	0.048
NM_005359	SMAD4	SMAD family member 4	0.75	0.039
NM_000029	AGT	Angiotensinogen	0.74	0.043
NM_003246	THBS1	Thrombospondin 1	0.74	0.019
NM_003998	NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells	0.72	0.027
NM_005901	SMAD2	SMAD family member 2	0.70	0.006
NM_003153	STAT6	Signal transducer activator of transcription 6	0.67	0.002
NM_02213	ITGB5	Integrin, beta 5	0.63	0.001
NM_004612	TGFB1	TGFβ receptor 1	0.63	0.018
NM_181501	ITGA1	Integrin, alpha 1	0.62	0.004
NM_000660	TGFB1	Transforming growth factor, beta 1	0.61	0.004
NM_000301	PLG	Plasminogen	0.57	0.004
NM_002211	ITGB1	Integrin, beta 1	0.53	0.004
NM_005163	AKT1	V-akt murine thymoma viral oncogene	0.51	0.004
NM_007315	STAT1	Signal transducer activator of transcription 1	0.51	0.001
NM_003255	TIMP2	TIMP metalloproteinase inhibitor 2	0.51	0.008
NM_004517	ILK	Integrin-linked kinase	0.39	0.001
NM_002317	LOX	Lysyl oxidase	0.38	0.001
NM_138473	SP1	Sp1 transcription factor	0.38	0.001
NM_003254	TIMP1	TIMP metalloproteinase inhibitor 1	0.38	0.001
NM_005194	CEBPC	CCAAT/enhancer binding protein	0.37	0.021
NM_0004995	MMP14	Matrix metalloproteinase14	0.33	0.007
NM_000602	SERPINE1	Serpin peptidase inhibitor, clade E, 1	0.28	0.001
NM_002210	ITGAV	Integrin, alpha V	0.11	0.024
NM_031479	INHBE	inhibin, beta E	0.05	0.001

molecular pathway in which each protein participates and its respective interaction and cross-linking with other signaling pathways, like the *TGFβ* signaling pathway, ECM organization, and serine-type endopeptidase inhibitor among others (Fig. 5B, D)

3.6. Molecular signaling pathways induced by HCV core expression in Huh7 co-cultured with LX2

Enriched signaling pathways analysis reveals that Core-Huh7 cells are most associated with signaling by *TGFβ* receptor and signaling by interleukins pathways at 48 and 72 h of co-culture with LX2

respectively. Furthermore, GO analysis reveals that binding to TGFβ receptor and cytokine activity as a molecular function enrichment at 48 and 72 h of co-culture respectively. The granule lumen was the cellular component more associated in both, 48 and 72 h of co-culture. The TGFβ receptor pathway and negative regulation of transcription were the biological process enriched at 48 h and 72 h of co-culture respectively (Fig. 6A, C). Additionally, the STRING database showed the protein interaction in molecular pathway associated with over and subexpressed genes, at 48 h of co-culture the TGFβ receptor signaling was induced, while at 72 h, this molecular pathway was associated with subexpressed genes, and the signaling by interleukins was induced by overexpressed genes (Fig. 6B, D).

4. Discussion

Chronic Hepatitis C virus infection induces liver damage that causes fibrosis which can be reversible when the disease etiology is resolved. However, despite direct-acting antiviral implementation to resolve HCV infection, in certain cases, fibrosis development continues to occur and unfortunately, no drug has been approved for liver fibrosis treatment yet [10]. Therefore, understanding the molecular mechanisms triggered by HCV infection that induce fibrosis can help

to identify therapeutic targets for the development of antifibrotic drugs. Being co-culture of hepatocytes and HSCs one of the processes that promote fibrosis development, this study determined the HCV NS5A and Core protein role in the LX2 cells activation during the co-cultured with transfected Huh7 cells and the effect on the transcriptional regulation of diverse genes related to fibrotic processes.

We determined that pNluc-NS5A/HCV and pCore/HCV induce the activation of LX-2 due to the Collagen1, α-SMA, TGFβ1, and TIMP1 expression. These results agree with studies where the expression of different viral proteins induces the activation of HSCs [6–8,11,12]. It is important to note that transfection efficiency differ for each plasmid and by different co-culture times, resulting in higher expression of NS5A at 48 h, whereas the expression of Core was higher at 72 h of co-culture. This variation may influence the differential expression of the various markers evaluated during the study. However, we also determined whether co-culture hours influence activation by comparing the expression of the markers at 48 h with 72 h of co-culture in each group. The decrease in *timp1* expression in NS5A-Huh7 cells at 72 h compared to 48 h and no difference over hours in the control, resulted in no differences when compared NS5A-Huh7 to the control only at 72 h of co-culture. On other hand, in core transfection control, *timp1* expression showed an increase after 72 h of co-culture




Fig. 6. Functional Enrichment and protein-protein interaction networks analysis of DEGs induced by HCV Core expression in Huh7 co-cultured with LX2 cells. The fibrosis-related genes evaluated by the RT2 Profiler human fibrosis PCR Array were analyzed with Enrichr and STRING bioinformatic platforms. (A) Signaling pathway enrichment and Gene Ontology analysis of the DEGs genes at 48 h. (B) Interaction networks of proteins from DEGs at 48 h of co-culture associated with different cellular functions from STRING analysis. (C) Signaling pathway enrichment and Gene Ontology analysis of the DEGs genes at 72 h. (D) Interaction networks of proteins from DEGs at 72 h of co-culture associated with different cellular functions from STRING analysis.

compared to 48 h. Maybe that's why we couldn't detect any difference at 72 h of co-culture when comparing the control with the Core-Huh7 cells. On the other hand, no changes were observed in collagen1 protein level with pCore/HCV or pNluc-NS5A/HCV after 72 h of co-culture compared to the control group, we suggest that there is a post-transcriptional regulation and/or that some metalloproteinases (MMPs) could be activated and hence degrading collagen since it has been determined that HSC can regulate the levels of some MMPs under specific stimuli [13–15]. We observed that pNluc-NS5A/HCV and pCore/HCV seem to counteract α -SMA mRNA overexpression, since an increase in expression is seen at 48 h compared with 72 h of co-culture only in the control group. According to previous research, the expression of *acta2* increases during hours of Huh7 co-cultured with LX2 [16]. Finally, we observe that the HSC activation markers increase after 48 h of co-culture with Core-Huh7 cells, on the other hand, transfection with pNluc-NS5A/HCV induces an increase in the activation markers after 72 h of co-culture. Therefore, our results suggest that Core/HCV expression could induce activation of LX2 earlier than pNluc-NS5A/HCV.

Likewise, the fibrotic-related genes regulation was evaluated with the qPCR Array and the expression fold changes of different genes was determined at 48 and 72 h of co-culture. According to the bioinformatic analyses, with pNluc-NS5A/HCV transfection, we found that both at 48 and 72 h of co-culture, pathways such as TGF β 1 and ECM organization were enriched. However, not all overexpressed genes were the same at different times. It should be noted that overexpressed antifibrotic genes such as *IFNG* and *IL10* were obtained only at 48 h of co-culture [17,18]. On the other hand, transfection of pCore/HCV generates subexpression of diverse fibrosis-related genes at 72 h of co-culture. This could be related to the activation state of the LX2 during the co-culture, since at 72 h the activation markers were not increased. It would be necessary to determine the production of non-coding RNAs also to establish their participation in the regulation of the different molecular mechanisms triggered by intercellular communication during co-culture.

In general, we found that *SERPINE1*, *SMAD7*, *TGFB2*, *INHBE*, *LTBP1* and *THBS1* genes were overexpressed during co-cultures and this is consistent with gene expression induced by *in vitro* HCV infection in Huh7.5 cells [19]. Therefore, we can infer that these genes are regulated by the HCV proteins expression. On the other hand, *CCN2*, *COL3A1*, *TIMP3*, *LOX*, *CXCR4*, *DCN*, *TGFB2*, *THBS1* and *STAT1* genes that were found overexpressed during co-culture agree with genes overexpressed in cirrhotic liver tissue derived from HCV infection compared to healthy liver tissue [20,21]. Based on this, we suggest that the microenvironment responsible for regulating these genes is generated by the intercellular interaction during the HCV proteins expression. It is important to emphasize the limitation of this work, in our co-culture system we could not consider the role of soluble factors larger than 3 μ m, due to the pore size of the transwell membrane. Likewise, it is important to further evaluate the role of exosomes derived from both cell lines that may be present in the conditioned medium of the co-culture, to develop a more comprehensive understanding of the potential molecular mechanisms involved.

5. Conclusions

Our results demonstrated that HCV NS5A and Core proteins expression on Huh7 cells, regulates the activation of LX2 cells. Likewise, both cells under the same microenvironment, regulate diverse processes that can induce fibrosis. We highlight the importance of including studies that demonstrate the intercellular communication effect in molecular mechanisms regulation associated with HCV infection. In addition, we identified genetic signatures related to fibrotic pathways derived from the expression of HCV NS5A and Core proteins in Huh7 cells co-cultured with LX2. Although, additional studies are necessary to verify the molecular function of these genes

in liver fibrosis in order to be considered as potential antifibrotic targets.

Funding

The study was funded by the Department of Biochemistry and Molecular Medicine and CIiViM, School of Medicine, from Universidad Autónoma de Nuevo León (UANL), Monterrey, Nuevo León 64460, Mexico.

Conflicts of interests

None.

Author contributions

Tania G. Heredia-Torres: Funding acquisition, Writing – original draft, Formal analysis. **Veronica Alvarado-Martínez:** Writing – original draft, Formal analysis. **Ana R. Rincón-Sánchez:** Writing – review & editing, Funding acquisition, Project administration. **Sonia A. Lozano-Sepúlveda:** Formal analysis, Writing – original draft, Supervision, Project administration. **Kame A. Galán-Huerta:** Writing – review & editing, Formal analysis, Project administration. **Daniel Arellanos-Soto:** Supervision, Writing – review & editing, Project administration. **Ana M. Rivas-Estilla:** Conceptualization, Methodology, Funding acquisition, Supervision, Project administration.

Supplementary materials

Supplementary material associated with this article can be found in the online version at doi:10.1016/j.aohep.2024.101517.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- Dubuisson J, Cosset FL. Virology and cell biology of the hepatitis C virus life cycle - an update. *J Hepatol* 2014;61:S3–13. <https://doi.org/10.1016/j.jhep.2014.06.031>.
- Irshad M, Gupta P, Irshad K. Molecular basis of hepatocellular carcinoma induced by hepatitis C virus infection. *World J Hepatol* 2017;9:1305–14. <https://doi.org/10.4254/wjh.v9.i36.1305>.
- Roehlen N, Crouchet E, Baumert TF. Liver fibrosis: mechanistic concepts and therapeutic perspectives. *Cells* 2020;9. <https://doi.org/10.3390/cells9040875>.
- Tsushima T, Friedman SL. Mechanisms of hepatic stellate cell activation. *Nat Rev Gastroenterol Hepatol* 2017;14:397–411. <https://doi.org/10.1038/nrgastro.2017.38>.
- Coenen M, Nischalke HD, Krämer B, Langhans B, Glässner A, Schulte D, et al. Hepatitis C virus core protein induces fibrogenic actions of hepatic stellate cells via toll-like receptor 2. *Lab Invest* 2011;91:1375–82. <https://doi.org/10.1038/labinvest.2011.78>.
- Ríos-Ocampo W.A., Navas M.-C., Faber K.N., Daemen T., Moshage H. Huh7 cells expressing hepatitis C virus Core or NS3/4A protein activate human LX-2 hepatic stellate cells through paracrine signaling: preliminary results n.d.
- Cheng JC, Tseng CP, Liao MH, Peng CY, Yu JS, Chuang PH, et al. Activation of hepatic stellate cells by the ubiquitin C-terminal hydrolase 1 protein secreted from hepatitis C virus-infected hepatocytes. *Sci Rep* 2017;7:4448. <https://doi.org/10.1038/s41598-017-04259-7>.
- Xie Z, Bailey A, Kuleshov MV, Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene set knowledge discovery with enrichr. *Curr Protoc* 2021;1:e90. <https://doi.org/10.1002/cpz1.90>.
- Böttcher K, Pinzani M. Pathophysiology of liver fibrosis and the methodological barriers to the development of anti-fibrogenic agents. *Adv Drug Deliv Rev* 2017;121:3–8. <https://doi.org/10.1016/j.addr.2017.05.016>.
- Schulze-Krebs A, Preimel D, Popov Y, Bartschlagler R, Lohmann V, Pinzani M, et al. Hepatitis C virus-replicating hepatocytes induce fibrogenic activation of hepatic stellate cells. *Gastroenterology* 2005;129:246–58. <https://doi.org/10.1053/j.gastro.2005.03.089>.
- Li W, Duan X, Zhu C, Liu X, Jeyarajan AJ, Xu M, et al. Hepatitis B and Hepatitis C virus infection promote liver fibrogenesis through a TGF- β 1-induced OCT4/Nanog pathway. *J Immunol* 2022;208:672–84. <https://doi.org/10.4049/jimmunol.2001453>.
- Liu T, Wang P, Cong M, Zhang D, Liu L, Li H, et al. Matrix metalloproteinase-1 induction by diethylthiocarbamate is regulated via Akt and ERK/miR222/ETS-1 pathways in hepatic stellate cells. *Biosci Rep* 2016;36. <https://doi.org/10.1042/BSR20160111>.

- [14] Lee EH, Park K II, Kim KY, Lee JH, Jang EJ, Ku SK, et al. Liquiritigenin inhibits hepatic fibrogenesis and TGF- β 1/Smad with Hippo/YAP signal. *Phytomedicine* 2019;62:152780. <https://doi.org/10.1016/j.phymed.2018.12.003>.
- [15] Robert S, Gicquel T, Bodin A, Lagente V, Boichot E. Characterization of the MMP/TIMP imbalance and collagen production induced by IL-1 β or TNF- α release from human hepatic stellate cells. *PLoS ONE* 2016;11:e0153118. <https://doi.org/10.1371/journal.pone.0153118>.
- [16] Liu Y, Yang X, Jiang D, Hu R, Huang F, Zou X, et al. 3D biomimetic tumor microenvironment of HCC to visualize the intercellular crosstalk between hepatocytes, hepatic stellate cells, and cancer cells. *Smart Mater Med* 2023;4:384–95. <https://doi.org/10.1016/j.smaim.2022.12.002>.
- [17] Weng H, Mertens PR, Gressner AM, Dooley S. IFN- γ abrogates profibrogenic TGF- β signaling in liver by targeting expression of inhibitory and receptor Smads. *J Hepatol* 2007;46:295–303. <https://doi.org/10.1016/j.jhep.2006.09.014>.
- [18] Zhang LJ, Zheng W Da, Chen YX, Huang YH, Chen ZX, Zhang SJ, et al. Antifibrotic effects of interleukin-10 on experimental hepatic fibrosis. *Hepatogastroenterology* 2007;54:2092–8.
- [19] Woodhouse SD, Narayan R, Latham S, Lee S, Antrobus R, Gangadharan B, et al. Transcriptome sequencing, microarray, and proteomic analyses reveal cellular and metabolic impact of hepatitis C virus infection in vitro. *Hepatology* 2010;52:443–53. <https://doi.org/10.1002/hep.23733>.
- [20] Mas VR, Maluf DG, Archer KJ, Yanek K, Kong X, Kulik L, et al. Genes involved in viral carcinogenesis and tumor initiation in hepatitis C virus-induced hepatocellular carcinoma. *Mol Med* 2009;15:85–94. <https://doi.org/10.2119/molmed.2008.00110>.
- [21] He L, Yuan H, Liang J, Hong J, Qu C. Expression of hepatic stellate cell activation-related genes in HBV-, HCV-, and nonalcoholic fatty liver disease-associated fibrosis. *PLoS ONE* 2020;15:e0233702. <https://doi.org/10.1371/journal.pone.0233702>.