



Dental characterization of colombian children with non syndromic cleft lip and palate

Caracterización dental de niños colombianos con hendiduras labio palatinas no sindrómicas

Herney Alonso Rengifo Reina*

ABSTRACT

Introduction: When compared to general population, subjects afflicted with cleft lip and palate present alterations in craniofacial growth and development as well as high incidence of dental anomalies which vary according to studied population; agenesia, presence of supernumerary teeth, abnormal crown morphology and taurodontism can be counted amongst them. Objective: To assess prevalence of dental anomalies found in Colombian children with non syndromic cleft lip and palate sequels, being treated at health providing institutions. **Methodology:** A cross-sectioned descriptive, observational study was conducted on a sample of 258 medical histories and panoramic X-rays of Colombian children treated at different health providing institutions in the city of Bogota, Colombia. The sample was composed of 156/258 males (60.55%) and 102/258 (39.5%) females. Average age was 9.8 years (\pm 3.3 years). **Results:** Based on studied X-rays, it was determined that 38.4% (99/258) children presented full left unilateral cleft lip and palate sequels, 31.0% (80/258) exhibited bilateral cleft and 30.6% (79/258) suffered right unilateral cleft lip and palate. Most frequent dental anomalies found were; dental agenesia, supernumerary teeth and size anomalies. Prevalence for said anomalies were: dental agenesia, over 90%; supernumerary teeth: 40% and size anomalies: 30%. **Conclusion:** High prevalence of dental anomalies was found in children with cleft lip and palate in Bogota in concordance with information reported in scientific literature.

Key words: Cleft lip, cleft palate, dental agenesia.

Palabras clave: Labio hendido, paladar hendido, agenesia dental.

INTRODUCTION

Cleft lip and palate cases (CLP) are the most common craniofacial malformations; they constitute congenital structural deficiencies caused by defects in the fusion of cranio-facial processes which form primary and secondary palate.^{1,2} They possess multifactorial etiology and varied frequency, according to environmental and socio-cultural factors. Thus, prevalence has been reported as 1 out of 800 live births in South America, 1,8 in 1,000 live births in Europe, 1 in 750 in Asia and 1 in 1200 in Africa. In

RESUMEN

Introducción: Comparados con la población general, los sujetos con labio y paladar hendido presentan alteraciones en su crecimiento y desarrollo craneofacial y una alta prevalencia de anomalías dentales, que varía según la población estudiada, entre ellas: agenesias, presencia de dientes supernumerarios, morfología coronal anormal y taurodontismo. Objetivo: Evaluar la prevalencia de anomalías dentales encontradas en niños colombianos con secuelas de hendiduras labio palatinas no sindrómicas, atendidos en instituciones prestadoras de salud. **Metodología:** Se realizó un estudio observacional descriptivo transversal en una muestra de 258 historias clínicas y radiografías panorámicas de niños colombianos de diferentes instituciones prestadoras de salud de la ciudad de Bogotá-Colombia. De los cuales 60.5% (156/258) eran hombres y 39.5% (102/258) mujeres. El promedio de edad fue 9.8 (\pm 3.3) años. **Resultados:** De las radiografías evaluadas se determinó que 38.4% (99/258) de los niños presentaban secuelas de labio y paladar hendido unilateral izquierdo completo, 31.0% (80/258) bilateral y 30.6% (79/258) con unilateral derecho. Las principales anomalías dentales encontradas fueron: agenesias dentales, dientes supernumerarios, anomalías de tamaño. La prevalencia encontrada para cada una de ellas fue: agenesias dentales: mayor del 90%. Dientes supernumerarios: 40% y en anomalías de tamaño estuvo alrededor del 30%. **Conclusión:** Se encontraron altas prevalencias en anomalías dentales en los niños con labio y paladar hendido en Bogotá, similar a lo reportado en la literatura científica.

Colombia, according to the III National Study in Oral Health (1988) ENSAB III, prevalence close to 0.2% has been reported, (13-17). Reports in ENSAB IV for

* Graduate in Maxillary Orthopedics and Orthodontics, National University of Colombia Hospital de la Misericordia Foundation (HOMI).

Received: September 2015. Accepted: January 2016.

This article can be read in its full version in the following page: <http://www.medigraphic.com/facultadodontologiaunam>

2014 revealed 0.07%, although margins of error were high due to the sample's characteristics.³⁻⁶

CLP sequels generate esthetic, psychological and functional alterations. Esthetic alterations are mostly related to lack of continuity in the upper lip as well as scars from surgical interventions. Psychological alterations are caused by disorders in the feelings of subjects with CLP sequels as well as frame of mind disturbances and lack ability to relate to other people. Lastly, functional alterations are within the framework of phonation, deglutition and mastication disorders.

When compared to general population, CLP subjects exhibit alterations in craniofacial development and growth^{1,2,7} as well as high prevalence of dental anomalies (agenesis, presence of supernumerary teeth, abnormal crown morphology and taurodontism,^{5,8} which varies according to studied population. In general, in this type of population a 90% increase of dental anomalies is observed when compared to non-affected subjects.⁹

Within reported frequencies of dental anomalies for this population, we can find the following; microdontia: 37%, dental agenesis: over 20%.^{8,10,11} K uchler¹² reported a 15.2% taurodontism frequency. Likewise, higher presence of supernumerary teeth has been reported for CLP population when compared to general population.

Within this context, the target of the present research project was to assess prevalence of dental anomalies found in Bogota in children with sequels of non syndromic palate and lip clefts, who were treated in Health Care providing institutions in that city.

MATERIAL AND METHODS

A cross-sectioned, descriptive and observational study was conducted in children with sequels of cleft lip and palate treated in public and private health care institutions. Children's ages ranged from 5-15 years, children had not received definitive oral rehabilitation or restorative care, and were equally lacking pre-existent orthodontics and/ or maxillary orthopedic treatment. Sample size was 258 children and was calculated by way of approximation to expected frequencies (CLP) prevalence and dental anomalies.

All patients attending dental services in selected institutions and within aforementioned age ranges were accepted to form the sample. HPI statistical records were assessed in order to obtain the names of all these patients which met inclusion criteria; they had at least been subjected to first care clinical examination at the dental service; panoramic x-rays were examined at the time of attending the service

as well as patients' full medical history taken at each institution

In the study, independent variables were: socio-demographic characteristics of the child and type of cleft, result variables were dental characteristics of number, size and root development.

Based on state-of-the art circumstances, a data collecting instrument was designed in the study. Before application, the instrument was subjected to a pilot test; according to test results, adjustments were undertaken until achieving the final instrument.

X-rays were selected according to criteria of suitable sharpness, density and contrast. X-rays should lack stains, double images, marks or scratches. X-rays which were deteriorated to the point of preventing analysis of number and shape of teeth to be assessed were discarded.

Dental development was assessed according to Nolla methodology,¹³ referenced by Infante.¹⁴ Congenital tooth absence and supernumerary teeth presence were determined according to Na-Youngkim¹⁵ study, which examined congenital absence of upper lateral incisors or presence of supernumerary teeth.

In 27 patients the size of teeth adjacent to the cleft was assessed. This procedure was conducted in two ways: one with study models, assessing integrity of each model; considering as integrity the fact that represented dental tissues did not exhibit fissures or lack of continuity. A fine-point digital gauge with 0.01mm precision was used. This gauge was employed to measure the greatest mesio-distal and buccal-palatal dimension of each tooth in the arch, especially teeth which were adjacent to the cleft.

In order to collect data for the study, researchers were trained and qualified, radiographic and clinical criteria tests were undertaken, an intra-class correlation coefficient was obtained, with kappa superior to 0.8 (it was conducted with an N which corresponded to 10% of the total sample (± 25 subjects) where X-rays of subjects which were different to the final sample were assessed.

Research teams recorded information of gathering instruments. During information collection typing, quality control was assessed by means of filters, checkers and program checking lists. Gathered information was included in a database designed in Access format. It was then exported to Stata version 11.0 for analysis and procedure. Before analysis, data were checked and cleansed. Whenever inconsistencies were found the instrument was selected and compared with the appropriate diagnostic aid in order to minimize errors.

With respect to information analysis, an exploratory analysis was initially conducted, using descriptive statistics techniques, in order to determine variable distribution; a bivariate analysis was conducted using contingency tables, square chi (χ^2) and Fisher tests according to the case and distribution in number and normalcy. Paired t test was conducted for root development.

Statistical significance lesser than 0.05 had to be presented in the evaluation bivariate analysis.

During development of the present study ethical considerations of the Helsinki Declaration as well as Resolution Number 000843016¹⁶ of the Colombian Ministry of Health were observed. This research project was considered risk-free and received the endorsement of the Ethics Committee of the School of Dentistry, National University of Colombia.

RESULTS

Information was gathered on 258 children with cleft lip and palate sequels; 60% were male (156/258) and 39.5% were female (102/256). Average age was 9.8 years (± 3.3), mean was 10, minimum age 5 and maximum age 15 years. Over 70% of patients counted with some sort of affiliation to some type of Social Security Regime (30.2% contributory, 44.6% subsidized) 8.5% attended the service as private patients; the rest did not inform of any affiliation (16.7%). With respect to origin it can be said that 96.5% (249) of subjects participating in the study were from the Capital and 9 (3.5%) came from another Colombian city.

In assessed children, 38.4% (99/258) exhibited full left unilateral CLP sequels, 31.0% (80/258) presented bilateral CLP and 30.6% (79/258) full right unilateral CLP. With respect to distribution amongst gender and CLP type, no significant differences were observed between both genders ($p = 0.9$) males exhibited greater numbers. Left Unilateral CLP was the most frequently found circumstance in both male and female patients (Table I).

Dental agenesis: Prevalence of agenesis in studied population was 93.0% (240/258) regardless

of missing tooth, with an average of 1.8 (± 1.0) missing teeth. With respect to agenesis and gender distribution, greater number of dental agenesis in males was observed (58.7%) compared with females (44.2%) with significant differences ($p = 0.04$)

Left upper lateral incisors were the most frequently absent teeth (61.6%) followed by their contra-lateral counterparts (58.1%) (Table II). The most frequent combination of missing teeth were those found between lateral and premolar in the affected site (8.5%) followed by central and premolar. (7.7%). No significant differences were found with respect to distribution of agenesis and type of CLP ($p = 0.23$ square chi (χ^2), Fisher)

With respect to dental agenesis type and their relationship with CLP types, significant differences were found for right unilateral CLP at the right lateral incisor ($p = 0.00$); with respect to left unilateral CLP differences were found for left lateral incisor ($p = 0.00$), left central incisor ($p = 0.02$), left upper canine ($p = 0.05$) and left first premolar ($p = 0.05$).

Supernumerary teeth: Prevalence of supernumerary teeth in studied population was 42.6% with an average of 1.4 teeth (± 0.7) Most children presented only one supernumerary tooth, whereas 13.2% exhibited more than one, there even was a patient with six supernumerary teeth. No significant differences were found with respect to gender and supernumerary teeth ($p = 0.36$).

Significant differences were found with respect to CLP type and amount of supernumerary teeth ($p = 0.02$); the greatest number of supernumerary teeth were found in left unilateral CLP. With respect to supernumerary tooth type, significant differences ($p < 0.05$) were found only in upper lateral supernumerary teeth.

Upper lateral central Incisors were the most frequent supernumerary teeth found (its presence was simultaneously found with the normal-nomenclature tooth) (Table III). It is important to note that in the present case, the object of observation were teeth of CLP subjects.

Tooth size anomalies: Anomalies which were clearly different from normal tooth size (not associated

Table I. Distribution of cleft lip and palate type according to gender.

Gender	Right		Left		Bilateral		Total
	N	%	N	%	N	%	
Female	32	40.5	39	39.4	31	38.7	102
Male	47	59.5	60	60.6	49	61.3	156
Total	79	100	99	100	80	100	258

to radiographic distortion) were radiographically assessed and determined. Each tooth was compared with its contra-lateral opponent, Thus, microdontia and macrodontia were observed in 75 subjects (29.1%).

Proportion of dental size anomalies in studied subjects were 4% macrodontia and 96% microdontia. Microdontia cases were mainly found in the left upper lateral incisor adjacent to the cleft (supernumerary teeth smaller than expected size were excluded from the analysis).

A sub-sample of 27 study models of this population was equally taken. Upper incisor size was measured with a digital gauge, it was observed that 82.5% exhibited tooth-size differences. Of these, 69% presented a difference lesser than 0.6 mm, the rest of

the percentage exhibited differences greater than 0.6 mm taurodontism was included within the categories of size anomalies; it was observed in 19.4% (50) of cases, its prevalence was higher in upper permanent molars (Table IV).

No significant differences were found ($p = 0.37$) with respect to CLP type and dental size anomalies (microdontia, macrodontais and taurodontism) (Figure 1). Differences according to gender did not exhibit statistical significance ($p = 0.22$).

Anomalies in eruption sequence: Anomalies in eruption sequence were observed in 12% (31/258) of all patients. No statistical differences were found according to gender (0.32), but they were encountered with respect to CLP type ($p = 0.02$). Greater frequency of sequence anomalies were found in children with left unilateral CLP sequels (58.1%) especially in upper canines.

With respect to Nolla stage, it was observed that there were canine-to-canine differences among homologous teeth, but no significant differences were found. In canines, differences of up to two Nolla root development stages were found, but this did not exhibit statistical significance.

Table II. Distribution of dental agenesis according to tooth type.

Missing tooth	Frequency*	Percentage
Left upper lateral incisor	159	61.6
Right upper lateral incisor	150	58.1
Right upper second premolar	32	12.4
Left lateral second premolar	30	11.6
Left upper central incisor	11	4.3
Left upper first premolar	11	4.3
Right upper first premolar	9	3.5
Left lower second premolar	8	3.2
Right upper central incisor	8	3.1
Left upper canine	7	2.7
Right lower second premolar	4	1.6
Right upper canine	4	1.5
Left lower first premolar	2	0.8
Right lower lateral incisor	1	0.4

* Only population which exhibited supernumerary teeth.

Table III. Frequency of supernumeraries according to tooth type.*

Of tooth	Frequency	Percentage
Type	5	3.3
Right upper central	52	34.4
Right upper lateral	75	49.7
Left upper lateral	0	0.0
Left upper canine	5	3.3
Right second premolar	6	4.0
Left upper central	1	0.7
Left upper central	4	2.6
Right lower lateral	1	0.7
Left second premolar	1	0.7
Right third molar	1	0.7
Total	151	100

* Only population which exhibited supernumerary teeth.

Table IV. Taurodontism distribution according to tooth presenting it.

Tooth	Frequency*	Percentage
Right upper first molar	46	32.2
Left upper first molar	44	30.7
Right lower first molar	26	18.2
Left lower first molar	26	18.2
Second molars	1	0.7

* Number of teeth.




Figure 1. Percentile distribution of anomalies of tooth size and cleft lip and palate type.

Position anomalies: Presence of position anomaly was observed in 22.1% cases (57/258) of studied population; rotations were the most common position anomalies (72%) followed by mesial and distal versions (17%) and (10.5%).

Transposition anomaly was observed in 4.3% of children; the most frequent (2.7%) was seen in cases when the upper canine assumed the position of the upper first premolar. Tooth impaction was observed in 3.8% of cases (10/258); the most frequent impaction was observed at the left upper first molar (1.2%). Likewise, impaction of upper and lower molars was observed, but in percentages lower than 1%.

Other findings: In this group of studied children presence of peg teeth was detected (especially lateral incisors), as shown in their clinical history. Likewise, radiographic findings also exhibited marked root laceration and root dwarfism among others (*Table V*).

DISCUSSION

High values of dental agenesis were found (93.0%). These values were similar to those reported by Akcam⁹ in 2010, who indicated 97.1% frequencies at that time. Values of the present study were above those reported by Shapira¹⁷ in 2000 (77%), Aizenbud¹⁸ in 2011 (68%), Jamal⁵ in 2010 (67%) and Slayton³⁰ in 2003 (47.5%). *Table VI* depicts a comparison of our findings with other findings reported in literature.

Lateral incisors were the most frequent absent teeth; this was similar to Baek's¹⁹ reports in Korea, Menezes,²⁰ Genovez¹⁰ in Brazil and Jamal.⁵ Percentages of dental agenesis (over 50%) of lateral incisors were similar to those reported by Wu²¹ in 2011.

In scientific literature, different authors^{18,22} mention that the upper second premolar (USP) was the most commonly found absent tooth, with percentages above 10%; these percentages were similar to those

reported by Shapira²³ in 1999 , Halpern²⁴ in 2010 and Camporesi²⁵ in 2010. Frequencies found in the present study were well below the percentage reported by Menezes,²⁰ who indicated a percentage above 30%, and by Tortora²⁶ (25%). With respect to central incisor absence, results of our study concurred with those found in literature:²¹ they were close to 3%.

With respect to mandibular teeth agenesis, findings of the present study were similar to those reported in 2010 with percentages close to 4% for premolars, but they were below reported frequency for incisors (0.4% in this study, whereas literature reports frequencies above 2%).

With respect to supernumerary teeth, in the present study, prevalence found was in the 43% range: they were similar to those reported by Menezes²⁰ for Brazil and Amaral²⁷ in India; they differ from reports by Genovez¹⁰ who indicated 12% prevalence of supernumerary teeth, Jamal⁵ reported 17% frequency and Camporesi and Baccetti²⁵ reported 5.2% frequency in 2010. Ramos²⁸ found significant differences with respect to supernumerary teeth and gender; he established greater number in males than in females, this fact was not observed in the present study.

Literature²¹ reports frequency of microdontia-type dental anomalies, where, like in the present study, greater frequency in teeth adjacent to the cleft²⁹ as well as in lateral incisors are observed.¹² In the present study, microdontia percentages were similar to those reported by Jamal,⁵ Genovez¹⁰ and Akcam⁹ in studies published in 2010, where they reported prevalences above 30%.

With respect to microdontia in lateral incisors a 28% frequency was observed, this was similar to reports of Genovez¹⁰ and Akcam⁹ who indicated presence of 29% frequency. Additionally, it can be said that frequency found in the present study was below that reported by Peteka in 1993³¹ who informed of a prevalence close to 60% in a population of Portuguese children.

With respect to Taurodontism prevalence, results of the present study (20%) were very different from those reported in literature for this type of population: literature shows ambivalent percentages, there are studies¹² reporting 15.2% frequencies and others which inform of an over 70% frequency.^{5,30} Literature reports did not individualize teeth which presented taurodontism as was the case in this project.

When describing patients with oral and facial clefts, literature³² informs of delays in development and eruption patterns of the permanent dentition, and alterations in different degrees; similar findings were observed in the present study. Nevertheless, no evidence was found in differences by gender, in contraposition to other findings in literature which show greater female prevalence.

Table V. Distinct tooth characteristics in cleft lip and palate patients.

Finding	Frequency	Percentage
Root dwarfism	7	2.7
Peg tooth**	4	1.5
Pronounced dilacerations***	2	0.8
Submerged primary tooth	3	1.1
Tooth fusion*	3	1.1
Mesiodens	4	1.7

* Fusion took place between lower lateral incisors and lower incisors of the same dental quadrant. ** Dilacerations were observed in upper lateral incisors as well as *** peg-shaped crown.

Table VI. Comparison of present study's findings with other findings in literature.

Assessed ITEM	Rengifo-Yesioro		Literature		
	Prevalence	N	Prevalence		Year
Agenesis	93.0	122	97	Akcam ⁹	2010
		278	77	Shapira ¹⁷	2000
		179	68	Aizenbud ¹⁸	2011
		44	67	Jamal ⁵	2010
		120	48	Slayton ³⁰	2003
		205	70	Genovez ¹⁰	2010
Left laterals	61.6	164	29	Olin ²²	1964
		146	12	Menezes ²⁰	2007
		205	55	Genovez ¹⁰	2010
Right laterals	58.1	87	49	Tortora ²⁶	2008
		196	50	Wu ²¹	2011
Premolars	10	87	25	Tortora ²⁶	2008
		146	30	Menezes ²⁰	2008
		278	18	Shapira ¹⁷	2000
		38	12	Halpern ²⁴	2010
		156	5	Camporesi ²⁵	2010
Mandibular teeth	3.2		4.0	Kontos ³⁵	2001
Microdontia	29.1	205	29	Genovez ¹⁰	2010
		122	29	Akcam ⁹	2010
Lateral Microdontia	27.9	205	29	Genovez ¹⁰	2010
		122	29	Akcam ⁹	2010
			59	Peterka ³¹	1983
Peg-shaped teeth	1.5	316	18-34	Walker ²⁹	2009
		196	No data	Wu ²¹	2011
Taurodontism	19.4	120	15.2	Slayton ³⁰	2003
		44	70	Jamal ⁵	2010
Supernumerary teeth	43	146	41	Menezes ²⁰	2007
		96	62	Amaral ²⁷	2007
		205	12	Genovez ¹⁰	2010
		44	17	Jamal ⁵	2010
		156	5	Camporesi ²⁵	2010
Transposition	4.3	44	30	Jamal ⁵	2010

Rodrigues,³³ in 2005, when studying root development, found differences in lateral incisors, whereas in the present study these differences were found in canines. With respect to CLP type, no differences were found in root development, this concurs with studies executed by Solis³² since 1998.

Jamal⁵ reported transposition prevalence in the vicinity of 30%. These figures were well above those found in the present study (4.3%). Nevertheless, concurring with literature,³⁴ greater transposition frequency between canine and upper first premolar was found.

With respect to tooth impaction, the present study is one of the first to evidence percentages of 3.8% since literature²¹ only mentions impaction frequency but does not report values for these impactions, underlining statistical differences ($p = 0.01$) where

higher impaction frequency was observed in patients with Left Unilateral CLP.

Several findings of the present study show similarities with literature reports when studying peg-shaped^{26,35} laterals, but with different frequencies. Concurrence with literature reports were also found with anomalies such as dilacerations and root dwarfism.⁵

CONCLUSIONS

In order to conclude the present article it is important to highlight that high prevalence of dental anomalies were found in Bogota in children with left cleft lip and palate, in a similar proportion as that reported in scientific literature. Within the limited scope of the present study, use of panoramic X-rays from a single radiological center was established, nevertheless, due

to Colombian Health System and characteristics of care provided for Colombian population, standardization was impossible, nevertheless, minimization of this variability was attempted with the valuator's standardization (kappa over 80%) as well as use of radiographs before initiating any type of treatment. It is important to continue characterizing population with sequels of cleft lip and palate and care, targeting day-today improvement of care offered to this population.

REFERENCES

- Avery J, Chiego D. *Principios de histología y embriología bucal*. 3a ed. Elsevier.com/Avery/histology. 1997.
- Moore K, Persaud TV. *Embriología básica*. 5a ed. McGraw-Hill Interamericana; Ciudad de México: 1998.
- República de Colombia, Ministerio de Salud. *III Estudio Nacional de Salud Bucal ENSAB III, II Estudio Nacional de Factores de Riesgo de Enfermedades Crónicas ENFREC II*. 1999. Santa fe de Bogotá.
- República de Colombia. Ministerio de Salud y Protección Social. *IV Estudio de Salud Bucal-ENSAB IV*. Bogotá 2014.
- Al Jamal GA, Hazza'a AM, Rawashdeh MA. Prevalence of dental anomalies in population of cleft lip and palate patients. *Cleft Palate Craniofac J*. 2010; 47 (4): 413-420.
- Haring FN. Dental development in cleft and noncleft subjects. *Angle Orthod*. 1976; 46 (1): 47-50.
- Murray JC. Gene/environment causes of cleft lip and/or palate. *Clin Genet*. 2002; 61 (4): 248-256.
- Lai MC, King NM, Wong HM. Dental development of chinese children with cleft lip and palate. *Cleft Palate Craniofac J*. 2008; 45 (3): 289-296.
- Akcam MO, Evirgen S, Uslu O, Memikoğlu UT. Dental anomalies in individuals with cleft lip and/or palate. *Eur J Orthod*. 2010; 32 (2): 207-213.
- Tereza GP, Carrara CF, Costa B. Tooth abnormalities of number and position in the permanent dentition of patients with complete bilateral cleft lip and palate. *Cleft Palate Craniofac J*. 2010; 47 (3): 247-252.
- Thilander B, Pena L, Infante C, Parada SS, de Mayorga C. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod*. 2001; 23 (2): 153-167.
- Küchler EC, da Motta LG, Vieira AR, Granjeiro JM. Side of dental anomalies and taurodontism as potential clinical markers for cleft subphenotypes. *Cleft Palate Craniofac J*. 2011; 48 (1): 103-108.
- Nolla C. The development of the permanent teeth. *J Dent Chi*. 1960; 27: 254-266.
- Infante C, Rosas LM, Benavides B. *Manual de ortopedia maxilar. Modelo diagnostico de maloclusiones para pacientes en crecimiento*. Bogotá: Universidad Nacional de Colombia. Facultad de Odontología. 2010.
- Kim NY, Baek SH. Cleft sidedness and congenitally missing or malformed permanent maxillary lateral incisors in Korean patients with unilateral cleft lip and alveolus or unilateral cleft lip and palate. *Am J Orthod Dentofacial Orthop*. 2006; 130 (6): 752-758.
- República de Colombia. Ministerio de Salud. Resolución 8430. Bogotá. 1993.
- Shapira Y, Lubit E, Kuflinec MM. Hypodontia in children with various types of clefts. *Angle Orthod*. 2000; 70 (1): 16-21.
- Aizenbud D, Coval M, Hazan-Molina H, Harari D. Isolated soft tissue cleft lip: epidemiology and associated dental anomalies. *Oral Dis*. 2011; 17 (2): 221-231.
- Baek SH, Kim NY. Congenital missing permanent teeth in Korean unilateral cleft lip and alveolus and unilateral cleft lip and palate patients. *Angle Orthod*. 2007; 77 (1): 88-93.
- Letra A, Menezes L, Granjeiro JM, Viera AR. Defining subphenotypes for oral clefts based on dental development. *J Dent Res*. 2007; 86 (10): 986-991.
- Wu TT, Chen PK, Lo LJ, Cheng MC, Ko EW. The characteristics and distribution of dental anomalies in patients with cleft. *Chang Gung Med J*. 2011; 34 (3): 306-314.
- Olin WH. Dental anomalies in cleft lip and palate patients. *Angle Orthod*. 1964; 34 (2): 119-123.
- Shapira Y, Lubit E, Kuflinec MM. Congenitally missing second premolars in cleft lips and cleft palate children. *Am J Orthod Dentofacial Orthop*. 1999; 115 (4): 396-400.
- Halpern RM, Noble J. Location and presence of permanent teeth in a complete bilateral cleft lip and palate population. *Angle Orthod*. 2010; 80 (3): 591-596.
- Camporesi M, Baccetti T, Marinelli A, Defraia E, Franchi L. Maxillary dental anomalies in children with cleft lip and palate: a controlled study. *Int J Paediatr Dent*. 2010; 20 (6): 442-450.
- Tortora C, Meazzini MC, Garattini G, Brusati R. Prevalence of abnormalities in dental structure, position, and eruption pattern in a population of unilateral and bilateral cleft lip and palate patients cleft palate. *Cleft Palate Craniofac J*. 2008; 45 (2): 154-162.
- Deepti A, Muthu MS, Kumar NS. Root development of permanent lateral incisor in cleft lip and palate children: a radiographic study. *Indian J Dent Res*. 2007; 18 (2): 82-86.
- Da Silva AP, Costa B, Carvalho LF, Carrara CF. Dental anomalies of number in the permanent dentition of patients with bilateral cleft lip: radiographic study. *Cleft Palate Craniofac J*. 2008; 45 (5): 473-476.
- Walker SC, Mattick CR, Hobson RS, Steen IN. Abnormal tooth size and morphology in subjects with cleft lip and/or palate in the north of England. *Eur J Orthod*. 2009; 31: 68-75.
- Slayton RL, Williams L, Murray JC, Wheeler JJ, Lidral AC, Nishimura CJ. Genetic association studies of cleft lip and/or with hypodontia outside the cleft region. *Cleft Palate Craniofac J*. 2003; 40 (3): 274-279.
- Peterka M, Müllerová Z. Tooth size in children with cleft lip and palate. *Cleft Palate J*. 1983; 20 (4): 307-313.
- Solis A, Figueroa AA, Cohen M, Polley JW, Evans CA. Maxillary dental development in complete unilateral alveolar clefts. *Cleft Palate Craniofacial J*. 1998; 35 (4): 320-328.
- Pioto NR, Costa B, Gomide MR. Dental development of the permanent lateral incisor in patients with incomplete and complete unilateral cleft lip. *Cleft Palate Craniofac J*. 2005; 42 (5): 517-520.
- Pair J. Transposition of a maxillary canine and a lateral incisor and use of cone-beam computed tomography for treatment planning. *Am J Orthod Dentofacial Orthop*. 2011; 139 (6): 834-844.
- Kontos K, Friede H, Cintras H, Celso LB, Lilia J. Maxillary development and dental occlusion in patients with unilateral cleft lip and palate after combined velar closure and lip-nose repair at different ages. *Scand J Plast Reconstr Surg Hand Surg*. 2001; 35 (4): 377-386.

Mailing address:

Herney Alonso Rengifo Reina

E-mail: hrengifo@gmail.com