



Research note

***Thompsodinium intermedium* (Dinophyceae), a freshwater planktonic species in Mexico**

***Thompsodinium intermedium* (Dinophyceae), una especie planctónica de agua dulce en México**

José A. Aké-Castillo[✉]

Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana. Calle Hidalgo 617, Col. Río Jamapa, 94290 Boca del Río, Veracruz, México.

[✉] aake@uv.mx

Abstract. Morphological description of *Thompsodinium intermedium* found in a karstic sinkhole Dzityá, in Yucatán is provided. This is the only locality in Mexico where the species has been recorded. Taxonomic circumscription of the species has changed considerably since its original description. An emended diagnosis for the species is provided, as well as for the genus. Worldwide distribution of *Thompsodinium intermedium* is provided.

Key words: dinoflagellate, emended diagnosis, karstic sinkhole, thecae.

Resumen. Se proporciona la descripción morfológica de *Thompsodinium intermedium* encontrado en el cenote Dzityá en Yucatán, único registro en México. La circunscripción de la especie ha cambiado considerablemente desde su descripción original, por lo que se propone una diagnosis enmendada para la especie y para el género. Se muestra la distribución mundial de *Thompsodinium intermedium*.

Palabras clave: dinoflagelado, enmienda, cenote, teca.

Cenote Dzityá (or Chen-Ha) is a small karstic sinkhole (10.89 x 7.40 m, 4.60 m mean depth) located close to Mérida City, Yucatán Mexico (21.06° N, 89.69° W). In 2 samples of phytoplankton collected in April and May 1999 from this sinkhole, the occurrence of a dinoflagellate was observed. This microalga was studied by light (Leica dml) and electron microscopes (SEM-JEOL-5600). The morphology indicates that the species belongs to the genus *Thompsodinium* Bourrelly. Description of the species follows:

Thompsodinium intermedium (Thompson) Bourrelly 1970: the cells are 20-32 µm in width and 22-33 µm in length with a ratio width/height of 0.7-0.95, so the cells are ovate or spherical. The cells are dorsoventrally slightly compressed. Many small chloroplasts and a small eyespot are present. Tabulation is Po, x, 4', 3a, 6'', 6c, 4s, 5''', 2'''' (Figs. 1, 2). The second intercalary plate and the third apical plate can be in "remotum" or "contactum" position. The sulcal posterior plate is large extending to the antapex looking like a plate of hypotheca. The third and fourth

postcigular plates are large and the antapical plates are dissimilar in size and shape.

Diagnostic characters of the species have changed in the interpretation of the type of hypothecal and sulcal plates (Table 1). Nowadays, the intercalary plate at hypotheca described by Thompson (1950) is known to be a large posterior sulcal plate (Sp) (Carty 1989). Since the original description is not correct anymore, an emended diagnosis of the species (and therefore the monotypic genus) is necessary to attain an adequate taxonomic circumscription of the species:


Thompsodinium Bourrelly emend. Aké-Castillo, herein.

Cells with tabulation Po, 4', 3a, 6'', 6c, 4s, 5''', 2''''. Arrangement of the 2a and 3' plates in remotum, contactum or conjunction position.

Type species: *Thompsodinium intermedium* (Thompson) Bourrelly emend. Aké-Castillo, herein.

Thompsodinium intermedium (Thompson) Bourrelly emend. Aké-Castillo, herein.

Basynonym: *Peridinium intermedium* Thompson 1950 Lloydia 13:277-299 p. 298 figs 80-88 (non *Peridinium intermedium* Playfair 1919).



Figures 1-2. *Thompsodinium intermedium* (scanning electron microscope). 1, Plate arrangement in remotum position; 2, sulcal plates: anterior plate (Sa), small left plate (Ss), right plate (Sd), large anterior plate (Sp). Scale bars 10 µm.



Figure 3. Worldwide distribution of *Thompsodinium intermedium*. Records in chronological order: 1) Thompson, 1950; 2) Popovský, 1970; 3) Couté and Iltis, 1984; 4) Carty, 1989; 5) Ngô and Pfiester, 1990; 6) Carty, 1993; 7) Carty and Wujek, 2003; 8) Johansen et al., 2007; 9) Cardoso and Torgan, 2007; 10) Krakhmalny, 2011; 11) this work.

Synonyms (taxonomic): *Peridinium pseudointermedium* Couté and Iltis 1984, *Thompsodinium pseudointermedium* (Couté and Iltis) Krakhmalny 2011.

Ovate to spherical cells with 20-42 µm wide and 22-47 µm in length. Tabulation Po, x, 4', 3a, 6'', 6c, 4s, 5''', 2'''' with a distinctive large posterior sulcal plate.

Lectotype: Figs. 80-88 in Thompson 1950 (designated by Bourrelly 1970).

Type locality: Chanute, Kansas, USA.

Distribution: freshwater environments in USA, Mexico, Belize, Brazil, Ivory Coast and Ukraine (Fig. 3).

The occurrence of this dinoflagellate in Mexico is known until now only from cenote Dzityá since 1988. It is the dominant species (Sánchez-Molina et al., 1994) and causes blooms with cell densities between 254 860 and 550 370 cells/l (Pacheco, 1988).

I thank Víctor Ontiveros who sampled cenote Dzityá; Nery Segovia, from Biblioteca-Ciencias Biológicas y

Table 1. Thecal information of *Thompsodinium intermedium*

Species	Measures (μm)	Plate formula*	Additional character	Location	Reference
<i>Peridinium intermedium</i>	32-39 long, 33-40 wide	4', 3a, 6'', 5''', 1p, 2''', remotum or contactum	No	Abandoned clay pit, Chanute, Kansas, USA	Thompson (1950)
<i>Peridinium intermedium</i>	30-46 long, 28-40 diameter	4', 3a, 6'', 5''', 2'''	Wing-like rim	Laguna del Tesoro, Cuba	Popovský (1970)
<i>Thompsodinium intermedium</i>	No information	4', 3a, 6'', 5''', 1p, 2'''	No	No information	Bourrelly (1970)
<i>Peridinium pseudointermedium</i>	36-40 long, 30-33 wide	Po, 4', 3a, 6'', 5''', 2'''	No	Ivory Coast, Maraoué, Africa	Couté and Iltis (1984)
<i>Thompsodinium intermedium</i>	No information	Po, 4', 3a, 6'', 6c, 4s, 5''', 2''', remotum or conjunctum	Wing like flange	Small ponds in Brazos County, Texas, USA	Carty (1989)
<i>Thompsodinium intermedium</i>	32-39 x 33-40	3', 4a, 6'', 5''', 1p, 2'''' or 4', 3a, 6'', 5''', 2p, 1''''	No	USA and Cuba	Popovský and Pfiester (1990)
<i>Thompsodinium intermedium</i>	No information	4', 3a, 6'', 5'', 2'''	antapical plates with fine teeth subtending the sulcus	Ponds in Tiffin, Ohio, USA	Carty (1993)
<i>Thompsodinium intermedium</i>	39.6-46.2 × 29.0-41.8	Po, x,, 4', 3a, 6'', 5c, 4s, 5''', 2'''	Sometimes 4 small spines	Lake Buchack Ukraine	Krakhmalny (2011)
<i>Thompsodinium intermedium</i>	22-33 length 20-32 wide	Po, x, 4', 3a, 6'', 6c, 4s, 5''', 2''', remotum or contactum	No	Cenote Dzityá, Mexico	This work

*given or based in information provided by the author

Agropecuarias, UADY; Angela Pacheco from Biblioteca Central, UNAM and Arlette Fuentes from INECOL facilitated acquisition of literature. A. F. Krakhmalny kindly sent basic papers. Tiburcio Láez assisted at the Scanning Electron Microscope Unit, INECOL, A. C.

Literature cited

- Bourrelly, P. 1970. Les algues d'eau douce: initiation à la systématique. 3. Les algues vertes. Boubée N. Paris. 572 p.
- Carty, S. 1989. *Thompsodinium* and two Species of *Peridiniopsis* (Dinophyceae): taxonomic notes based on scanning electron micrographs. Transactions of the American Microscopical Society 108:64-73.
- Carty, S. 1993. Contribution to the dinoflagellate flora of Ohio. Ohio Journal of Science 93:140-146.
- Carty, S. and D. Wujek. 2003. A new species of *Peridinium* and new records of dinoflagellates and silica-scaled chrysophytes from Belize. Caribbean Journal of Science 39:136-139.
- Cardoso, L. S. and L. C. Torgan. 2007. Dinoflagelados em diversos habitats e hidropériodos na zona costeira do sul do Brasil. Acta Botanica Brasilica 21:411-419.
- Couté, A. and A. Iltis. 1984. Mise au point sur la flore peridiniale Algae, Pyrrhophyta d'eau douce de Côte d'Ivoire. Revue d'Hydrobiologie Tropicale 17:53-64.
- Johansen, J. R., R. L. Lowe, S. Carty, K. Fučíková, C. E. Olsen, M. H. Fitzpatrick, J. A. Ress and P. C. Furey. 2007. New algal species records for Great Smoky Mountains National Park, with an annotated checklist of all reported algal taxa for the park. Southeastern Naturalist, Special issue 1:99-134.
- Krakhmalny, A. F. 2011. New record of *Thompsodinium intermedium* (R. H. Thomps.) Bourr. (Dinophyta) in Ukraine. International Journal of Algae 13:117-123.
- Ngô, H. M. and L. A. Pfiester. 1990. Freshwater dinoflagellates from North Deming Pond, Minnesota, USA. Transactions of the American Microscopical Society 109:380-398.
- Pacheco, H. G. 1988. Florecimiento masivo de un dinoflagelado en el cenote Dzityá (Chen-Ha), Yucatán. Tesis, Escuela de Química, Universidad Autónoma de Yucatán. Mérida. 57 p.
- Popovský, J. 1970. Some thecate dinoflagellates from Cuba. Archiv für Protistenkunde 112:252-258.

Popovský, J. and L. Pfiester. 1990. Dinophyceae (Dinoflagellida). Band 6. Reprinted 2008 Süßwasserflora von Mitteleuropa. Spektrum Akademischer, Heidelberg. 272 p.

Sánchez-Molina, I., C. Zetina-Moguel, R. Medina-González and L. Pérez-Aranda. 1994. Phytoplankton composition in cenote Dzitya, Yucatán, Mexico. Abstracts of presentations,

North American Lake Management Society, Florida. Lake Reservoir Management 9:111.

Thompson, R. H. 1950. A new genus and new records of freshwater Pyrrhophyta in the Desmokontae and Dinophyceae. Lloyd 13:277-299.