



Available online at www.sciencedirect.com

Revista Mexicana de Biodiversidad

Revista Mexicana de Biodiversidad 88 (2017) 715–734



www.ib.unam.mx/revista/

Biogeography

Updated checklist and zoogeographic remarks of benthic amphipods (Crustacea: Peracarida: Amphipoda) of two coastal lagoons in the western Gulf of Mexico

*Lista de especies actualizada y notas zoogeográficas de anfípodos bentónicos (Crustacea:
Peracarida: Amphipoda) de dos lagunas costeras en el golfo de México occidental*

Andrea Raz-Guzmán*, Luis A. Soto

Laboratorio de Ecología del Bento, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Apartado postal 70-305, 04510
México City, Mexico

Received 6 November 2015; accepted 22 March 2017

Available online 12 August 2017

Abstract

Estuarine amphipods of the western Gulf of Mexico represent mostly eurythermal species. Sampling took place in Summer and Winter on seagrass beds, macroalgae and bare substrates, at 75 localities in Laguna Madre and 34 in Laguna de Tamiahua, Mexico, with a Renfro beam net and an otter trawl. A total of 19,398 specimens of 19 species were collected. *Cymadusa compta* was dominant in both lagoons with 63.4% of the total abundance. First records are 7 species for Laguna Madre and 11 species for Laguna de Tamiahua. A literature review updated the number of species from the 19 collected to 29. An analysis of the species' distribution produced 3 zoogeographic patterns: Warm and Caribbean, Warm Temperate and Caribbean, and temperate. A hierarchical cluster analysis of the geographic distribution data of the 29 species produced 2 assemblages: one with an affinity for the warmer regions of the Gulf of Mexico, Caribbean Sea and Brazil, and another associated with the temperate conditions of the northern Gulf of Mexico, the eastern USA, Canada and Argentina, each group indicating a preference for either higher or lower water temperatures.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Gammaridea; Corophiidea; Zoogeography; Laguna Madre; Laguna de Tamiahua; Mexico

Resumen

Los anfípodos estuarinos del golfo de México occidental representan en su mayoría especies euritermas. El muestreo se llevó a cabo en verano e invierno en pastos marinos, macroalgas y sustratos sin vegetación, en 75 localidades de laguna Madre y 34 de laguna de Tamiahua, México, con una red Renfro y una red camarnera. Se recolectaron 19,398 ejemplares de 19 especies. *Cymadusa compta* fue dominante en ambas lagunas con el 63.4% de la abundancia total. Se anotaron como primeros registros 7 especies en la laguna Madre y 11 en la laguna de Tamiahua. Se actualizó el número de especies de las 19 recolectadas a 29 mediante una revisión de la literatura. El análisis de la distribución de las especies proporcionó 3 patrones zoogeográficos: cálido-caribeño, cálido-templado-caribeño y templado. El análisis de conglomerados de la distribución geográfica de las 29 especies formó 2 grupos: uno con afinidad por las regiones más cálidas del golfo de México, Caribe y Brasil, y otro asociado a las condiciones templadas del norte del golfo de México, el este de EE. UU., Canadá y Argentina, indicando cada grupo una preferencia por temperaturas más cálidas o más frescas.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave: Gammaridea; Corophiidea; Zoogeografía; Laguna Madre; Laguna de Tamiahua; México

* Corresponding author.

E-mail address: razguzman@gmail.com (A. Raz-Guzmán).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

<http://dx.doi.org/10.1016/j.rmb.2017.07.002>

1870-3453/© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Amphipods, together with polychaetes and bivalves, make up 77% of the abundance in benthic communities of continental shelves and deep seas (Probert & Grove, 1998). They are a diverse, abundant and species-rich group of peracarid crustaceans. Most are free-living marine species, though some are freshwater, brackish or terrestrial. They are mostly eurythermal and euryhaline. They inhabit a wide variety of habitats and their geographic distribution is mostly cosmopolitan. They influence the structure of estuarine and coastal communities through their high abundance (Baldo, Arias, & Drake, 2001; Saunders, Attrill, Shaw, & Rowden, 2003; Tanner, 2006) and varied reproductive strategies and behaviour (Johnson, Stevens, & Watling, 2000; Thiel, 1997). They also affect community function by playing a part in the biogeochemical cycles of sedimentary organic matter (Gardner, Escobar, Cruz, & Rowe, 1993; Thiel & Hinojosa, 2009), promoting bioturbation and sediment stabilisation in soft substrates, as parasites of some vertebrates and vectors in the transmission of diseases and microorganisms, as well as through their feeding habits (DeBlois & Leggett, 1993; Duffy, 1990; Kamermans et al., 2002), nutrient recycling, linking foraging and detritivore food webs, and being prey of fish, birds and other crustaceans (Escobar & Soto, 1997; Hill & Elmgren, 1992; Zupo & Nelson, 1999). They are also excellent bioindicators of environmental quality (Montagna & Harper, 1996; Thomas, 1993b; Winfield & Ortiz, 2003).

Laguna Madre in Tamaulipas and Laguna de Tamiahua in Veracruz, Mexico, are large coastal lagoon systems that serve as nursery and feeding grounds for freshwater, estuarine and marine species. Their high biodiversity supports artisanal fisheries of oysters, shrimp, swimming crabs and mullet. Laguna Madre also provides feeding and refuge for migratory bird populations (Tunnell & Judd, 2002). The salinity gradients, submerged aquatic vegetation and different substrate types favour a variety of habitats that are available for resident and visiting species. In particular, the part that aquatic vegetation plays in the recruitment of many estuarine invertebrate and fish species has long been recognised (Heck & Crowder, 1991). High values of abundance have been recorded for species of invertebrates and fish that are temporal or permanent community components in these 2 lagoons (Raz-Guzmán & Barba, 2000). However, few studies have focused on the amphipods of these lagoons, and their zoogeography has not been reported. A literature review of studies on amphipods of the Mexican Gulf of Mexico coastal lagoons provided only the papers of Corona, Soto, and Sánchez (2000) and Cházaro, Winfield, Ortiz, and Álvarez (2002) for Laguna de Términos in Campeche, Winfield, Ortiz, Franco, and Bedia (1997), Winfield, Escobar, and Álvarez (2001), Winfield, Cházaro-Olvera, and Álvarez (2007), and Winfield and Ortiz (2011) for Laguna de Alvarado, Cházaro et al. (2002) and Winfield and Ortiz (2011) for Laguna Camaronera, and Winfield and Ortiz (2011) for Laguna de Son tecoman in Veracruz, and Barba and Sánchez (2005) and Ortega (2013) for Laguna Madre in Tamaulipas. The only study found for Laguna de Tamiahua was that of Winfield and Ortiz (2011).

The zoogeographic provinces along the Western Atlantic and the Gulf of Mexico have been described by Hedgpeth (1953), Briggs (1974), and Barnwell and Thurman (1984). Zoogeographic province boundaries are important barriers to the dispersal of tropical shallow marine species (Thurman, 1987). The boundaries along the Western Atlantic include the northern tropical-temperate boundary along an imaginary line that spans from Cabo Rojo in Veracruz to Tampa Bay and Cape Canaveral in Florida and the southern tropical-temperate boundary at Cabo Frío in Brazil. Laguna de Tamiahua is zoogeographically significant as its island barrier of Cabo Rojo lies at the transition between the Caribbean-tropical and the Carolinean-Temperate provinces in the western Gulf of Mexico (Thurman, 1987).

The recording of species along a geographical gradient provides information on the biogeographical and systematic structure of life, and the mapping of the geographic distribution of species contributes to understanding ecological and evolutionary patterns and processes. Cluster analyses assemble species with similar geographic localities, as well as geographic localities with a similar species composition, and support the establishment of zoogeographic distribution patterns (Craw, Grehan, & Heads, 1999). Considering that amphipod studies in Mexican coastal lagoons are few, the purpose of this study was to define the zoogeographic patterns of 29 amphipod species recorded for Laguna Madre and Laguna de Tamiahua, based on updated data of the species' geographic distribution that includes both published records for the Western Atlantic and the localities sampled in this study.

Materials and methods

Laguna Madre is the largest coastal lagoon in Mexico with an area of 200,000 ha. It lies at 23°45'–25°27' N, 97°23'–97°52' W. It is limited to the north by the delta of the Río Bravo and to the south by the Río Soto la Marina. In 1996, the sand barrier was breached by 5 inlets: Boca de Mezquital, Boca Ciega, Boca de Catán, Boca de Caballo and Boca Soto la Marina. As a result of its location in a semi-arid area, its limited communication with the sea and the reduced input of freshwater from the Río San Fernando, it is a hypersaline system with salinities of 33–62‰, water temperatures of 25–35 °C in Summer and 19–25 °C in Winter, and a maximum depth of 6 m (measured with a rope and dead weight). Further south, Laguna de Tamiahua is the third largest coastal lagoon in the Mexican Gulf of Mexico with 88,000 ha. It lies at 21°16'–22°05' N and 97°23'–97°43' W. It is limited to the north by the Río Pánuco and to the south by the Río Tuxpan, with the sandy barrier of Cabo Rojo to the east and 2 inlets, Tampachichi to the north and Corazones to the south. Six streams discharge into the lagoon along its western banks and contribute freshwater seasonally. For this reason, salinity varies from 22 to 38‰, water temperatures from 28 to 35 °C in Summer and 22 to 27 °C in Winter, and the maximum depth is 4 m (Fig. 1). Both lagoons have 2 marked climatic seasons, a dry Summer, and a rainy Winter when strong northerly winds (northerns) are common (Salcedo, Soto, Estradas-Romero, & Botello, 2017).

The submerged aquatic vegetation in both lagoons is characterised by the shoalgrass *Halodule wrightii* Aschers that settles




Figure 1. Laguna Madre, Tamaulipas and Laguna de Tamiahua, Veracruz, Mexico.

along the inner margins of the sand barriers, and macroalgae along the western margins of the lagoons that include the green algae *Cladophora* sp. and *Halimeda incrassata* (Ellis) Lamouroux syn. *H. tridens*, the brown algae *Dictyota dichotoma* (Hudson) Lamouroux, *Padina* sp. and *Rosenvingea intricata* (J. Agardh) Börgensen, and the red algae *Hypnea cervicornis* J. Agardh, *Gracilaria blodgettii* Harvey, *Gingicithara cylindrica* Börgensen, *Chondria littoralis* Harvey and *Digenea simplex* (Wulff) C. Agardh. Fringing mangroves in Laguna Madre are present only in the extreme south and are mostly *Avicennia germinans* (L.) L., while in Laguna de Tamiahua one finds small mangrove forests of *Rhizophora mangle* L., *A. germinans*, *Laguncularia racemosa* (L.) L. F. Gaertner and *Conocarpus erectus* L.

Specimens were collected in Summer (August) and Winter (December) of 1996 from 75 localities in August and 32 in December in Laguna Madre, and 34 localities in August and 23 in December in Laguna de Tamiahua (Fig. 1). Regarding the sampling date, it is important to make 3 considerations after having consulted the only 3 more recent studies available: 1) that only 3 species of 13 collected by Barba and Sánchez (2005) and 3 species of 12 collected by Ortega (2013) in Laguna Madre, as well as 5 species of 6 collected by Winfield and Ortiz (2011) in Laguna de Tamiahua, were not collected in our study; 2) that in our study we recorded 9 more species than Barba and Sánchez (2005), 11 more species than Ortega (2013) and 11 more species than Winfield and Ortiz (2011), and 3) that all the species recorded in these 3 papers plus the species collected in our study are included here, in view of which our species list may be considered to be taxonomically updated and valid.

Water temperature and salinity were recorded with a field thermometer and a refractometer respectively. Two nets were used to obtain specimens of different sizes and habitats: a Renfro beam net (1 mm mesh size, 50 m² sampling area) and an otter trawl (1 cm mesh size, 1.5–3 min capture per unit effort). The amphipods were preserved in 10% formalin in the field, after which they were transferred to a solution of glycerin and 70% alcohol for whitening previous to dissection.

The specimens were observed in detail in order to insure a correct identification of the species. The families are listed in alphabetical order. The assignment of genera to families follows Bousfield (1973), Martín and Díaz (2003) and LeCroy (2002, 2004, 2007). Exceptions are the Corophiidae that follow Myers and Lowry (2003) and the talitridan *stat. nov.* that follow Serejo (2004). The arrangement of genera and families included in the 2 latter works were adopted since they provide more recent phylogenetic evidence of relationships within the corophiid and talitridan amphipods. The genera and species were identified following Barnard (1969), Bousfield (1973), Myers (1982), Myers and McGrath (1984), Ledoyer (1986), Conlan (1990), Barnard and Karaman (1991), Thomas (1993a), LeCroy (2000, 2002, 2004, 2007), Myers and Lowry (2003), Appadoo and Myers (2004), and Serejo (2004). The species list follows Martin and Davis (2001) for the Suborder Gammaridea, Myers and Lowry (2003) for the Suborder Corophiidea, with an updating following Ahyong et al. (2011) and Horton, Lowry, and De Broyer (2013).

An Olmstead-Tükey analysis was applied to the abundance and spatial frequency data in order to identify dominant species. These and first records of species and families are reported for each lagoon, as well as the species collected in this study together

Table 1

Temperature, salinity, seasonal abundance (ind), number of species (spp.) and number of families (fam) of amphipods collected in each lagoon in this study.

	Laguna Madre	Laguna de Tamiahua
Temperature	19–35 °C	22–35 °C
Salinity	33–62‰	22–38‰
Abundance-species-families	13,052 ind – 18 spp. – 14 fam	6,346 ind – 12 spp. – 9 fam
Summer	5,367 ind – 14 spp.	4,209 ind – 10 spp.
Winter	7,685 ind – 13 spp.	2,137 ind – 11 spp.

with 3 species reported by Barba and Sánchez (2005) for Laguna Madre, 3 species reported by Ortega (2013) for the same lagoon (his *Protohyale* sp. B was not included in the analysis), and 5 species reported by Winfield and Ortiz (2011) for Laguna de Tamiahua. An updating was carried out in January 2015 for all species' names (<http://www.itis.gov>; Martin & Davis, 2001; Myers & Lowry, 2003) and geographic distributions. The specimens were kept in the Laboratorio de Ecología del Bentos, ICMyL, UNAM.

A map of the Western Atlantic including the important zoogeographic landmarks of Cape Cod, Cape Hatteras, Cape Canaveral and Tampa Bay in the United States, Cabo Rojo in Mexico, and Cabo Frío in Brazil, was prepared for each species considering the published data on the geographic distribution of each species (the geographic localities and the authors that reported them for each species are presented in Appendix 1), as well as the updated records for Laguna Madre and Laguna de Tamiahua. The Primer v.6 programme (Clarke & Gorley, 2006) was used to analyse a presence-absence matrix for species-geographic localities, and group geographic points according to Pearson's average linkage correlation to form clusters. A cluster analysis to group species provided no clear pattern.

Results

Laguna Madre and Laguna de Tamiahua experience marked seasonal changes in water temperature throughout the year. In Winter, values drop to 19 °C in Madre and to 22 °C in Tamiahua, while in Summer, there is a maximum of 35 °C in both lagoons. Salinity varied from 33 to 62‰ in Madre and from 22 to 38‰ in Tamiahua (Table 1).

A total of 19,398 specimens of 19 amphipod species were collected in the 2 lagoons. Laguna Madre provided 13,052 amphipods of 18 species and 14 families, of which 5,367 specimens and 14 species were collected in August and 7,685 specimens and 13 species in December. In the case of Laguna de Tamiahua, the amphipods collected included 6,346 specimens of 12 species and 9 families, with 4,209 specimens and 10 species recorded in August, and 2,137 specimens and 11 species in December (Table 1). Three species of the 13 recorded by Barba and Sánchez (2005), 3 species of the 12 recorded by Ortega (2013), and 5 species of the 6 recorded by Winfield and Ortiz (2011) that were not collected in the present study were added to the list, providing a total of 24 species for Laguna Madre and 17 species for Laguna de Tamiahua. In total, 29 species were included in the zoogeographic analysis (Table 2).

The amphipod species of Laguna Madre and Laguna de Tamiahua, together with the anomuran and brachyuran crabs of the 4 largest estuarine systems of the western and south-western Gulf of Mexico, contribute to the highly diversified fauna of the Tropical Western Atlantic (Raz-Guzmán & Sánchez, 1992a, 1992b, 1996, 1998; Guzmán, Sánchez, Soto, & Álvarez, 1986; Guzmán, Sánchez, & Soto, 1992; Raz-Guzmán, Sánchez, Peralta, & Florido, 2004). It is interesting to note that in spite of Cabo Rojo being the barrier island of Laguna de Tamiahua and representing a critical boundary for the northern dispersal of tropical shallow water fauna (Thurman, 1987), this lagoon and Laguna Madre further north share 12 of the 29 amphipod species recorded here, a 41% affinity that may indicate that these 2 lagoons constitute habitats of positive population growth from which amphipods can successfully disperse to neighbouring areas.

Cymadusa compta (S.I. Smith, 1873) was dominant in both lagoons with 63.4% of the overall total abundance. Of the 18 species collected in this study in Laguna Madre, 2 constitute 82.3% of the total abundance: *C. compta* (61.7%) and *Elasmopus levis* (S.I. Smith, 1873) (20.6%), and are followed by *Batea catharinensis* F. Müller, 1865 (6.1%), *Ericthonius brasiliensis* (Dana, 1853) (5.2%), *Hourstonius laguna* (McKinney, 1978) (4.5%), and the other 13 species with 0.008–0.912%. Of the 12 species collected in Laguna de Tamiahua, *C. compta* is markedly more abundant than the others with 66.8%, and is followed by *Gammarus mucronatus* Say, 1818 (13.4%), *Apohyale prevostii* (H. Milne Edwards, 1830) (12.5%), *E. brasiliensis* (3.9%), and the other 8 species with 0.016–1.261% (Table 3).

Seven species reported here for Laguna Madre were not recorded by Barba and Sánchez (2005) or Ortega (2013). *Ampelisca vadorum* Mills, 1963, *H. laguna*, *Ampithoe longimana* S.I. Smith, 1873, *Ampithoe valida* S.I. Smith, 1873, *Melita nitida* S.I. Smith, 1873, *Eobrolgus spinosus* (Holmes, 1905) and *Orchestia grillus* (Bosc, 1802) may thus be considered first records for this lagoon, as well as 2 families, the Amphilochidae and Phoxocephalidae. In the case of Laguna de Tamiahua, with the exception of *G. mucronatus* which was recorded previously by Winfield and Ortiz (2011), the other 11 species, *A. vadorum*, *H. laguna*, *A. longimana*, *A. valida*, *C. compta*, *Monocorophium acherusicum* (Costa, 1851), *Nototropis minikoi* (A.O. Walker, 1905), *A. prevostii*, *E. brasiliensis*, *E. levis* and *M. nitida*, are first records, together with the family Amphilochidae.

The mapping of the geographic distribution data of each of the 29 species recorded here made it possible to visually define 3 zoogeographic distribution patterns along the Western Atlantic from Newfoundland to Argentina: Warm and Caribbean

Table 2

Amphipod species and abundance in Laguna Madre (LM) and Laguna de Tamiahua (LT) (+ Barba & Sánchez, 2005; ++ Winfield & Ortiz, 2011; +++ Ortega, 2013).

Family	Species	LM	LT
Subphylum Crustacea Brünich, 1772			
Class Malacostraca Latreille, 1802			
Subclass Eumalacostraca Grobben, 1892			
Superorder Peracarida Calman, 1904			
Order Amphipoda Latreille, 1816			
Suborder Gammaridea Latreille, 1802 (sensu Martin & Davis, 2001)			
Ampeliscidae Krøyer, 1842	<i>Ampelisca vadorum</i> Mills, 1963	2	2
	<i>Ampelisca venetiensis</i> Shoemaker, 1916	++	
	<i>Ampelisca verrilli</i> Mills, 1967	++	
Amphilochidae Boeck, 1871	<i>Hourstonius laguna</i> (McKinney, 1978)	589	7
Bateidae Stebbing, 1906	<i>Batea catharinensis</i> F. Müller, 1865	794	
Dexaminidae Leach, 1814	<i>Nototropis minikoi</i> (A.O. Walker, 1905)	119	2
Gammaridae Leach, 1814	<i>Gammarus mucronatus</i> Say, 1818	42	847
Hyalidae Bulycheva, 1957	<i>Apohyale prevostii</i> (H. Milne Edwards, 1830)	5	791
Melitidae Bousfield, 1983	<i>Parhyale hawaiensis</i> (Dana, 1853)	+++	
	<i>Elasmopus levius</i> (S.I. Smith, 1873)	2684	80
	<i>Elasmopus pectenircus</i> (Bate, 1862)	+++	
	<i>Elasmopus rapax</i> Costa, 1853	+++	
	<i>Melita nitida</i> S.I. Smith, 1873	1	54
Phoxocephalidae G.O. Sars, 1891	<i>Eobrolgus spinosus</i> (Holmes, 1905)	1	++
Talitridae Rafinesque, 1815	<i>Orchestia gammarellus</i> (Pallas, 1766)	+	
	<i>Orchestia grillus</i> (Bosc, 1802)	13	
Suborder Corophiidea Leach, 1814 (sensu Myers & Lowry, 2003)			
Infraorder Corophiida Leach, 1814			
Superfamily Aoroidea Stebbing, 1899			
Aoridae Stebbing, 1899	<i>Grandidierella bonnieroides</i> Stephensen, 1948	55	
Unciolidae Myers & Lowry, 2003	<i>Lembos websteri</i> Bate, 1857	+	
	<i>Unciola serrata</i> Shoemaker, 1945		++
Superfamily Corophioidea Leach, 1814			
Ampithoidae Stebbing, 1899	<i>Ampithoe longimana</i> S.I. Smith, 1873	10	1
	<i>Ampithoe valida</i> S.I. Smith, 1873	4	76
Corophiidae Leach, 1814	<i>Cymadusa compta</i> (S.I. Smith, 1873)	8054	4240
	<i>Monocorophium acherusicum</i> (Costa, 1851)		1
	<i>Monocorophium tuberculatum</i> (Shoemaker, 1934)	202	
Infraorder Caprellida Leach, 1814			
Superfamily Microprotozoidea Myers & Lowry, 2003			
Microprotopidae Myers & Lowry, 2003	<i>Microprotopus raneyi</i> Wigley, 1966	1	
Superfamily Photoidea Boeck, 1871			
Ischyroceridae Stebbing, 1899	<i>Cerapus tubularis</i> Say, 1817	1	
	<i>Ericthonius brasiliensis</i> (Dana, 1853)	674	245
Photidae Boeck 1871	<i>Jassa falcata</i> (Montagu, 1808)	+	
	<i>Photis longicaudata</i> (Bate & Westwood, 1862)		++

Table 3

Dominant amphipod species in Laguna Madre (LM) and Laguna de Tamiahua (LT) (% of total abundance).

Species	Laguna Madre	Laguna de Tamiahua
<i>Cymadusa compta</i>	61.7	66.8
<i>Elasmopus levius</i>	20.6	
<i>Gammarus mucronatus</i>		13.4
<i>Apohyale prevostii</i>		12.5
<i>Batea catharinensis</i>	6.1	
<i>Ericthonius brasiliensis</i>	5.2	3.9
<i>Hourstonius laguna</i>	4.5	
The other 13 species in LM	0.008–0.912	
The other 8 species in LT		0.016–1.261




Figure 2. Distribution of Warm and Caribbean species (*Ampelisca venetiensis*, *Hourstonius laguna*, *Parhyale hawaiensis*, *Photis longicaudata*, *Elasmopus pectenircus*) along the Western Atlantic (■) including the Mexican lagoons (●).

with 5 species (17%), Warm Temperate and Caribbean with 18 species (62%) and Temperate with 6 species (21%). A hierarchical cluster analysis of the presence-absence matrix of geographic distribution data of the 29 amphipod species produced 2 groups at a correlation level of -0.05 (with other smaller groups at higher correlation levels). One group included the warmer regions of the Gulf of Mexico, Caribbean Sea and Brazil (+), and another group included the colder regions of the northern Gulf of Mexico, the eastern and northeastern United States, Canada, Greenland, and Argentina (●).

Discussion

The first records reported here for Laguna Madre (*A. vadourum*, *H. laguna*, *A. longimana*, *A. valida*, *M. nitida*, *E. spinosus* and *O. grillus*) and Laguna de Tamiahua (*A. vadourum*, *H. laguna*, *A. longimana*, *A. valida*, *C. compta*, *M. acherusicum*, *N. minikoi*, *A. prevostii*, *E. brasiliensis*, *E. levis* and *M. nitida*) respond to the fact that previous studies on amphipods in the state of Veracruz have focused on Laguna Camaronera (Cházaro et al., 2002; Winfield & Ortiz, 2011), Laguna de Alvarado (Winfield & Ortiz, 2011; Winfield et al., 1997, 2001; Winfield, Cházaro-Olvera, et al., 2007), Laguna de Sontecomapan (Winfield & Ortiz, 2011), the Veracruz Reef System (Winfield & Ortiz, 2011; Winfield, Abarca-Arenas, & Cházaro-Olvera, 2007; Winfield, Cházaro-Olvera, Horta-Puga, Lozano-Aburto, & Arenas-Fuentes, 2010; Winfield, Cházaro-Olvera, Ortiz, & Palomo-Aguayo, 2011; Winfield, 2013) and the Veracruz continental shelf (Winfield & Ortiz, 2011), with only that of Winfield and Ortiz (2011) for

Laguna de Tamiahua, while only those of Barba and Sánchez (2005) and Ortega (2013) have dealt with Laguna Madre in the state of Tamaulipas.

The 5 Warm and Caribbean species, *Ampelisca venetiensis* Shoemaker, 1916, *H. laguna*, *Parhyale hawaiensis* (Dana, 1853), *Photis longicaudata* (Bate & Westwood, 1862) and *Elasmopus pectenircus* (Bate, 1862), are mostly distributed along the eastern coast of Florida and the Gulf of Mexico (except for *A. venetiensis*), throughout the Caribbean Sea, and along the coast of Brazil (except for *A. venetiensis* and *H. laguna*). These species include a few records along the Temperate northern Gulf of Mexico which may be explained as the result of dispersal by the shallow coastal currents that flow from Louisiana and Texas (the LATEX current) to Campeche during the Winter and back during the Summer (Fernández, Zavala, & Romero, 1998) (Fig. 2). The capacity to disperse is important in the life history of many species. Benthic invertebrates usually disperse through a swimming larval phase. However, many species with a direct development have wide geographic distributions (Castilla & Guílez, 2000; Thiel, 2003a). Floating and rafting have been identified as highly efficient dispersal mechanisms in these species (Highsmith, 1985; Thiel, 2003b), which include shallow-water isopods and amphipods (Locke & Corey, 1989), many of which brood their young. This has been supported by Havermans, De Broyer, Mallefet, and Zintzen (2007) who recorded how *Jassa herdmani* (Walker, 1893), a tube-building amphipod, may travel long distances between hard-substrate areas, and by records of brooding benthic peracarids found in plankton nets in areas far from their native habitats (Williams




Figure 3. Distribution of Warm Temperate and Caribbean species. Group A1 (*Unciola serrata*, *Ampelisca verrilli*, *Cymadusa compta*, *Elasmopus levis*, *Eobrolgus spinosus*, *Microprotopus raneyi*) along the Western Atlantic (■) including the Mexican lagoons (●).




Figure 4. Distribution of Warm Temperate and Caribbean species. Group A2 (*Gammarus mucronatus*, *Ampelisca vadorum*, *Ampithoe longimana*, *Melita nitida*) along the Western Atlantic (■) including the Mexican lagoons (●).




Figure 5. Distribution of Warm Temperate and Caribbean species. Group B1 (*Grandidierella bonnieroides*, *Nototropis minikoi*, *Elasmopus rapax*) along the Western Atlantic (■) including the Mexican lagoons (●).

& Bynum, 1972). Most common floaters are small individuals that may remain suspended on the water surface for long periods of time. Small animals are also more likely to live on seagrass and macroalgae, and may become rafters when the plants break (Norton & Mathieson, 1983). However, gravid females have also been observed rendering rafting a potential dispersal mechanism. It is important to note that *P. hawaiensis*, a species used in developmental and genetic studies, may actually occur as a species complex (Myers, 1985).

The warm affinity of these species is supported by records from other warm localities that include Hawaii, the Philippines and the South China Sea for *P. hawaiensis*, and Hawaii, the Red Sea, Madagascar, Mauritius, India, Indonesia, the South China Sea, New Guinea and Australia for *E. pecteniferus*. *P. longicaudata* has been recorded in the Indian Ocean, however it is also present in colder localities including Sweden, Ireland, Scotland, England, Belgium, France, Spain, Greece, the Mediterranean Sea, the Black Sea and Japan (see Appendix 1).

The 18 Warm Temperate and Caribbean species may be grouped into 5 sub-groups. Six species in group A1, *Unciola serrata* Shoemaker, 1945, *Ampelisca verrilli* Mills, 1967, *C. compta*, *E. levis*, *E. spinosus* and *Microprotopus raneyi* Wigley, 1966, present a mostly continuous distribution from Maine, south along the eastern coast of the United States, the Gulf of Mexico and throughout the Caribbean Sea (except for *U. serrata* which at present has no records in the Caribbean) (Fig. 3). The species in group A2 follow the same distribution but include temperate habitats in Nova Scotia and a northernmost point in the Gulf of St Lawrence, Canada. These are *G. mucronatus*, *A. vadorum*, *A. longimana* and *M. nitida*. It is interesting to note that *G. mucronatus* has 1 single record in the Caribbean, in Quintana Roo, Mexico (Fig. 4).

The species in group B1 are distributed along the southern half of the eastern coast of the United States, the Gulf of Mexico, the Caribbean Sea and the coast of Brazil, and include *Grandidierella bonnieroides* Stephensen, 1948, *N. minikoi* and




Figure 6. Distribution of Warm Temperate and Caribbean species. Group B2 (*Ericthonius brasiliensis*, *Monocorophium acherusicum*, *Batea catharinensis*) and group C (*Ampithoe valida*, *Cerapus tubularis*) along the Western Atlantic (■) including the Mexican lagoons (●).

Elasmopus rapax Costa, 1853 (Fig. 5). In particular, regarding *G. bonnieroides*, Myers (1970) reviewed its status and stated that specimens collected in the Gulf of Mexico, Caribbean Sea, Colombia, Venezuela, and Trinidad corresponded to this species. Asari and Myers (1982) later observed variations in specimens collected in the Western Atlantic and concluded that a complex of species was present. However, the 55 specimens from Laguna Madre were identified by A. Corona (pers comm), an expert in amphipod taxonomy, leaving no doubt as to the identity of the species. Three of the Warm Temperate and Caribbean species, *C. compta*, *A. longimana* and *E. rapax*, have also been recorded in Bermuda. The species in group B2, *E. brasiliensis*, *M. acherusicum* and *B. catharinensis*, have the same distribution but extend northwards to Nova Scotia (Fig. 6). The coastline between Venezuela and Brazil with no records in the case of some species most probably indicates a lack of studies throughout the area rather than a disjunct distribution of the species.

The species in group C, *A. valida* and *Cerapus tubularis* Say, 1817, are distributed along the eastern coast of the United States from Maine to Florida, throughout the Gulf of Mexico, the Caribbean Sea, southern Brazil (*C. tubularis*) and include some southernmost points in Argentina (Fig. 6).

The warm and temperate affinity of these species is reflected in the wide range of geographic locations where they have been

recorded and indicates a high ecologic plasticity, as Melo (1990) stated for South American brachyurans. The geographic locations previously recorded for these species, which latitudinally include the Gulf of Mexico, validate their presence in this area. As an example of this, 12 species reported here are amphiamerican or cosmopolitan: *E. spinosus* (Washington to California, USA; Baja California and Gulf of California, Mexico), *G. mucronatus* (California, USA), *A. longimana* (Baja California and Gulf of California, Mexico), *M. nitida* (British Columbia-Canada, Washington to California, USA, Baja California, Sinaloa and Michoacán, Mexico, Costa Rica, Panama, Ecuador, United Kingdom, Netherlands, Germany), *G. bonnieroides* (Tanzania, Mozambique, South Africa, Madagascar, Mauritius, India, Australia), *N. minikoi* (East India, South China Sea, Indo-Pacific), *E. rapax* (United Kingdom, Italy, Mediterranean Sea, Turkey, Red Sea, Australia), *E. brasiliensis* (California, USA, Baja California, Mexico, Norway, United Kingdom, France, Spain, Italy, Mediterranean Sea, Mauritius, Indian Ocean, South China Sea), *M. acherusicum* (Alaska to California, USA, Baja California, Mexico, Hawaii, United Kingdom, Netherlands, Mediterranean Sea, France, Italy, Egypt, Tunisia, Algeria, Senegal, Cape Colony, Dar es Salaam, Mauritius, Indian Ocean, Japan, South China Sea, Australia, New Zealand), *B. catharinensis* (Baja California, Mexico), *A. valida* (British Columbia, Canada to California, USA, Japan, Korea), and *C. tubularis*




Figure 7. Distribution of temperate species (*Lembos websteri*, *Monocorophium tuberculatum*, *Orchestia grillus*, *Apohyale prevostii*, *Jassa falcata*, *Orchestia gammarellus*) along the Western Atlantic (■) including the Mexican lagoons (●).

(California, USA to Baja California, Mexico, Japan, Korea) (Appendix 1).

Among the 6 temperate species (Fig. 7), *Lembos websteri* Bate, 1857, *Monocorophium tuberculatum* (Shoemaker, 1934)

and *O. grillus* are distributed from Maine to eastern Florida and in the Gulf of Mexico. *L. websteri* also has records in an extremely northerly point in Greenland, while *O. grillus* extends northwards to the Newfoundland coast and southwards




Figure 8. Cluster diagram of geographic points recorded for the 29 amphipod species of Laguna Madre, Tamaulipas and Laguna de Tamiahua, Veracruz, Mexico, showing one group of warmer regions and another group of colder regions. Eight smaller groups may be defined as: A, eastern and northeastern United States; B, Argentina; C, northeastern United States, Canada and Argentina (and ? Panama); D, northeastern Brazil; E, Caribbean; F, northwestern Gulf of Mexico; G, western, southern and eastern Gulf of Mexico, and H, Caribbean. Three interrogation marks (?) indicate odd localities within groups.

to Colombia (this last point is considered an isolated record). *M. tuberculatum* is included in this group in spite of having 1 record in Florida Bay, in consideration of its clearly northern affinity and its absence from typically tropical habitats such as Laguna de Términos, Campeche where Ledoyer (1986) and Corona et al. (2000) did not record it in spite of thorough sampling. These species present a disjunct distribution at the tip of Florida where ocean water masses are more characteristically tropical, as is the case of the Carolinian zoogeographic province (Thurman, 1987).

The other 3 temperate species, *A. prevostii*, *Jassa falcata* (Montagu, 1808) and *Orchestia gammarellus* (Pallas, 1766), also present a northerly distribution up to Newfoundland. *A. prevostii* extends south to Laguna de Tamiahua, Veracruz (with 791 specimens collected in this study) at the northern tropical-temperate boundary for shallow water fauna within the western Gulf of Mexico, and includes an isolated record in eastern Brazil. *J. falcata* and *O. gammarellus* also present a southern temperate distribution with records in southern Brazil (*J. falcata*), an isolated point in eastern Brazil (*O. gammarellus*) and Argentina (both species). The presence of species in widely separated localities such as these generally responds to dispersal processes governed by ocean currents, rafting on plant debris, logs, sargassum and macroalgae, and transportation in ships' ballast water (Appadoo & Steele, 1998).

L. websteri, *M. tuberculatum*, *O. grillus* and *J. falcata*, previously recorded in the northern Gulf of Mexico, may disperse west and south to the northern tropical-Temperate boundary at Cabo Rojo, Veracruz, by floating and/or rafting on the LATEX coastal current, as is discussed above in the section on the "Warm and Caribbean species" (Castilla & Guílez, 2000; Fernández et al., 1998; Havermans et al., 2007; Highsmith, 1985; Locke & Corey, 1989; Norton & Mathieson, 1983; Thiel, 2003a, 2003b; Williams & Bynum, 1972). The 6 temperate species were recorded in Laguna Madre, Tamaulipas, which lies north of this zoogeographic boundary.

As a confirmation of the temperate affinity of these species, other temperate localities have been reported for 4 of these species: for *L. websteri* Norway, England, France, Spain, Italy, Yugoslavia, Greece, Turkey, Israel, Egypt, the Mediterranean Sea and West Africa, for *A. prevostii* the Arctic Ocean, Iceland, Ireland, Norway, the United Kingdom, the Faroe Islands, the Netherlands, Belgium, France and Spain, for *J. falcata* Ireland, Norway, the United Kingdom, Spain, the Mediterranean Sea, British Columbia, Canada to California-United States and New Zealand, and for *O. gammarellus* Iceland, Norway, the United Kingdom, Spain, the Mediterranean Sea, the Black Sea, Turkey and SW Africa (Appendix 1).

The area with no records that may be seen for some species along the Central American coasts and the northern and northeastern South American coasts may, in general, be attributed to a lack of local studies. Future studies in these areas should thus contribute to filling the information gaps, particularly in the case of species in the Warm Temperate and Caribbean distribution pattern.

The grouping of warmer (+) and colder (●) geographic localities generated by the hierarchical cluster analysis based on the




Figure 9. Grouping of the warmer (+) and colder (●) regions along the Western Atlantic based on the geographic points recorded for the 29 amphipod species of Laguna Madre, Tamaulipas and Laguna de Tamiahua, Veracruz, Mexico. The 2 slotted lines represent the zoogeographic boundaries that separate the warmer tropical regions from the colder Temperate regions along the Western Atlantic. The northern boundary spans along an imaginary line from Cabo Rojo in Veracruz to Tampa Bay and Cape Canaveral in Florida, and the southern boundary lies at Cabo Frio in Brazil.

presence of gammarid and corophiid amphipods along the Western Atlantic (Figs. 8 and 9) indicates that these species, despite most being eurythermal, have a physiological preference for higher or lower water temperatures (Barba & Sánchez, 2005; d'Udekem-d'Acoz, & Verheyen, 2013).

This is the first comprehensive study of gammarid and corophiid amphipods of 2 major coastal lagoons in the western Gulf of Mexico that are characterised by their unique environmental conditions and are located at the northern tropical-temperate zoogeographical boundary that restricts the dispersion of tropical shallow water fauna. It may be stated that these estuarine amphipods may be further analysed, not only due to their abundance and diversity but because some species

may be considered good indicators of zoogeographic provinces along the Western Atlantic.

Acknowledgements

To A. Corona for sample processing, identification of species and the note on *G. bonnieroides*; N. López for identifying the macroalgae; L. Huidobro and D. Salcedo for running the cluster analyses; M. Thiel for reviewing an earlier version of the manuscript, and especially I. Winfield for reviewing the final draft. Conabio provided the funds to carry out this research (project H258).

Appendix 1. Geographic distribution and references per species (+ Barba & Sánchez, 2005; ++ Winfield & Ortiz, 2011; +++ Ortega, 2013). Lesser Antilles: St. Thomas, Tortola, Anegada, Jost van Dyke, Virgin Gorda, St. Croix, Anguilla, St. Martin, St. Barthélemy, Saba, St. Eustatius, Barbuda, St. Kitts, Nevis, Antigua, Montserrat, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Grenadines, Grenada. References:

A = www.geocities.com/ajuarroero/amphipods.html
B = www.bdt.fat.org.br/workshop/costa/praias/tabela3
C = www.acd.ofrj.br/mndi/Carcinologia/hp/Text/Amphipoda1.htm

Geographic distribution	References
<i>Ampelisca venetiensis</i> ++ Figure 2 Laguna de Tamiahua in Veracruz, Mexico; Costa Rica; Caribbean Sea; Colombia.	Warm and Caribbean Escobar, Winfield, Ortiz, Gasca, & Suárez, 2002; Escobar & Winfield, 2003; Ortiz, Martín, & Díaz, 2007; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013
<i>Hourstonius laguna</i> Figure 2 St John's River, Indian River Lagoon, Biscayne Bay, Florida Bay in Florida, Mississippi, and Galveston, San Antonio Bay, Corpus Christi Bay, Laguna Madre in Texas, US; Laguna Madre in Tamaulipas, Laguna de Tamiahua, Laguna Camaronera and Laguna de Alvarado in Veracruz, Laguna de Términos in Campeche, Quintana Roo, Mexico; Bermuda; Caribbean Sea; Cuba; Colombia; Venezuela. – California, US.	McKinney, 1978; Ledoyer, 1986; Barnard & Karaman, 1991; Ortiz & Lalana, 1993; Nelson, 1995; Camp, Lyons, & Perkins, 1998; Corona et al., 2000; Cházaro et al., 2002; LeCroy, 2002; Ortiz et al., 2007; Winfield, Cházaro-Olvera, et al., 2007; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013
<i>Parhyale hawaiensis</i> +++ Figure 2 Florida, Louisiana, US; Gulf of Mexico; Soto la Marina in Tamaulipas, Quintana Roo, Mexico; Caribbean Sea; Cuba; Haiti; Dominican Republic; Puerto Rico; Venezuela; Aruba; Curaçao; Bonaire; Lesser Antilles; Rio de Janeiro to Paraná, Brazil. – Circumtropical; California, US; Atlantic and Pacific oceans; Hawaii; Fiji; Philippines; South China Sea; Mediterranean Sea.	Shoemaker, 1956; Thomas, 1976; Myers, 1985; Camp et al., 1998; Wakabara & Serejo, 1998; Serejo, 1999; Lowry, 2000; Ortiz et al., 2007; Somaio & Moreira, 2008; LeCroy, Gasca, Winfield, Ortiz, & Escobar-Briones, 2009; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Ortega, 2013
<i>Photis longicaudata</i> ++ Figure 2 Florida, US; Laguna de Tamiahua in Veracruz, Quintana Roo, Mexico; Caribbean Sea; Venezuela; Pernambuco, Alagoas, Espírito Santo, Rio de Janeiro, São Paulo, Rio Grande do Sul, Brazil. – Atlantic and Indian oceans; Sweden; Ireland; Scotland; England; Belgium; France; Spain; Greece; Mediterranean Sea; Black Sea; Japan.	Wakabara, Tararam, Valério-Berardo, Duleba, & Pereira, 1991; Camp et al., 1998; Wakabara & Serejo, 1998; Escobar et al., 2002; Ortiz et al., 2007; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013; www.marinespecies.org
<i>Elasmopus pecteniferus</i> +++ Figure 2 Florida, Florida Keys, US; Gulf of Mexico; Laguna Madre in Tamaulipas, Quintana Roo, Mexico; Caribbean Sea; Cuba; Lesser Antilles; west, central and east coasts of Venezuela; Ceará to Paraná, Brazil. – Circumtropical; Atlantic, Pacific and Indian oceans; Hawaii; Mediterranean Sea; Turkey; Red Sea; western and southern Africa; Madagascar; Mauritius; India; Indonesia; South China Sea; New Guinea; Australia.	Barnard, 1970; McKinney, 1977; Ortiz, 1978; Karaman, 1982; Wakabara, Tararam, & Takeda, 1983; Hay, Duffy, Fenical, & Gustafson, 1988; Appadoo & Steele, 1998; Camp et al., 1998; Wakabara & Serejo, 1998; Lowry, 2000; Díaz & Martín, 2001; Escobar et al., 2002; Appadoo & Myers, 2003; Martín & Díaz, 2003; Ortiz et al., 2007; LeCroy et al., 2009; Bakir, Sezgin, & Katagan, 2010; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Martín et al., 2013; Ortega, 2013; Ortega & Martín, 2013
<i>Unciola serrata</i> ++ Figure 3 Cape Cod in Massachusetts to Delaware Bay and Virginia, Florida, US; Laguna de Tamiahua in Veracruz, Mexico.	Warm Temperate and Caribbean Dickinson, Wigley, Brodeur, & Brown-Leger, 1980; Camp et al., 1998; Winfield & Ortiz, 2011; www.marinespecies.org
<i>Ampelisca verrilli</i> ++ Figure 3 Cape Cod in Massachusetts to Delaware Bay and Chesapeake Bay, Virginia to Cape Hatteras in North Carolina, Indian River Lagoon in Florida, US; Laguna de Tamiahua and Veracruz reef system in Veracruz, Campeche, Quintana Roo, Mexico; Caribbean Sea.	Dickinson et al., 1980; Nelson, 1995; Camp et al., 1998; Eggleston, Elis, Etherington, Dahlgren, & Posey, 1999; Escobar et al., 2002; Escobar & Winfield, 2003; Winfield, Escobar, & Morrone, 2006; Ortiz et al., 2007; Winfield & Escobar, 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013; www.marinespecies.org

Geographic distribution	References
<i>Cymadusa compta</i> Figure 3	Watling & Maurer, 1972; Bousfield, 1973; Ledoyer, 1986; Ortiz & Lalana, 1993; Thomas, 1993a; Nelson, 1995; Camp et al., 1998; Eggleston et al., 1999; Corona et al., 2000; Cházaro et al., 2002; LeCroy, 2002; Varela, Manuel, & Rogelio, 2003; Barba & Sánchez, 2005; Ortiz et al., 2007; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Martín et al., 2013; Ortega, 2013
<i>Elasmopus levis</i> Figure 3	Watling & Maurer, 1972; Bousfield, 1973; McKinney, 1977; Dickinson et al., 1980; Ledoyer, 1986; Ortiz & Lalana, 1993; Camp et al., 1998; Eggleston et al., 1999; Corona et al., 2000; LeCroy, 2000; Escobar et al., 2002; Barba & Sánchez, 2005; Ortiz et al., 2007; Winfield, Abarca-Arenas, et al., 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013; Ortega, 2013; Paz-Ríos & Ardisson, 2013
<i>Erobolus spinosus</i> Figure 3	Barnard, 1970; Watling & Maurer, 1972; Bousfield, 1973; Barnard & Barnard, 1981; Ledoyer, 1986; Camp et al., 1998; Eggleston et al., 1999; Corona et al., 2000; Díaz & Martín, 2001; Martín, Ortiz, & Díaz, 2002; Escobar & Winfield, 2003; Martín & Díaz, 2003; Winfield et al., 2006; Ortiz et al., 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Ortiz & Lalana, 2010; LeCroy, 2011; Winfield & Ortiz, 2011; Martín et al., 2013; Paz-Ríos, Simões, & Ardisson, 2013
<i>Micropotopus raneyi</i> Figure 3	Lowry, 1972; Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Barnard & Karaman, 1991; Rakocinski, Heard, LeCroy, McLelland, & Simons, 1993; Camp et al., 1998; Eggleston et al., 1999; Díaz & Martín, 2001; Barba & Sánchez, 2005; Ortiz et al., 2007; Dubois, Gelpi, Condrey, Grippo, & Fleeger, 2009; Miloslavich et al., 2010; Martín et al., 2013; Ortega & Martín, 2013
<i>Gammarus mucronatus</i> Figure 4	Watling & Maurer, 1972; Bousfield, 1973; Thomas, 1976; Lincoln, 1979; Dickinson et al., 1980; Heard, 1982; Ledoyer, 1986; Ortiz & Lalana, 1993; Nelson, 1995; Winfield et al., 1997; Camp et al., 1998; Eggleston et al., 1999; Cházaro et al., 2002; Escobar & Winfield, 2003; Barba & Sánchez, 2005; Ortiz et al., 2007; Winfield, Cházaro-Olvera, et al., 2007; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013; Ortega, 2013
<i>Ampelisca vadorum</i> Figure 4	Mills, 1963; Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Ledoyer, 1986; Barnard & Karaman, 1991; Ortiz & Lalana, 1993; Nelson, 1995; Camp et al., 1998; Eggleston et al., 1999; Corona et al., 2000; LeCroy, 2002; Escobar & Winfield, 2003; Winfield et al., 2006; Ortiz et al., 2007; Winfield & Escobar, 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Martín et al., 2013; Paz-Ríos & Ardisson, 2013
<i>SW Gulf of St Lawrence, Nova Scotia, Canada; Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina to Florida and Florida Keys, Louisiana, Laguna Madre in Texas, US; northern Gulf of Mexico; Laguna Madre in Tamaulipas, Mexico; Caribbean Sea; Colombia; Venezuela.</i>	
<i>– California, US.</i>	
<i>SW Gulf of St Lawrence, Nova Scotia, Canada; Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina, South Carolina, Georgia, Florida and Florida Keys, US; Laguna Madre in Tamaulipas, Laguna de Tamiahua and continental shelf off Alvarado in Veracruz, southwestern Gulf of Mexico, Laguna de Términos and Bay of Campeche in Campeche, Yucatán, Quintana Roo, Mexico; Caribbean Sea; Cuba.</i>	

Geographic distribution	References
<i>Ampithoe longimana</i> Figure 4	Barnard, 1963; Bousfield, 1973; Ortiz & Lalana, 1993; Nelson, 1995; Camp et al., 1998; Eggleston et al., 1999; LeCroy, 2002; Martín & Díaz, 2003; Ortiz et al., 2007; Miloslavich et al., 2010; Winfield & Ortiz, 2011; Winfield et al., 2011; Martín et al., 2013; Winfield, 2013
<i>Melita nitida</i> Figure 4	Watling & Maurer, 1972; Bousfield, 1973; Thomas, 1976; Lincoln, 1979; Dickinson et al., 1980; Heard, 1982; Ortiz, 1983; Chapman, 1988; Ortiz & Lalana, 1993; Nelson, 1995; Camp et al., 1998; Eggleston et al., 1999; Corona et al., 2000; LeCroy, 2000; Corona & Raz-Guzmán, 2003; Faasse & van Moorsel, 2003; Ortiz et al., 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Winfield et al., 2011; Martín et al., 2013; Winfield, 2013; A
<i>Grandidierella bonnieroides</i> Figure 5	Fox & Bynum, 1975; Thomas, 1976; Leite, Tararam, & Wakabara, 1980; Asari & Myers, 1982; Heard, 1982; Ledoyer, 1986; Wakabara et al., 1991; Ortiz & Lalana, 1993; Nelson, 1995; Winfield et al., 1997; Camp et al., 1998; Wakabara & Serejo, 1998; Corona et al., 2000; Ortiz, 2001; Winfield et al., 2001; Cházaro et al., 2002; LeCroy, 2002; Martín & Díaz, 2003; Appadoo & Myers, 2004; Ortiz et al., 2007; Winfield, Cházaro-Olvera, et al., 2007; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Winfield & Ortiz, 2011; Martín et al., 2013; Ortega, 2013; Paz-Ríos & Ardisson, 2013; A, B, C
<i>Nototropis minikoi</i> Figure 5	Stephensen, 1948; Oliveira, 1953; Bousfield, 1973; Bynum & Fox, 1977; Naomi, 1979; Leite et al., 1980; Ledoyer, 1986; Lagarde, 1987; Barnard & Karaman, 1991; Wakabara et al., 1991; Ortiz & Lalana, 1993; Wakabara, Nicoletti, & Tararam, 1996; Wakabara & Serejo, 1998; Lowry, 2000; Cházaro et al., 2002; Lowry & Stoddart, 2003; Martín & Díaz, 2003; Barba & Sánchez, 2005; McLaughlin et al., 2005; Ortiz et al., 2007; LeCroy et al., 2009; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Winfield & Ortiz, 2011; Martín et al., 2013; Ortega & Martín, 2013; Paz-Ríos et al., 2013; A, B, C
<i>Elasmopus rapax</i> +++ Figure 5	Shoemaker, 1933; Barnard, 1962; Bousfield, 1973; McKinney, 1977; Ortiz, 1978; Lincoln, 1979; Ortiz, 1979; Karaman, 1982; Ruffo, 1982; Galan, 1984; González, 1991; Wakabara et al., 1991; Procaccini & Scipione, 1992; Camp et al., 1998; Wakabara & Serejo, 1998; LeCroy, 2000; Escobar et al., 2002; LeCroy, 2002; Martín & Díaz, 2003; Ortiz et al., 2007; Winfield & Escobar, 2007; LeCroy et al., 2009; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Martín et al., 2013; Ortega, 2013; Paz-Ríos & Ardisson, 2013; http://www.skaphandrus.com
<i>Eriichtonius brasiliensis</i> Figure 6	Barnard, 1970; Watling & Maurer, 1972; Bousfield, 1973; McKinney, 1977; Lincoln, 1979; Dickinson et al., 1980; Leite et al., 1980; Ruffo, 1982; Myers & McGrath, 1984; Wakabara et al., 1991; Procaccini & Scipione, 1992; Ortiz & Lalana, 1993; Rakocinski et al., 1993; Thomas, 1993a; Nelson, 1995; Appadoo & Steele, 1998; Camp et al., 1998; Wakabara & Serejo, 1998; Eggleston et al., 1999; Lowry, 2000; Cházaro et al., 2002; Escobar & Winfield, 2003; Martín & Díaz, 2003; Varela et al., 2003; Appadoo & Myers, 2004; Barba & Sánchez, 2005; Winfield et al., 2006; Ortiz et al., 2007; Winfield, Abarca-Arenas, et al., 2007; Dubois et al., 2009; Cerrato et al., 2010; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Winfield et al., 2010; Winfield & Ortiz, 2011; Martín et al., 2013; Ortega, 2013; Paz-Ríos & Ardisson, 2013; A, B, C

Geographic distribution	References
<i>Monocorophium acherusicum</i> Figure 6	Shoemaker, 1934; Barnard, 1970, 1971, 1972; Watling & Maurer, 1972; Bousfield, 1973; Lincoln, 1979; Myers, 1982; Ruffo, 1982; Barnard & Karaman, 1991; Wakabara et al., 1991; Procaccini & Scipione, 1992; Ortiz & Lalana, 1993; Nelson, 1995; Appadoo & Steele, 1998; Camp et al., 1998; Wakabara & Serejo, 1998; Eggleston et al., 1999; Lowry, 2000; Appadoo & Myers, 2004; LeCroy, 2004; Winfield et al., 2006; Ortiz et al., 2007; Somaio & Moreira, 2008; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Winfield et al., 2011; Martín et al., 2013; Ortega & Martín, 2013; Winfield, 2013; A, B
Bay of Fundy and Nova Scotia, Canada; Maine, New Hampshire, Massachusetts, Rhode Island, Connecticut, New Jersey, Delaware Bay, Chesapeake Bay, North Carolina, Georgia, Florida and Florida Keys, west Florida to Texas, US; Laguna de Tamiahua and Veracruz reef system in Veracruz, southwestern Gulf of Mexico, Quintana Roo, Mexico; Sigsbee abyssal plain; Cuba; Puerto Rico; Lesser Antilles; Venezuela; Pernambuco to Santa Catarina, Brazil. – Alaska to California, US; Bahía San Quintín, Baja California, Mexico; Hawaii; United Kingdom and the Netherlands south to the Mediterranean Sea including France, Italy, Egypt, Tunisia and Algeria; Senegal; Cape Colony; Dar es Salaam; Mauritius; Indian Ocean; Japan; South China Sea; Australia; New Zealand.	
<i>Batea catherinensis</i> Figure 6	Barnard, 1972; Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Wakabara et al., 1991; Ortiz & Lalana, 1993; Nelson, 1995; Camp et al., 1998; Wakabara & Serejo, 1998; Eggleston et al., 1999; Escobar & Winfield, 2003; LeCroy, 2004; Barba & Sánchez, 2005; Ortiz et al., 2007; Cerrato et al., 2010; Miloslavich et al., 2010; Ortiz & Lalana, 2010; Martín et al., 2013; Ortega, 2013; A, B, C
Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, Long Island Sound in New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina, South Carolina, Georgia, Florida, Tampa Bay, Mississippi Sound, Louisiana, US; Laguna Madre in Tamaulipas, Quintana Roo, Mexico; Caribbean Sea; Cuba; Barbados to São Paulo and Santa Catarina, Brazil. – Cedros Island and Magdalena Bay in Baja California, Mexico.	
<i>Ampithoe valida</i> Figure 6	Watling & Maurer, 1972; Bousfield, 1973; Fox & Bynum, 1975; Alonso, Tablado, López-Gappa, & Magaldi, 1995; Camp et al., 1998; Eggleston et al., 1999; LeCroy, 2002; Martín & Díaz, 2003; López Gappa et al., 2006; Ortiz et al., 2007; Miloslavich et al., 2010; Martín et al., 2013
Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, Long Island Sound in New York, New Jersey, Delaware Bay, Chesapeake Bay, North Carolina, South Carolina, Georgia, Cape Canaveral in Florida, Apalachee Bay in Florida to Galveston Bay in Texas, US; Laguna Madre in Tamaulipas, Laguna de Tamiahua in Veracruz, Quintana Roo, Mexico; Caribbean Sea; Venezuela; Quequén, Rio Negro, Chubut = Group A: 36°–43°S, Argentina. – British Columbia and Vancouver, Canada to Newport Bay in California, US; Japan; Korea.	
<i>Cerapus tubularis</i> Figure 6	Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Barnard & Karaman, 1991; Wakabara et al., 1991; Rakocinski et al., 1993; Nelson, 1995; Camp et al., 1998; Wakabara & Serejo, 1998; Barba & Sánchez, 2005; LeCroy, 2007; Ortiz et al., 2007; Dubois et al., 2009; Miloslavich et al., 2010; A, B, C
Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina, Georgia, east Florida, Perdido Key, Louisiana, Texas, US; Laguna Madre in Tamaulipas, Laguna de Tamiahua and Laguna de Alvarado in Veracruz, Quintana Roo, Mexico; Cuba; Colombia; São Paulo and Paraná, Brazil; Río Deseado, Argentina. – San Diego in California, US to Ensenada in Baja California, Mexico; Japan; Korea.	
<i>Lembos websteri</i> + Figure 7	Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Ruffo, 1982; Barnard & Karaman, 1991; Ortiz & Lalana, 1993; Camp et al., 1998; Eggleston et al., 1999; LeCroy, 2002; Escobar & Winfield, 2003; Barba & Sánchez, 2005; Ortiz et al., 2007
Greenland; Maine, New Hampshire, Cape Cod, Vineyard Sound in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina to Georgia and north Florida, west Florida, US; Laguna Madre in Tamaulipas, Mexico. – Norway; England; France; Spain; Italy; Yugoslavia; Greece; Turkey; Israel; Egypt; Algeria in the Mediterranean Sea; west Africa.	
<i>Monocorophium tuberculatum</i> Figure 7	Shoemaker, 1934, 1947; Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Bousfield & Hoover, 1997; Camp et al., 1998; LeCroy, 2004; Ortiz et al., 2007; Dubois et al., 2009; Ortega, 2013
Minas Basin, Bay of Fundy and Nova Scotia, Canada; Maine, New Hampshire, Cape Cod in Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina, Georgia, Sebastian inlet in Florida, Florida Bay to Chandeleur Sound in Louisiana, US; Laguna Madre in Tamaulipas, Mexico.	
<i>Orchestia grillus</i> Figure 7	Watling & Maurer, 1972; Bousfield, 1973; Thomas, 1976; Heard, 1982; Nelson, 1995; Camp et al., 1998; Ortiz et al., 2007; Miloslavich et al., 2010; Martín et al., 2013

Geographic distribution	References
<i>Apohyale prevostii</i> Figure 7 Labrador, Gulf of St Lawrence, Nova Scotia, Canada; Maine, New Hampshire, Massachusetts, Rhode Island, Connecticut, US; Laguna Madre in Tamaulipas, Laguna de Tamiahua in Veracruz, Mexico; Bermuda; Pernambuco, Brazil. – Arctic Ocean; Iceland; Ireland; United Kingdom; Faroe Islands; Norway; Netherlands; Belgium; France; Spain.	Bousfield, 1973; Lincoln, 1979; Wakabara & Serejo, 1998; Chavanich & Wilson, 2000; Barba & Sánchez, 2005; Lowry, 2014
<i>Jassa falcata</i> + Figure 7 Newfoundland and Nova Scotia, Canada; Maine, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Delaware Bay, Chesapeake Bay, Virginia, North Carolina, Georgia, Florida, Texas, US; Laguna Madre in Tamaulipas, Mexico; São Paulo, Brazil; Group A: 36°–43°S, Argentina. – Ireland; United Kingdom; Norway; Spain; Mediterranean Sea; British Columbia, Canada to California, US; South China Sea; New Zealand.	Barnard, 1972; Watling & Maurer, 1972; Bousfield, 1973; Dickinson et al., 1980; Leite et al., 1980; Barnard & Karaman, 1991; Wakabara et al., 1991; Nelson, 1995; Camp et al., 1998; Wakabara & Serejo, 1998; Lowry, 2000; Barba & Sánchez, 2005; López Gappa et al., 2006; Ortiz et al., 2007; Cerrato et al., 2010; C
<i>Orchestia gammarellus</i> + Figure 7 Newfoundland to Cape Breton Island, Nova Scotia, Bay of Fundy, Canada to Maine, New Hampshire, Massachusetts, Rhode Island, Connecticut, US; Laguna Madre in Tamaulipas, Mexico; Pernambuco, Brazil; Group B: 36°–58°S, Argentina. – Iceland; Norway; United Kingdom; Spain; Mediterranean Sea; Black Sea, Turkey; southwest Africa.	Bousfield, 1973; Soares, 1979; Wakabara & Serejo, 1998; Barba & Sánchez, 2005; López Gappa et al., 2006; Ortega, 2013; B, C

References

- Ahyong, S. T., Lowry, J. K., Alonso, M., Bamber, R. N., Boxshall, G. A., Castro, P., et al. (2011). Subphylum Crustacea Brünnich, 1772. In Z. Q. Zhang (Ed.), *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness* (3148) (pp. 165–191). Zootaxa.
- Alonso, G., Tablado, A., López-Gappa, J., & Magaldi, N. (1995). Seasonal changes in an intertidal population of the amphipod *Ampithoe valida* Smith, 1873. *Oebalia, XXI*, 77–91.
- Appadoo, C., & Myers, A. A. (2003). The genus *Elasmopus* (Crustacea: Amphipoda: Melitidae) from Mauritius (Indian Ocean) with description of five new species. *Records of the Australian Museum*, 55, 61–84.
- Appadoo, C., & Myers, A. A. (2004). Corophiidea (Crustacea: Amphipoda) from Mauritius. *Records of the Australian Museum*, 56, 331–362.
- Appadoo, C., & Steele, D. (1998). Shallow-water marine gammaridean amphipods of Mauritius Island. *Crustaceana*, 71, 633–645.
- Asari, K. P., & Myers, A. A. (1982). Taxonomic studies on the genus *Grandidierella* Coutiere (Crustacea, Amphipoda). IV. Indian species. *Bulletin du Muséum National d'Histoire Naturelle, Paris*, 4, 237–256.
- Bakir, K., Sezgin, M., & Katagan, T. (2010). Alien amphipods on the Turkish coasts. *Zoologica Baetica*, 21, 191–196.
- Baldo, F., Arias, A. M., & Drake, P. (2001). The macrobenthic community of the Guadalquivir estuary. *Boletín del Instituto Español de Oceanografía*, 17, 137–148.
- Barba, E., & Sánchez, A. J. (2005). Peracarid crustaceans of central Laguna Madre, Tamaulipas in the southwestern Gulf of Mexico. *Gulf of Mexico Science*, 23, 241–247.
- Barnard, J. L. (1962). Benthic marine Amphipoda of Southern California: 1. Families Aoridae, Photidae, Ischyroceriae, Corophiidae, Podoceridae, 2. Families Tironidae to Gammaridae, 3. Families Amphilochidae, Leucothoidae, Stenothoidae, Argissidae, Hyalidae. *Pacific Naturalist*, 3, 1–163.
- Barnard, J. L. (1963). Los anfípodos bentónicos marinos de la costa occidental de Baja California. *Revista de la Sociedad Mexicana de Historia Natural*, XXIV, 205–273.
- Barnard, J. L. (1969). The families and genera of marine gammaridean Amphipoda. *US Natural Museum Bulletin*, 271, 1–535.
- Barnard, J. L. (1970). Sublittoral gammaridean (Amphipoda) of the Hawaiian Islands. *Smithsonian Contributions in Zoology*, 34, 1–286.
- Barnard, J. L. (1971). Keys to the Hawaiian marine Gammaridea, 0–30 meters. *Smithsonian Contributions in Zoology*, 58, 1–135.
- Barnard, J. L. (1972). The marine fauna of New Zealand: algae-living littoral Gammaridea (Crustacea Amphipoda). *New Zealand Department of Scientific and Industrial Research, Bulletin*, 210, 1–216 [New Zealand Oceanographic Institute Memoir No. 62].
- Barnard, J. L., & Barnard, C. M. (1981). The amphipod genera *Eobrolgus* and *Eyakia* (Crustacea: Phoxocephalidae) in the Pacific Ocean. *Proceedings of the Biological Society of Washington*, 94, 295–313.
- Barnard, J. L., & Karaman, G. S. (1991). The families and genera of marine gammaridean Amphipoda (except marine Gammaroids). *Records of the Australian Museum (Supplement)*, 13, 1–866.
- Barnwell, F. H., & Thurman, C. L., II. (1984). Taxonomy and biogeography of the fiddler crabs (Ocypodidae: genus *Uca*) of the Atlantic and Gulf coasts of eastern North America. *Zoological Journal of the Linnean Society*, 81, 23–87.
- Bousfield, E. L. (1973). *Shallow-water gammaridean amphipoda of New England*. London: Cornell University Press.
- Bousfield, E. L., & Hoover, P. M. (1997). The amphipod superfamily Corophioidea on the Pacific coast of North America. Part 5. Family Corophiidae: Corophiinae, new subfamily. Systematics and distributional ecology. *Amphipacifica*, 2, 67–139.
- Briggs, J. C. (1974). *Marine zoogeography*. New York: McGraw-Hill.
- Bynum, K. H., & Fox, R. S. (1977). New and noteworthy amphipod crustaceans from North Carolina, US. *Chesapeake Science*, 18, 1–33.
- Camp, D. K., Lyons, W. G., & Perkins, T. H. (1998). *Checklists of selected shallow-water marine invertebrates of Florida*. Technical Report TR-3. St. Petersburg: Florida Marine Research Institute.
- Castilla, J. C., & Guikez, R. (2000). Disjoint geographical distribution of intertidal and nearshore benthic invertebrates in the Southern Hemisphere. *Revista Chilena de Historia Natural*, 73, 585–603.
- Cerrato, R. M., Flood, R. D., & Holt, L. C. (2010). *Benthic mapping for habitat classification in the Peconic estuary: phase III ground truth studies*. Stony Brook, NY 11794-5000. MSRC Special Report #137: Final report to the Suffolk County Department of Health Services Office of Ecology. Marine Sciences Research Center, School of Marine and Atmospheric Sciences, Stony Brook University.
- Chapman, J. W. (1988). Invasions of the northeast Pacific by Asian and Atlantic Gammaridean amphipod crustaceans, including a new species of *Corophium*. *Journal of Crustacean Biology*, 8, 364–382.

- Chavanich, S., & Wilson, K. A. (2000). Rocky intertidal zonation of gammaridean amphipods in Long Island Sound, Connecticut. *Crustaceana-International Journal of Crustacean Research*, 73, 835–846.
- Cházaro, S., Winfield, I., Ortiz, M., & Álvarez, F. (2002). Peracard crustaceans from three inlets in the southwestern Gulf of Mexico: new records and range extensions. *Zootaxa*, 123, 1–16.
- Clarke, K. R., & Gorley, R. N. (2006). PRIMER v6: user manual/tutorial. Plymouth, UK: PRIMER-E.
- Conlan, K. E. (1990). Revision of the crustacean amphipod genus *Jassa* Leach (Corophioidea: Ischyroceridae). *Canadian Journal of Zoology*, 68, 2031–2075.
- Corona, A., & Raz-Guzmán, A. (2003). Distribución geográfica de los anfípodos e isópodos (Crustacea: Peracarda: Amphipoda e Isopoda) de los sistemas estuarinos de Michoacán, México. In M. E. Hendrickx (Ed.), *Contribuciones al estudio de los crustáceos del Pacífico Este 2* (pp. 219–225). Mexico, D.F.: Instituto de Ciencias del Mar y Limnología, UNAM.
- Corona, A., Soto, L. A., & Sánchez, A. J. (2000). Epibenthic amphipod abundance and predation efficiency of the pink shrimp *Farfantepenaeus duorarum* (Burkenroad, 1939) in habitats with different physical complexity in a tropical estuarine system. *Journal of Experimental Marine Biology and Ecology*, 253, 33–48.
- Craw, R. C., Grehan, J. R., & Heads, M. J. (1999). *Panbiogeography*. Oxford: Oxford University Press.
- DeBlois, E. M., & Leggett, W. C. (1993). Impact of amphipod predation on the benthic eggs of marine fish: an analysis of *Calliopius laeviusculus* bioenergetic demands and predation on the eggs of a beach spawning osmeriid (*Mallotus villosus*). *Marine Ecology Progress Series*, 93, 205–216.
- Díaz, Y. J., & Martín, A. (2001). New records of amphipods (Crustacea: Amphipoda) from shallow waters of the Caribbean coast of Venezuela. *Revista de Biología Tropical*, 49, 1271–1276.
- Dickinson, J. J., Wigley, R. L., Brodeur, R. D., & Brown-Leger, S. (1980). *Distribution of gammaridean Amphipoda (Crustacea) in the Middle Atlantic Bight Region*. US Department of Commerce, NOAA Technical Report NMFS SSRF-741.
- Dubois, S., Gelpi, C. G., Jr., Condrey, R. E., Grippo, M. A., & Fleeger, J. W. (2009). Diversity and composition of macrobenthic community associated with sandy shoals of the Louisiana continental shelf. *Biodiversity and Conservation*, 18, 3759–3784.
- d'Udekem-d'Acoz, C., & Verheyen, M. (2013). Taxocoenoses of amphipod crustaceans. In J. Gutt (Ed.), *The expedition of the research vessel "Polarstern" to the Antarctic in 2013 (ANT-XXIX/3). Berichte zur Polar – und Meeresforschung* (Vol. 665) (pp. 57–67).
- Duffy, J. E. (1990). Amphipods on seaweeds: partners or pests? *Oecologia*, 83, 267–276.
- Eggleson, D. B., Elis, W. E., Etherington, L. L., Dahlgren, C. P., & Posey, M. H. (1999). Organism responses to habitat fragmentation and diversity: habitat colonization by estuarine macrofauna. *Journal of Experimental Marine Biology and Ecology*, 236, 107–132.
- Escobar, E., & Soto, L. A. (1997). Continental shelf benthic biomass in the western Gulf of Mexico. *Continental Shelf Research*, 17, 585–604.
- Escobar, E., & Winfield, I. (2003). Checklist of the benthic Gammaridea and Caprellidea (Crustacea: Peracarda: Amphipoda) from the Gulf of Mexico continental shelf and slope. *Belgian Journal of Zoology*, 133, 37–44.
- Escobar, E., Winfield, I., Ortiz, M., Gasca, E., & Suárez, E. (2002). Amphipoda. In J. Llorente-Bousquets, & J. J. Morrone (Eds.), *Biodiversidad, taxonomía y biogeografía de artrópodos de México: hacia una síntesis de su conocimiento actual* (Vol. III) (pp. 341–371). México, D.F.: Conabio/UNAM.
- Faasse, M., & van Moorsel, G. (2003). The North-American amphipods, *Melita nitida* Smith, 1873 and *Incisocalliope aestuarius* (Watling and Maurer, 1973) (Crustacea: Amphipoda: Gammaridea), introduced to the Western Scheldt estuary (The Netherlands). *Aquatic Ecology*, 37, 13–22.
- Fernández, A., Zavala, J., & Romero, R. (1998). *Circulación de invierno en la plataforma de Tamaulipas y áreas adyacentes. IX Reunión Nacional SELPER-MEXICO 1998, Desarrollo Sustentable. Zacatecas, México. Memorias en CD-ROM*.
- Fox, R. S., & Bynum, K. H. (1975). The amphipod crustaceans of North Carolina estuarine waters. *Chesapeake Science*, 16, 223–237.
- Galan, A. (1984). *A systematic study of Amphipoda (Crustacea) of the Caribbean coast of Venezuela* (Ph.D. thesis). London: Department of Pure and Applied Biology, Imperial College of Science & Technology, London University.
- Gardner, W., Escobar, E., Cruz, E., & Rowe, G. (1993). Ammonium excretion by benthic invertebrates and sediment-water nitrogen flux in the Gulf of Mexico near the Mississippi River outflow. *Estuaries*, 16, 799–808.
- González, E. (1991). The genus *Hyale* in Chile. *Spixiana*, 14, 125–142.
- Havermans, C., De Broyer, C., Mallefet, J., & Zintzen, V. (2007). Dispersal mechanisms in amphipods: a case study of *Jassa herdmani* (Crustacea Amphipoda) in the North Sea. *Marine Biology*, 153, 83–89.
- Hay, M. E., Duffy, J. E., Fenical, W., & Gustafson, K. (1988). Chemical defense in the seaweed *Dictyopteris delicatula*: differential effects against reef fishes and amphipods. *Marine Ecology Progress Series*, 48, 185–192.
- Heard, R. W. (1982). *Guide to common tidal marsh invertebrates of the northeastern Gulf of Mexico*. Mississippi Alabama Sea Grant Consortium, MASGP-79-004.
- Heck, K. L., Jr., & Crowder, L. B. (1991). Habitat structure and predator-prey interactions in vegetated aquatic systems. In S. S. Bell, E. D. McCoy, & H. R. Mushinsky (Eds.), *Habitat structure: the physical arrangement of objects in space* (pp. 281–299). London: Chapman & Hall.
- Hedgpeth, J. W. (1953). pp. 107–224. *An introduction to the zoogeography of the northwestern Gulf of Mexico with reference to the invertebrate fauna* (3) Publications of the Institute of Marine Science, University of Texas.
- Highsmith, R. C. (1985). Floating and algal rafting as potential dispersal mechanisms in brooding invertebrates. *Marine Ecology Progress Series*, 25, 169–179.
- Hill, C., & Elmgren, R. (1992). Predation by the isopod *Saduria entomon* on the amphipods *Monoporeia affinis* and *Pontoporeia hoyi*: experiments on prey vulnerability. *Oecologia*, 91, 153–156.
- Horton, T., Lowry, J., & De Broyer, C. (2013). *World Amphipoda database*. Retrieved from <http://www.marinespecies.org/amphipoda/>
- Johnson, W. S., Stevens, M., & Watling, L. (2000). Reproduction and development of marine peracarideans. *Advances in Marine Biology*, 39, 107–220.
- Kamermans, P., Malta, E. J., Verschueren, J. M., Schrijvers, L., Lentz, L. F., & Lien, A. T. A. (2002). Effect of grazing by isopods and amphipods on growth of *Ulva* spp. (Chlorophyta). *Aquatic Ecology*, 36, 425–433.
- Karaman, G. S. (1982). Family Gammaridae. In S. Ruffo (Ed.), *The Amphipoda of the Mediterranean, Part 1. Gammaridea (Acanthonoto-zomatidae to Gammaridae)* (pp. 245–364). Monaco: Mémoires de l'Institut Oceanographique 13.
- Lagarde, G. (1987). Anfípodos Gammaridea del litoral de Golfo Triste y áreas adyacentes. *Caribbean Journal of Science*, 23, 260–277.
- LeCroy, S. E. (2000). *An illustrated identification guide to the nearshore marine and estuarine gammaridean Amphipoda of Florida*. Vol. 1. Families Gammaridae, Hadziidae, Isaedae, Melitidae and Oedicerotidae. Tallahassee, Florida: Annual Report for the Department of Environmental Protection. Contract Number WM724.
- LeCroy, S. E. (2002). *An illustrated identification guide to the nearshore marine and estuarine gammaridean Amphipoda of Florida*. Vol. 2. Families Ampeliscidae, Ampelochidae, Ampithoidae, Aoridae, Argissidae and Hausoriidae. Tallahassee, Florida: Annual Report for the Department of Environmental Protection. Contract Number WM724.
- LeCroy, S. E. (2004). *An illustrated identification guide to the nearshore marine and estuarine gammaridean Amphipoda of Florida*. Vol. 3. Families Bateidae, Biancolinidae, Cheluridae, Colomastigidae, Corophiidae, Cypridoidea and Dexaminidae. Tallahassee, Florida: Annual Report for the Department of Environmental Protection. Contract Number WM724.
- LeCroy, S. E. (2007). *An illustrated identification guide to the nearshore marine and estuarine gammaridean Amphipoda of Florida*. Vol. 4. Families Anamixidae, Eusiridae, Hyalellidae, Hyalidae, Iphimediidae, Ischyroceridae, Lysianassidae, Megaluropidae and Melphidippidae. Tallahassee, Florida: Annual Report for the Department of Environmental Protection. Contract Number WM724.
- LeCroy, S. E. (2011). *An illustrated identification guide to the nearshore marine and estuarine gammaridean Amphipoda of Florida*. Vol. 5. Families Leucothoidae, Liljeborgiidae, Neomegamphopidae, Ochlesidae, Phliantidae, Phoxocephalidae, Platyschnopidae, Pleustidae, Podoceridae,

- Pontoporeiidae, Sebidae, Stenothoidae, Synopiidae and Talitridae. Tallahassee, Florida: Annual Report for the Department of Environmental Protection.
- LeCroy, S. E., Gasca, R., Winfield, I., Ortiz, M., & Escobar-Briones, E. (2009). Amphipoda (Crustacea) of the Gulf of Mexico. In D. Felder, & D. Camp (Eds.), *Gulf of Mexico. Origin, waters and biota, Volume I, Biodiversity* (pp. 941–972). College Station: Texas A&M University Press.
- Ledoyer, M. (1986). Faune mobile des herbiers de phanérogames marines (*Halodule* et *Thalassia*) de la Lagune de Términos (Mexique, Campeche). II. Les gammariens (Crustacea). *Anales del Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México*, 13, 171–200.
- Leite, F. P. P., Tararam, A. S., & Wakabara, Y. (1980). Composição e distribuição da fauna de Gammaridea na região da Enseada da Fortaleza-Ubatuba, Estado de São Paulo. *Boletim do Instituto Oceanográfico*, 29, 297–299.
- Lincoln, R. J. (1979). *British marine Amphipoda: Gammaridea*. London: British Museum (Natural History).
- Locke, A., & Corey, S. (1989). Amphipods, isopods and surface currents: a case for passive dispersal in the Bay of Fundy Canada. *Journal of Plankton Research*, 11, 419–430.
- López-Gappa, J., Alonso, G. M., & Landoni, N. A. (2006). Biodiversity of benthic Amphipoda (Crustacea: Peracarida) in the southwest Atlantic between 35°S and 56°S. *Zootaxa*, 1342, 1–66.
- Lowry, J. K. (1972). Taxonomy and distribution of Microprotopus along the east coast of the United States (Amphipoda: Isaeidae). *Crustaceana (Suppl.)*, 3, 277–286.
- Lowry, J. K. (2000). Taxonomic status of amphipod crustaceans in the South China Sea with a checklist of known species. *The Raffles Bulletin of Zoology (Suppl.)*, 8, 309–342.
- Lowry, J. K. (2014). *Hyale prevostii* (Milne-Edwards, 1830). In T. Horton, J.K. Lowry, & C. de Broyer (2013 onwards) World Amphipoda. Retrieved from <http://www.marinespecies.org/amphipoda/aphia.php?p=taxdetails&id=102326>.
- Lowry, J. K., & Stoddart, H. E. (2003). Crustacea: Malacostraca: Peracarida: Amphipoda, Cumacea Mysidacea. In P. L. Beesley, & W. W. K. Houston (Eds.), *Zoological Catalogue of Australia* (Vol. 19.2B). Melbourne: CSIRO Publishing.
- Martin, J. W., & Davis, G. E. (2001). *An updated classification of the recent Crustacea* (Science Series, pp. 39). Natural History Museum of Los Angeles County.
- Martín, A., & Díaz, Y. J. (2003). La fauna de anfípodos (Crustacea: Amphipoda) de las aguas costeras de la región oriental de Venezuela. *Boletín del Instituto Español de Oceanografía*, 19, 327–344.
- Martín, A., Ortiz, M., & Díaz, Y. J. (2002). Nuevos registros crustáceos anfípodos (Gammaridea) colectados en las costas del caribe venezolano. *Boletín de Investigaciones Marinas y Costeras*, 31, 15–24.
- Martín, A., Díaz, Y., Miloslavich, P., Escobar-Briones, E., Guerra-García, J. M., Ortiz, M., et al. (2013). Regional diversity of Amphipoda in the Caribbean Sea. *Revista de Biología Tropical*, 61, 1681–1720.
- McKinney, L. D. (1977). *The origin and distribution of shallow water gammaridean Amphipoda in the Gulf of Mexico and Caribbean Sea with notes on their ecology*. Austin: Texas A&M University.
- McKinney, L. D. (1978). Amphilochidae (Crustacea: Amphipoda) from the western Gulf of Mexico and Caribbean Sea. *Gulf Research Reports*, 6, 137–143.
- McLaughlin, P. A., Camp, D. K., Angel, M. V., Bousfield, E. L., Brunei, P., Brusca, R. C., et al. (2005). *Common and scientific names of aquatic invertebrates from the United States and Canada: Crustaceans*. Bethesda, Maryland: American Fisheries Society Special Publication 31.
- Melo, G. A. S. (1990). A presença, no litoral sudeste brasileiro, de espécies de Brachyura (Crustacea: Decapoda) originárias das regiões biogeográficas Magelânica e Argentina do Atlântico Sul. *Atlântica*, 12, 71–83.
- Mills, E. L. (1963). A new species of *Ampelisca* (Crustacea: Amphipoda) from eastern North America, with notes on other species of the genus. *Canadian Journal of Zoology*, 41, 971–989.
- Miloslavich, P., Díaz, J. M., Klein, E., Alvarado, J. J., Díaz, C., Gobin, J., et al. (2010). Marine biodiversity in the Caribbean: regional estimates and distribution patterns. *PLoS ONE*, 5, e11916.
- Montagna, P., & Harper, E. (1996). Benthic faunal long term response to offshore production platforms in the Gulf of Mexico. *Canadian Journal of Fisheries and Aquatic Science*, 53, 2567–2588.
- Myers, A. A. (1970). Taxonomic studies on the genus *Grandidierella* Coutiere (Crustacea: Amphipoda) with a description of *G. dentimera* sp. nov. *Bulletin of Marine Science*, 20, 135–147.
- Myers, A. A. (1982). Family Corophiidae. In S. Ruffo (Ed.), *The Amphipoda of the Mediterranean, Part 1, Gammaridea (Acanthonotozomatidae to Gammaridae)* (13) (pp. 185–208). Monaco: Mémoires de l'Institut Océanographique.
- Myers, A. A. (1985). Shallow-water, coral reef and mangrove Amphipoda (Gammaridea) of Fiji. *Records of the Australian Museum (Suppl.)*, 5, 1–143.
- Myers, A. A., & Lowry, J. K. (2003). A phylogeny and new classification of the Corophiidae Leach 1814 (Amphipoda). *Journal of Crustacean Biology*, 23, 443–485.
- Myers, A. A., & McGrath, D. (1984). A revision of the north-east Atlantic species of *Ericthonius* (Crustacea: Amphipoda). *Journal of the Marine Biological Association of the United Kingdom*, 64, 379–400.
- Naomi, T. S. (1979). On a swarm of amphipods *Atylus minikoi* (Walker) in the shallow waters of the Karwar Bay. *Indian Journal of Fisheries*, 26, 227–228.
- Nelson, W. G. (1995). Amphipod crustaceans of the Indian River Lagoon: current status and threats to biodiversity. *Bulletin of Marine Science*, 57, 143–152.
- Norton, T. A., & Mathieson, A. C. (1983). The biology of unattached seaweeds. In F. E. Round, & D. J. Chapman (Eds.), *Progress in phycological research* (Vol. 2) (pp. 333–386). Amsterdam/New York/Oxford: Elsevier.
- Oliveira, L. P. H. (1953). Crustacea Amphipoda de Rio de Janeiro. *Memorias del Instituto Oswaldo Cruz*, 51, 289–376.
- Ortega, V. M. (2013). *Taxonomía y distribución ecológica de los crustáceos peracáridos de la Laguna Madre y regiones costeras adyacentes, Tamaulipas México* (M.Sc. thesis). Monterrey, N.L., México: Universidad Autónoma de Nuevo León.
- Ortega, I., & Martín, A. (2013). Suprabenthic amphipods from the littoral zone of Barlovento Venezuela: spatial distribution and seasonal variation. *Crustaceana*, 86, 1206–1223.
- Ortiz, M. (1978). Invertebrados marinos bentónicos de Cuba. I. Crustacea, Amphipoda, Gammaridea. *Revista de Investigaciones Marinas, Serie 8*, 38, 3–10.
- Ortiz, M. (1979). Contribución al estudio de los anfípodos (Gammaridea) del Mediterráneo americano. *Revista de Investigaciones Marinas, Serie 8*, 1–16.
- Ortiz, M. (1983). Los anfípodos (Gammaridea) de las costas del Mar Caribe de la República de Colombia. *Revista de Investigaciones Marinas*, 4, 23–31.
- Ortiz, M. (2001). Lista de invertebrados marinos, estuarinos y semiterrestres de la playa de Cojímar en la costa norte de la Provincia Ciudad de la Habana. *Revista de Investigaciones Marinas*, 22, 93–102.
- Ortiz, M., & Lalana, R. (1993). Adición a la lista de especies y bibliografía de los anfípodos (Crustacea: Amphipoda) del Mediterráneo Americano. *Revista de Investigaciones Marinas*, 14, 16–37.
- Ortiz, M., & Lalana, R. (2010). Distribución de los anfípodos (Crustacea, Malacostraca, Peracarida) de los subórdenes Gammaridea Caprellidea e Hyperiidea, presentes en el archipiélago cubano. *Revista de Investigaciones Marinas*, 31, 75–90.
- Ortiz, M., Martín, A., & Díaz, Y. J. (2007). Lista y referencias de los crustáceos anfípodos (Amphipoda: Gammaridea) del Atlántico occidental tropical. *Revista de Biología Tropical*, 55, 479–498.
- Paz-Ríos, C. E., & Ardisson, P. L. (2013). Benthic amphipods (Amphipoda: Gammaridea and Corophiidea) from the Mexican southeast sector of the Gulf of Mexico: checklist, new records and zoogeographic comments. *Zootaxa*, 3635, 137–173.
- Paz-Ríos, C. E., Simões, N., & Ardisson, P. L. (2013). Intertidal and shallow water amphipods (Amphipoda: Gammaridea and Corophiidea) from Isla Pérez Alacranes Reef, southern Gulf of Mexico. *Nauplius*, 21, 179–194.
- Probert, P. K., & Grove, S. L. (1998). Macrobenthic assemblages of the continental shelf and upper slope of the west coast of South Island, New Zealand. *Journal of the Royal Society of New Zealand*, 28, 259–280.
- Procaccini, G., & Scipione, M. B. (1992). Observations on the spatio-temporal distribution of crustacean amphipods in the Fusaro coastal lagoon (Central

- Tyrrhenian Sea Italy) and some notes on their presence in Mediterranean lagoons. *Marine Ecology*, 13, 203–224.
- Rakocinski, C. F., Heard, R. W., LeCroy, S. E., McLelland, J. A., & Simons, T. (1993). Seaward change and zonation of the sandy-shore macrofauna at Perdido Key, Florida, U.S.A. *Estuarine, Coastal and Shelf Science*, 36, 81–104.
- Raz-Guzmán, A., & Barba, E. (2000). Seagrass biomass, distribution and associated macrofauna in southwestern Gulf of Mexico coastal lagoons. *Biotología Marina Mediterranea*, 7, 271–274.
- Raz-Guzmán, A., & Sánchez, A. J. (1992a). Registros adicionales de cangrejos braquiuros (Crustacea: Brachyura) de Laguna de Términos, Campeche. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 63, 29–45.
- Raz-Guzmán, A., & Sánchez, A. J. (1992b). Registros adicionales de cangrejos braquiuros (Crustacea: Brachyura) del sistema lagunar de Alvarado, Veracruz. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 63, 273–277.
- Raz-Guzmán, A., & Sánchez, A. J. (1996). *Catálogo ilustrado de cangrejos braquiuros (Crustacea) de la Laguna de Tamiahua, Veracruz, México. Cuadernos*, 31. México, D.F.: Instituto de Biología, UNAM.
- Raz-Guzmán, A., & Sánchez, A. J. (1998). Catálogo con sinonimias y notas sobre el hábitat de los cangrejos ermitaños estuarinos del suroeste del Golfo de México. *Universidad y Ciencia*, 14, 17–31.
- Raz-Guzmán, A., Sánchez, A. J., Soto, L. A., & Álvarez, F. (1986). Catálogo ilustrado de cangrejos braquiuros y anomuros de Laguna de Términos, Campeche (Crustacea: Brachyura, Anomura). *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 57, 343–383.
- Raz-Guzmán, A., Sánchez, A. J., & Soto, L. A. (1992). *Catálogo ilustrado de cangrejos braquiuros y anomuros (Crustacea) de Laguna de Alvarado, Veracruz México. Cuadernos*, 14. México, D.F.: Instituto de Biología, UNAM.
- Raz-Guzmán, A., Sánchez, A. J., Peralta, P., & Florido, R. (2004). Zoogeography of hermit crabs (Decapoda: Diogenidae, Paguridae) from four coastal lagoons in the Gulf of Mexico. *Journal of Crustacean Biology*, 24, 625–636.
- Ruffo, S. (1982). *The Amphipoda of the Mediterranean Part 1. Gammaridea (Acanthonotozomatidae to Gammaridae)*. pp. 13. Monaco: Mémoires de l'Institut Oceanographique.
- Salcedo, D. L., Soto, L. A., Estradas-Romero, A., & Botello, A. (2017). Interannual variability of soft-bottom macrobenthic communities of the NW Gulf of Mexico in relationship to the Deepwater Horizon oil spill. *Marine Pollution Bulletin*, 114, 987–994.
- Saunders, J. E., Attrill, M. J., Shaw, S. M., & Rowden, A. A. (2003). Spatial variability in the epiphytic algal assemblages of *Zostera marina* seagrass beds. *Marine Ecology Progress Series*, 249, 107–115.
- Serejo, C. S. (1999). Taxonomy and distribution of the family Hyaliidae (Amphipoda, Talitroidea) on the Brazilian coast. In F. R. Schram, & J. C. von Vaupel Klein (Eds.), *Crustaceans and the biodiversity crisis* (pp. 591–616). Amsterdam, The Netherlands, Koninklijke Brill NV, Leiden: Proceedings of the Fourth International Crustacean Congress.
- Serejo, C. S. (2004). Cladistic revision of talitroidean amphipods (Crustacea Gammaridea), with a proposal of a new classification. *Zoologica Scripta*, 33, 551–586.
- Shoemaker, C. R. (1933). Amphipoda from Florida and the West Indies American Museum. *Novitates*, 598, 1–24.
- Shoemaker, C. R. (1934). The amphipod genus *Corophium* on the east coast of America. *Proceedings of the Biological Society of Washington*, 47, 23–32.
- Shoemaker, C. R. (1947). Further notes on the amphipod genus *Corophium* from the east coast of America. *Journal of the Washington Academy of Sciences*, 37, 47–63.
- Shoemaker, C. R. (1956). Observations on the amphipod genus *Parhyale*. *Proceedings of the United States National Museum*, 106, 345–358.
- Soares, C. M. A. (1979). Estudo ecológico da região de Itamaracá, Pernambuco Brasil. III. Anfípodos das famílias Talitridae e Ampithoidae. (1). *Trabalhos Oceanográficos da Universidade Federal de Pernambuco*, 14, 93–104.
- Somaio, C., & Moreira, R. (2008). Introduced and cryptogenic species and their management in Paranaguá Bay Brazil. *Brazilian Archives of Biology and Technology*, 51, 623–633.
- Stephensen, K. (1948). . pp. 1–20. *Amphipods from Curaçao, Bonaire, Aruba and Margarita. Studies on the fauna of Curaçao, Bonaire, Aruba and the Venezuelan Islands* (Vol. 11) Foundation for Scientific Research in Surinam and The Netherlands Antilles.
- Tanner, J. E. (2006). Landscape ecology of interactions between seagrass and mobile epifauna: the matrix matters. *Estuarine, Coastal and Shelf Science*, 68, 404–412.
- Thiel, M. (1997). Reproductive biology of an epibenthic amphipod (*Dyopedos monacanthus*) with extended parental care. *Journal of the Marine Biological Association of the United Kingdom*, 77, 1059–1072.
- Thiel, M. (2003a). Extended parental care in crustaceans – an update. *Revista Chilena de Historia Natural*, 76, 205–218.
- Thiel, M. (2003b). Rafting of benthic macrofauna: important factors determining the temporal succession of the assemblage on detached macroalgae. *Hydrobiologia*, 503, 49–57.
- Thiel, M., & Hinojosa, I. A. (2009). Peracáridos – Anfípodos, Isópodos, Tanaidáceos y Cumáceos. In V. Häussermann, & G. Förster (Eds.), *Fauna marina bentónica de la Patagonia chilena* (pp. 674–678). Santiago de Chile: Nature in Focus.
- Thomas, J. D. (1976). A survey of gammarid amphipods of the Barataria Bay Louisiana region. *Contributions in Marine Science*, 20, 87–100.
- Thomas, J. D. (1993a). *Identification manual for marine Amphipoda (Gammaridea): I. Common coral reef and rocky bottom amphipods of South Florida*. Tallahassee, Florida. Department of Environmental Protection. Retrieved from <ftp://ftp.dep.state.fl.us/pub/labs/biology/biokeys/sflamps.pdf>
- Thomas, J. D. (1993b). Biological monitoring and tropical biodiversity in marine environments: a critique with recommendations and comments on the use of amphipods as bioindicators. *Journal of Natural History*, 27, 795–806.
- Thurman, C. L., II. (1987). Fiddler crabs (Genus *Uca*) of eastern Mexico (Decapoda, Brachyura Ocypodidae). *Crustaceana*, 53, 94–105.
- Tunnell, J. W., & Judd, F. W. (2002). *The Laguna Madre of Texas and Tamaulipas*. College Station: Texas A&M University Press.
- Varela, C., Manuel, O., & Rogelio, L. (2003). Crustáceos (Peracarida y Decapoda) de la costa sur de la península de Guanahacabibes Cuba. *Revista de Investigaciones Marinas*, 24, 73–76.
- Wakabara, Y., & Serejo, C. S. (1998). Malacostraca – Peracarida, Amphipoda, Gammaridea and Caprellidea. In P. S. Young (Ed.), *Catalogue of Crustacea of Brazil* (pp. 561–594). Rio de Janeiro: Museu Nacional, Série Livros # 6.
- Wakabara, Y., Tararam, A. S., & Takeda, A. M. (1983). Comparative study of the amphipod fauna living on *Sargassum* of two Itanhaém shores. *Brazilian Journal of Crustacean Biology*, 3, 602–607.
- Wakabara, Y., Tararam, A. S., Valério-Berardo, M. T., Duleba, W., & Pereira, F. P. (1991). Gammaridean and caprellidean fauna from Brazil. *Hydrobiologia*, 223, 69–77.
- Wakabara, Y., Nicoletti, M., & Tararam, A. S. (1996). Ingestion and selection of suprabenthic crustaceans by small-sized fishes in a lower saltmarsh system. *Revista Brasileira de Oceanografia*, 44, 89–103.
- Watling, L., & Maurer, D. (1972). Shallow water amphipods of the Delaware Bay region. *Crustaceana Supplement*, 3, 251–266.
- Williams, A. B., & Bynum, K. H. (1972). A ten year study of meroplankton in North Carolina estuaries: amphipods. *Chesapeake Science*, 13, 175–192.
- Winfield, I. (2013). *Catálogo de las especies de crustáceos anfípodos invasores del Parque Nacional Sistema Arrecifal Veracruzano (PNSAV) y la actualización de la base de datos (CONABIO) de los anfípodos en México*. Mexico, D.F: FES-Iztacala, UNAM. Final report SNIB-CONABIO. Project No. GN011.
- Winfield, I., & Escobar, E. (2007). Anfípodos (Crustacea: Gammaridea) del sector norte del Mar Caribe: listado faunístico, registros nuevos y distribución espacial. *Revista Mexicana de Biodiversidad*, 78, 51–61.
- Winfield, I., & Ortiz, M. (2003). *Anfípodos: un enfoque biológico*. México, D.F.: Editorial Universidad Nacional Autónoma de México, FES-Iztacala.
- Winfield, I., & Ortiz, M. (2011). Crustáceos con bolsa incubadora (Crustacea: Malacostraca: Peracarida). In Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (Ed.), *La Biodiversidad en Veracruz: estudio de estado* (Vol. II) (pp. 277–286). México, D.F: Conabio, Gobierno del Estado de Veracruz, Universidad Veracruzana, Instituto de Ecología, A.C.

- Winfield, I., Ortiz, M., Franco, J., & Bedia, C. (1997). Distribución y diversidad del superorden Peracarida asociado a pastos marinos de Alvarado Veracruz. *Cuadernos Mexicanos de Zoología*, 3, 1–8.
- Winfield, I., Escobar, E., & Álvarez, F. (2001). Crustáceos peracáridos asociados a praderas de *Ruppia maritima* (Ruppiaceae) en la laguna de Alvarado, México. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 72, 29–41.
- Winfield, I., Escobar, E., & Morrone, J. J. (2006). Updated checklist and identification of areas of endemism of benthic amphipods (Caprellidea and Gammaridea) from offshore habitats in the SW Gulf of Mexico. *Scientia Marina*, 70, 99–108.
- Winfield, I., Abarca-Arenas, L. G., & Cházaro-Olvera, S. I. (2007). Crustacean macrofoulers in the Veracruz coral reef system SW Gulf of Mexico: checklist, spatial distribution and diversity. *Cahiers de Biologie Marine*, 48, 287–295.
- Winfield, I., Cházaro-Olvera, S., & Álvarez, R. (2007). ¿Controla la biomasa de pastos marinos la densidad de los peracáridos (Crustacea: Peracarida) en lagunas tropicales? *Revista de Biología Tropical*, 55, 43–53.
- Winfield, I., Cházaro-Olvera, S., Horta-Puga, G., Lozano-Aburto, M. A., & Arenas-Fuentes, V. (2010). Macrocrustáceos incrustantes en el Parque Nacional Sistema Arrecifal Veracruzano: biodiversidad, abundancia y distribución. *Revista Mexicana de Biodiversidad (Suplem.)*, 80, S165–S175.
- Winfield, I., Cházaro-Olvera, S., Ortiz, M., & Palomo-Aguayo, U. (2011). Lista actualizada de las especies de anfípodos (Peracarida: Gammaridea y Corophiidea) marinos invasores en México. *Revista de Biología Marina y Oceanografía*, 46, 349–361.
- Zupo, V., & Nelson, W. G. (1999). Factors influencing the association patterns of *Hippolyte zostericola* and *Palaemonetes intermedius* (Decapoda: Natantia) with seagrasses of the Indian River Lagoon Florida. *Marine Biology*, 134, 181–190.