



ORIGINAL ARTICLE

Do we publish what we preach? Analysis of Spanish Shoulder and Elbow Surgery Society publication rates[☆]

J. Miquel^{a,*}, S. Fernández-Muñoz^a, A. Romero^a, X. Pelfort^a, C. Torrens^b

^a Servicio de Cirugía Ortopédica y Traumatología, Hospital d'Igualada, Consorci Sanitari de l'Anoia, Igualada (Barcelona), Spain

^b Servicio de Cirugía Ortopédica y Traumatología, Parc de Salut Mar, Barcelona, Spain

Received 16 January 2017; accepted 3 September 2017



CrossMark

KEYWORDS

Publication rate;
Congress;
Abstracts;
Concordance

Abstract

Introduction: The purpose of this study is to analyse the publication rate of studies presented as podium presentations in the Spanish Society of Shoulder and Elbow Surgery (SECHC) congresses. **Methods:** One hundred and twenty-two abstracts presented at the SECHC congresses held in 2007, 2009 and 2011 were included for the purpose of the study. The oral communications were categorised by study type, sample included and follow-up. In June 2017, possible publications of these studies were searched in PubMed. Type of study, delay in publication, journal and impact factor obtained were recorded. The concordance between the information presented at the congress and their subsequent full-text publications was analysed.

Results: The publication rate was 17.21% (21 of the 122 abstracts studied) after 6 years, with a mean time spent for publications of 36.71 months, and a mean impact factor of 1.51. There were no differences between results initially presented at the congresses and those subsequently published ($p > 0.05$).

Discussion: The majority of papers presented at SECHC congresses do not end up with a publication. The papers that are published do not usually contain significant differences compared to the content delivered at the congress.

© 2017 Published by Elsevier España, S.L.U. on behalf of SECOT.

PALABRAS CLAVE

Tasa de publicación;
Congreso;
Resúmenes;
Concordancia

¿Publicamos lo que predicamos? Tasa de publicación de los congresos de la Sociedad Española de Cirugía de Hombro y Codo

Resumen

Introducción: El propósito de este estudio es analizar la tasa de publicación de los estudios presentados como comunicaciones orales en los congresos de la Sociedad Española de Cirugía de Hombro y Codo (SECHC).

[☆] Please cite this article as: Miquel J, Fernández-Muñoz S, Romero A, Pelfort X, Torrens C. ¿Publicamos lo que predicamos? Tasa de publicación de los congresos de la Sociedad Española de Cirugía de Hombro y Codo. Rev Esp Cir Ortop Traumatol. 2018;62:35–46.

* Corresponding author.

E-mail address: joanmiquelnoguera@hotmail.com (J. Miquel).

Métodos: Se incluyeron 122 resúmenes presentados en los congresos SECHC celebrados en 2007, 2009 y 2011. Se categorizaron las comunicaciones orales por tipo de estudio, muestra incluida y seguimiento. En junio de 2017 se buscaron posibles publicaciones de estos trabajos en PubMed. El tiempo para la publicación, revista y factor de impacto fueron registrados. Se analizó la concordancia entre la información presentada en el congreso y las publicaciones posteriores. **Resultados:** Veintiuno de los 122 resúmenes analizados (17,21%) fueron publicados después de 6 años de seguimiento mínimo, con un tiempo medio empleado para las publicaciones de 36,71 meses y con un factor de impacto medio de 1,51. No se observaron diferencias significativas entre los datos expuestos inicialmente en el congreso y los posteriormente publicados ($p > 0,05$). **Conclusiones:** La mayoría de los trabajos presentados en congresos SECHC no acaban siendo publicados. Aquellos trabajos que logran publicarse no suelen presentar diferencias significativas respecto a lo que se expuso previamente en el congreso.

© 2017 Publicado por Elsevier España, S.L.U. en nombre de SECOT.

Introduction

Presenting an investigative work as a scientific communication at a congress does not necessarily lead to its subsequent publication, since the process of publication requires effort and involvement by the authors. Lack of time on the authors' part for preparing the article appears to be the principal barrier to publishing a scientific communication in a journal.¹ Despite this, publication in indexed journals is still essentially the standard method of spreading information through the scientific community, since it leads to greater dissemination of the work undertaken and recognition for the authors.

The rate of publications of oral communications in specialised journals and the factor of impact obtained are a reflection of the scientific quality of congresses.² For this reason, some Spanish medical associations have published their publication rates, with highly variable results.³⁻⁸ These publications rates range from between 17% and 51% for podium presentations and from between 20% and 50% for those studies which include scientific posters and podium presentations in their analysis. In recent years there has been an increasingly greater presence of Spanish authors, in international reference data bases, which has been calculated as a 600% rise between 1980 and 2003.⁹

Bibliometric information on congresses in the specialty or sub-specialty of orthopaedic surgery or traumatology celebrated in the Spanish language is unknown. Different international orthopaedic and trauma associations have published publication rates of their congress,¹⁰⁻³² because interest in the rate of publications has increased. The publication rate varied from between 48% and 73% depending on the congress analysed and the year of its celebration. With regard to the sub-speciality of shoulder and elbow, only the publication rate of the shoulder and elbow sessions of the specialty congress of the American Academy of Orthopaedic Surgery (AAOS)¹² is known, and is set at 58%. In this article, the changes in the order and number of authors, in the study sample or in the title do not usually have an impact on the conclusions of publication.

What is the probability of publication of an abstract that has been accepted in the congress of the Spanish Society of Shoulder and Elbow Surgery (SECHC)? Is any factor influential in the publication rate? Our study proposes to offer the publication rate of the congress of the SECHC, analysing the factors which could affect whether a research study

is published afterwards. Concordance of content between published studies and their corresponding podium presentations previously presented in the congress was also analysed.

Material and methods

The abstracts of the oral communications presented in 3 consecutive congresses of the SECHC in 2007, 2009 and 2011 were obtained through the society's website (www.sechc.es) and included in this study. Communications were classified in accordance with the type of study (treatment, diagnosis, prognosis or basic sciences) and the level of scientific study (I-V). The studies were classified according to subject matter: elbow injuries, proximal humerus fractures, rotator cuff injury, degenerative disease of the shoulder/joint replacements, glenohumerus instability, collarbone injury/acromioclavicular articulation, basic or miscellaneous sciences.

A minimum period of 6 years was fixed to evaluate the survival of the review processes and publication of the articles, although the majority of communications which are published manage to be published within 4 years after the celebration of the congress.^{11,14,16,17,23,32}

All the abstracts obtained through the SECHC web site were investigated using the PubMed data in the month of June 2017, which aimed at identifying those abstracts which were published afterwards. The search was undertaken with the name of the main author, and the key words that appeared in the title of the abstract. In the cases where no publications were found with the before-mentioned criteria, a search was made through the remaining authors prior to declaring a work as not published. The criteria used to declare a abstract as published was based on previously described criteria.^{3,27}

A total of 122 oral presentation abstracts were included in the SECHC website and were classified according to a combination of characteristics in order to determine the probabilities of obtaining publication in indexed journals. The indexed journal where the works were published and the impacting factor at the time of publication were recorded for those abstracts which passed the review process of the indexed journal, using CiteFactor.³³

In statistical analysis, the Wilcoxon rank-sum non parametric test was used for comparing groups, and the Spearman correlation test for assessing associations of

Table 1 Comparison of podium presentations with articles published afterwards, for those presentations which obtained indexed publications.

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Cadaveric and three-dimensional computed tomography study of the morphology of the scapula with reference to reversed shoulder prosthesis	2007	Abad et al.	108	NA	Ratio glenoid neck length Ant view: short 18.25% Post view: 52%; 64%	Cadaveric and three-dimensional computed tomography study of the morphology of the scapula with reference to reversed shoulder prosthesis ⁴⁹	Torrens et al.	108	NA	Ratio glenoid neck length Res Ant view: short 18%; long 82% Post view: short 60%; long 40%	J Orthop Surg Res	19	1.33
Orthopaedic treatment of proximal humeral fractures proximal	2007	Corrales et al.	58	12	NA	Functional and quality-of-life results of displaced and nondisplaced proximal humeral fractures ⁵¹	Torrens et al.	70	24	Constant scale 4p 33.66 3p 54.64 2p 66.88 1p 71	J Orthop Trauma	55	2.13
Treatment of latissimus dorsi transfer for irreparable posterosuperior rotator cuff tear	2007	De Casas et al.	7	13	△ Constant scale = 26	Clinic and electro-tromiographic results of latissimus dorsi transfer for irreparable posterosuperior rotator cuff tear ³⁹	De Casas et al.	14	36	△ Constant scale = 26	J Orthop Res	92	1.55
Study of concordance and reproducibility in the assessment of radiolucent lines in cemented shoulder hemi-arthroplasties	2007	Torres et al.	32	NA	Kappa index Category 1: 0.62 Category 2: 0.40 Category 3: 0.50 Category 4: 0.25 Category 5: 0.02 Category 6: 0.25 Category 7: 0.57	Assessment of radiolucent lines in cemented shoulder hemi-arthroplasties: study of concordance and reproducibility ⁵⁰	Torrens et al.	32	NA	Kappa index Category 1: 0.65 Category 2: 0.51 Category 3: 0.60 Category 4: 0.21 Category 5: 0.60 Category 6: 0.48 Category 7: 0.62	Int Orthop	21	0.94

Table 1 (Continued)

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Radial head fractures Mason grade II, III and IV treated with resection arthroplasties. Long-term follow-up	2009	Iftimie et al.	27	204	DASH 4.89 MEPS 95% of results excellent or good	Resection arthroplasty for radial head fractures: long-term follow-up ⁴¹	Iftimie et al.	27	204	DASH 4.89 MEPS 96.4% of results excellent or good	J Shoulder Elbow Surg	22	2.74
Quality of life of patients with reverse shoulder arthroplasties. Conditioning factors	2009	Margarit Martin et al.	26	40	△ Constant scale = 32.6 Scapular erosion 33%	Retrospective study of scapular notches in reverse shoulder arthroplasties ⁵²	Torrens et al.	36	40	△ Constant scale = No erosion 26.87 Erosion 29.79 Scapular erosion 36%	Am J Orthop (Belle Mead NJ)	51	0.77
Nondisplaced proximal humeral fractures: high incidence among outpatient-treated osteoporotic fractures and severe impact on upper extremity function and patient subjective health perception	2009	Morcillo et al.	912	NA	DASH 26.62 Euro-QOL 67.3% have pain	Nondisplaced proximal humeral fractures: high incidence among outpatient-treated osteoporotic fractures and severe impact on upper extremity function and patient subjective health perception ³⁷	Calvo et al.	912	NA	DASH 26.62 EuroQOL 67.3% have pain	J Shoulder Elbow Surg	24	2.74

Table 1 (Continued)

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Repair of type II SLAP lesion through a single anterior portal	2009	Rodríguez Vellando et al.	23	15	UCLA 91%	Vertical versus horizontal suture configuration for the repair of isolated type II SLAP lesion through a single anterior portal: a randomised controlled trial ⁴⁷	Silberberg et al.	32	37	ASSE 91.9 vs. ASSE 95.8	Arthroscopy	33	3.02
Shoulder dislocation in patients older than 60 years of age	2009	Rapariz	29	84	Recurrence 31.03%	Shoulder dislocation in patients older than 60 years of age ⁴⁴	Rapariz et al.	29	72	Recurrence 31.03%	Int J Shoulder Surg	19	-
The pectoralis major tendon as a reference for restoring humeral length and retroversion with hemi-arthroplasty for fracture	2009	Vilà et al.	20	NA	Distance pect. major-humeral head: 5.64 cm Distance insertion of pect major-posterior implant flange: 1.06 cm Pect major-implant flange angle: 24.65	The pectoralis major tendon as a reference for restoring humeral length and retroversion with hemi-arthroplasty for fracture ⁴⁸	Torrens et al.	20	NA	Distance pect. major-humeral head: 5.64 cm Distance insertion of pect major-posterior implant flange: 1.06 cm Pect major-implant flange angle: 24.65	J Shoulder Elbow Surg	-4	1.93

Table 1 (Continued)

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Functional outcomes after arthroscopic repair of massive rotator cuff tears: influence of the application of platelet-rich fibrin	2011	Antuña et al.	28	12	△ Constant scale PRP 26 No PRP 30	Platelet-rich fibrin in arthroscopic repair of massive rotator cuff tears: a prospective randomised pilot clinical trial ³⁵	Antuña et al.	28	12	△ Constant scale PRP 26 NO PRP 30	Acta Orthop Belg	21	0.63
Clinical outcome and prognostic factors of revision arthroscopic rotator cuff tear repair	2011	Calvo et al.	40	12	△ Constant scale 15.2	Clinical outcome and prognostic factors of revision arthroscopic rotator cuff tear repair ⁵⁴	Valencia et al.	51	25	△ Constant scale 26.42 △ Simple Shoulder Test 3,82	Knee Surg Sports Trauma Arthrosc	68	3.23
Medium term results of unipolar modular radial head arthroplasty	2011	Cecilia et al.	32	30	MEPS 84% excellent or good results; 3 cases of compromised radial nerve 3 cases of capitellum erosion 1 luxation 1 incomplete resection	Medium term results of unipolar modular radial head arthroplasty ⁴³	Martín Fuentes et al.	44	21	MEPS 82% excellent or good results; 2 loosening	Rev Esp Cir Ortop Traumatol	26	
Systematic review after 6 years of olecranon fractures. Clinical and functional results.	2011	Cervera et al.	93	36	87.4% Good or excellent Range of motion >50	Treatment of comminuted olecranon fractures with olecranon plate and structural iliac crest graft ³⁸	Cervera et al.	3	23	100% good or excellent Flexion 115 Extension -20 Pronation 71 Supination 80	Acta Orthop Belg	21	0.63

Table 1 (Continued)

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Use of adipose- derived stem cells in an experimental rotator cuff fracture	2011	Encinas et al.	56	2	No difference in tensile strength at 4 and 8 weeks	Use of adipose- derived stem cells in an experimental rotator cuff fracture animal model ³⁶	Barco et al.	44	2	No difference in tensile strength at 4 and 8 weeks	Rev Esp Cir Ortop Traumatol	42	0.09
Progression of deformity during fracture healing of humeral fractures treated conservatively	2011	Foruria et al.	106	12	↑ retroversion of the head 7.5 18.75% progression of greater tuberosity	Proximal humeral fractures treated conservatively settle during fracture healing ⁴⁰	Foruria et al.	89	12	↑ retroversion of the head 7.1 Trauma <20% progression of greater tuberosity	J Orthop Trauma	39	0.92
Treatment of fracture sequelae of the proximal humerus: comparison of hemiarthroplasty and reverse total shoulder arthroplasty.	2011	Isart et al.	14	32	HA Group △ Constant scale 18.16 PI Group △ Constant scale 22.19	Treatment of fracture sequelae of the proximal humerus: comparison of hemiarthroplasty and reverse total shoulder arthroplasty ³⁴	Alentorn Geli et al.	32	50	HA Group △ Constant scale 14.6 PI Group △ Constant scale 26.6	Arch Orthop Trauma Surg	41	2.33
Morbidity and mortality of surgically treated proximal humerus fractures	2011	Isart et al.	97	96	Autonomy 79.5% Mortality 14%	Morbidity and mortality of surgically treated proximal humerus fractures ⁴²	Isart et al.	94	96	Autonomy 79.5% Mortality 18.6%	Rev Esp Cir Ortop Traumatol	40	0.09

Table 1 (Continued)

Title of the presentation	Year	Author	No	FU	Results	Title of the published article	Author	No	FU	Results	Revista	Demora	FI
Plasma rich in growth factors (PRGF) in arthroscopic rotator cuff repair: a randomised, clinical trial with 69 patients	2011	Molano Muñoz et al.	69	12	PRFC Group 78% satisfaction Control group 77% satisfaction	Plasma rich in growth factors (PRGF) in arthroscopic rotator cuff repair: A randomised, double-blind, controlled clinical trial ⁴⁵	Ruiz Moneo et al.	69	12	PRGF group △UCLA: 8.3 Control group △UCLA: 9	Arthroscopy	22	3.29
Decreased axial coracoid inclination angle is associated with rotator cuff tears	2011	Sánchez et al.	82	NA	Rotator group Angle 1 (A1) 57.06 Angle 2 (A2) 49.17 Control group: A1: 132.52 A2: 144.34	Decreased axial coracoid inclination angle is associated with rotator cuff tears ⁵³	Torrens et al.	100	NA	Rotator Group Angle 1: 49.7 Angle 2: 76.45 Control group A1: 132.33 A2: 144.34	J Orthop Surg (Hong Kong)	47	0.77
Treatment of complex proximal humeral fractures in elderly patients with reverse shoulder arthroplasty versus hemi-arthroplasty. Prospective study	2011	Sebastiá-Forcada et al.	32	22.7	△ Anterior flexion 64 △ abduction 63	Reverse shoulder arthroplasty versus hemi-arthroplasty for acute proximal humeral fractures. A blinded, randomised, controlled, prospective study ⁴⁶	Sebastiá-Forcada et al.	62	28.5	△ Anterior flexion 41 △ abduction 34,2	J Shoulder Elbow Surg	43	2.75

IF, impact factor of the journal; FU, follow-up in months; No, sample size, NA, not applicable; NA, not available △, pre and postoperative differences.

Table 2 Publication rate of Spanish congresses.

Congress	Year	Overall PR in %	PR of podium presentations in %	PR of poster in %	Observation period (minimum) in years
Spanish Society of Cardiology ³	2002/2005/2008	NR	38.3	NR	4
Spanish Society of Anaesthesia and Resuscitation ⁴	1992	NR	17.1	NR	6
AEEH ⁵	1999	52.8	71.4	54	4
Spanish Society of Radiology ⁶	1994–1998	15	18	13	5
Spanish Pharmacology Congress ⁷	1994–1998	26	NR	NR	5
Spanish Society of Rheumatology ⁸	1996	21	45	15	5

NR, not reported; PR, publication rate.

continuous variables which followed a non-normal distribution. The chi-square test or the exact Fisher test was used to assess the association between dichotomous variables. 95% confidence intervals were calculated over the estimators. The R version 3.4.0 for Windows was the statistical package used to treat data and for analysis.

Results

Out of a total of 122 abstracts examined, 28 corresponded to elbow injury, 27 to proximal humerus fractures, 24 to rotator cuff injury, 19 to basic sciences or miscellaneous, 13 to glenohumeral instability, 7 to degenerative disease of the shoulder and 4 to collarbone or acromioclavicular injury. During the congress celebrated in 2007 30 oral communications were presented, 43 in the 2009 congress, and 49 in the 2011 congress.

With regard to the type of works presented at the SECHC congress, 87 abstracts corresponded to treatment assessment presentations (intervention); 15 to assessment of prognostic factors; 11 to assessment of diagnostic methods; 8 to basic sciences, whilst only one presentation consisted of a review of the literature. With regard to the level of scientific evidence, we found one presentation of level I, 9 of level II, 17 of level III, 85 of level IV, 2 of level V, whilst it was not possible to classify 8 by level of evidence. The mean sample of communications presented was of 59.66 patients per presentation (4–912), with mean follow-up of 27.06 (1–240) months.

Twenty one of the 122 oral communications assessed by the SECHC (17.21%) presented in the congresses were published in indexed journals, in accordance with the applied criteria.^{34–54} The 21 articles were published in 12 different journals, with a mean impact factor of 1.51 (0.09–3.29). The mean time which elapsed between the presentation of the oral communication and subsequent publication was 36.71 months (4–92 months). 4 communications out of 30 oral communications (13.6%) were published in reference to the 2007 congress, 6 out of 43 communications (13.95%) in reference to the 2009 congress, whilst of the 49 communications presented in 2011, 11 of them successfully had a publication in an indexed journal (22.45%). The tendency to

growing publication between congresses did not obtain any statistical significance ($p=0.29$).

Fifteen of the 21 articles published showed differences in the sample size (4 of the 15 reported lower samples, whilst higher samples were reported in 11 of the 15 samples included in the final presentation). With regards to follow-up, 3 of the 21 studies published lower follow-ups whilst 4 of the published presentations reported a higher follow-up than that presented at the SECHC congress. Nine of the 21 studies reported the same follow-up whilst 5 did not apply follow-up due to the study design. In 8 out of the 21 works published the results were the same as those previously expressed in the SECHC congress, whilst in 12 works the results were different. In one single work they could not obtain results from the oral communication in the SECHC (Table 1) website. Despite this, only one out of the 21 works published modified the conclusions. In this study,⁴⁷ the focus of the published work differed from the podium presentation. The changes in the sample, follow-up and results were not statistically significant ($p=0.26$; $p=0.17$ and $p=0.36$, respectively).

The publication rate was not significantly related to the study subject matter ($p=0.30$), neither were the type of study, level of evidence, sample included or follow-up of podium presentations determining factors of the probability of publication ($p=0.35$; $p=0.96$; $p=0.19$ and $p=0.17$, respectively). Lastly, delay in publication did not affect the impact factor which was obtained with regards to the indexed journal ($p=0.92$).

Discussion

The SECHC congress is celebrated biannually and is the benchmark state congress in Spain in shoulder and elbow disorders. This paper shows that at least 20% of the podium presentations are published in indexed journals.

The publication rate of different medical congresses celebrated in Spain^{3–8} is very varied (Table 2), and the majority are based on congresses celebrated during the last century (1992–1999). The Spanish Association for the Study of the Liver achieved the best publication rate of a Spanish Medical congress,⁵ with 71.4% of its podium presentations and

54% of its posters published in journals four years after the congress (1999). Only the Spanish Society of Cardiology³ has published results from congresses celebrated between 2002 and 2008, and has reported 38.4% as its publication rate of podium presentations at 4 years. In the sub-speciality of shoulder and elbow we only have knowledge of the publication rates of podium presentations in congress sessions of the AAOS, as being 66% after a minimum follow-up of 3 years.¹²

The reasons for the low SECHC publication rate found may differ. The fact that the SECHC congress is a sub-specialty of orthopaedic surgery and trauma (specialty) could explain the differences with regard to the publications rates mentioned for Spanish congresses. Different publications have proposed reasons for explaining the non publication of podium presentations. Sprague et al.¹ studied the reasons for non publications of abstracts presented at the AAOS congress. Out of the 306 works for which no publication was found, 199 authors responded to the survey. Out of the 199 authors, 72 (36.2%) declared that the work had been published whilst 71 authors (35.7%) had never sent off their presentations. The major obstacles stated by the authors for not sending off their scientific studies was lack of time¹ and lack of interest in publishing. Fear of rejection was another reason stated for not sending off a podium presentation for publication.⁵⁵

The majority of studies presented at the SECHC congress which were later published suffered from changes over the publication process. In our study, half of the works published result in a higher sample, whilst almost 25% reduce the sample in later publication (Table 1). The works by De Casas³⁹ and Sebastiá-Forcada⁴⁶ are the ones with a larger sample, with the patient numbers increasing by 100%. Bhandari et al.¹⁰ reported an almost 10% reduction of the sample in the works published compared to the abstract given at the AAOS congress, whilst Preston et al.⁵⁵ observed almost a 20% reduction in the sample of the studies previously expressed in the congress of the Orthopaedic Trauma Association. With regard to follow-up, all the studies except 3^{38,43,45} maintained the same follow-up time or increased it considerably. As an example, the work published by Torrens et al.⁵¹ changed from a 12 month follow-up recorded in the congress abstract to a 24 month follow-up in subsequent publication. De Casas³⁹ practically tripled follow-up from podium presentation to publication. With regard to the conclusions, only the work of Silberberg⁴⁷ showed different conclusions between podium presentation and publication. In this case, the podium presentation expressed the results of a surgical technique (SLAP injury repair through anterior portal vein), whilst the subsequent publication extended the study with a comparative work between 2 specific techniques. These findings appear to be in keeping with other publications which study the consistency between abstracts and posterior publications. The work of Preston et al.⁵⁵ delves into the differences between the podium presentations from the congress of the Orthopaedic Trauma Association between the years of 1994 and 1997. The authors of this study found that there was a 93.4% coincidence rate in the conclusions between congress abstracts and later publications. 56.3% of the works which were published used the same sample as in the previous podium presentation whilst 25.5% increased it. The reasons stated for changing the presented data were diverse. Often podium presentations were the preliminary

results of an investigation with regards to follow-up and sample size. Only 10% of the studies which contained changes to the content of the work offer an explanation, according to Preston et al.⁵⁵ Among them, the loss of follow-up of the patients included in the study was the main reason for change. Bandhari et al.,¹⁰ in contrast, identified the delay in publication as the only predisposing factor between the oral communication at a congress and subsequent publication.

The study topic was not related to the probability of gaining a posterior publication in our analysis. Schulte et al.²⁷ found differences in the publication rate according to the work design (prospective vs retrospective) and the level of scientific evidence (I-II vs III-IV) in the study of the German orthopaedics congress, but there were no differences in the study topic. Schulte et al.²⁷ and Kleine-Konig et al.²² found that the abstracts corresponding to works with a significant main outcome were more likely to be published than those with clearly non significant outcomes. Furthermore, multi-centre studies and biomechanical studies²⁷ obtained higher publication rates. Cifuentes et al.⁵ identified the type of presentation (oral or poster) as the only determining factor for final publication of an abstract presented at a meeting of the Spanish Association of the Study of Liver Diseases.

Scientific meetings in Spain, such as the Electrophysiology, Arrhythmias and cardiac stimulation meeting have strictly limited the time for oral presentations. Only those presentations which are deserving of awards due to their originality or level of scientific evidence are presented at the congress. This focus could help to improve the number of studies published which are presented previously in SECHC congresses. Another measure which would help to improve the rate of publication of the SECHC would be that or promoting the study of high levels of evidence: prospective randomised studies, multicentre or biomechanical studies, as has been determined by other authors.²⁷ Consolidation of the rising trend in the SECHC publication rate is expected, like that observed in this study in the 2007–2011 period due to the scientific development of society itself. Publications in Spain in the Science Citation Index Expanded data base increased by over 600% between 1980 and 2003, a far higher rise than the total increase of the data bases (100%).⁹ Despite the research internationalisation process and the drop in publications in the Spanish language,⁹ the indexation of the SPANISH JOURNAL OF ORTHOPAEDIC SURGERY AND TRAUMATOLOGY in Medline/Pubmed from 2013 will help Spanish authors to familiarise themselves with a publication in their own language. The Spanish Society of Cardiology has identified³ the “Revista Española de Cardiología” as the main publication for authors of works presented at their congresses. 37.1% of the works published from the congress were present in this journal.

This study has several limitations. Firstly, the reasons for non publication of communications have not been analysed nor is the number of studies which initiated some type of review process in indexed journals available. Secondly, the search for posterior publications through PubMed may lead to an underestimation of the analysed publication rate.

Conclusions

The majority of oral communications presented in the SECHC congresses are not published in indexed journals. The dif-

ferences between the published communications and their previous oral presentations do not appear to be significant.

Level of evidence

Level of evidence IV (systematic review).

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments have been performed on humans or animals in this research.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflict of interests

The authors have no conflict of interests to declare.

References

1. Sprague S, Bhandari M, Devereaux PJ, Swiontkowski MF, Tornetta P 3rd, Cook DJ, et al. Barriers to full-text publication following presentation of abstracts at annual orthopaedic meetings. *J Bone Joint Surg Am.* 2003;85:158–63.
2. Gormann RL, Olerda GM. Publication of presented abstracts at annual scientific meetings: a measure of quality. *Vet Hum Toxicol.* 1990;32:470–2.
3. Alonso-Arroyo A, Aleixandre-Benavent R, Vidal-Infer A, Anguita-Sánchez M, Chorro-Gascó FJ, Bolaños-Pizarro M, et al. Executive Committee of the Spanish Society of Cardiology. Subsequent full publication of abstracts presented in the annual meetings of the Spanish Society of Cardiology. *Rev Esp Cardiol.* 2014;67:15–21, <http://dx.doi.org/10.1016/j.recesp.2013.05.010>.
4. Castillo J, García-Guasch R, Cifuentes I. Publicaciones derivadas de las comunicaciones libres del XX Congreso de la Sociedad Española de Anestesiología y reanimación (Anestesia 92). *Rev Esp Anestesiol Reanim.* 2000;47:53–6.
5. Cifuentes I, Bartolí Solé R, Miquel Planas M, Cabré Gelad E, Morillas Cunill RM, Planas Vila R. How many reach their goal? Outcome of abstracts presented at the XXIV AAEEH (Spanish Liver Study Association) Congress and their progress over time. *Gastroenterol Hepatol.* 2007;30:263–7.
6. Miguel-Dasit A, Martí-Bonmatí L, Aleixandre R, Sanfeliu P, Valderrama JC. Publications resulting from Spanish radiology meeting abstracts: which, where and who. *Scientometrics.* 2006;66:467–80.
7. Montané E, Vidal X. Fate of the abstracts presented at three Spanish clinical pharmacology congresses and reasons for unpublished research. *Eur J Clin Pharmacol.* 2007;63:103–11, <http://dx.doi.org/10.1007/s00228-006-0235-7>.
8. Olivé A, Cifuentes I, Vázquez Corbacho D, Roca J. Rumbo a lo desconocido: Destino final de los resúmenes presentados a un congreso de reumatología. *Rev Esp Reumatol.* 2004;31:122–5.
9. Bordons M. Hacia el reconocimiento de las publicaciones científicas españolas. *Rev Esp Cardiol.* 2004;57:799–802.
10. Bhandari M, Devereaux PJ, Guyatt GH, Cook DJ, Swiontkowski MF, Sprague S. An observational study of orthopaedic abstracts and subsequent full-text publications. *J Bone Joint Surg Am.* 2002;84:615–21.
11. Daluiski A, Kuhns CA, Jackson KR, Lieberman JR. Publication rate of abstracts presented at the annual meeting of the Orthopaedic Research Society. *J Orthop Res.* 1998;16:645–50.
12. DeMola PM, Hill DL, Abboud JA. Publication rate of abstracts presented at the shoulder and elbow session of the American Academy of Orthopaedic Surgery. *Clin Orthop Relat Res.* 2009;467:1629–33, <http://dx.doi.org/10.1007/s11999-008-0474-2>.
13. Donegan DJ, Kim TW, Lee GC. Publication rates of presentations at the Annual Meeting of the American Academy of Orthopaedic Surgeons. *Clin Orthop Relat Res.* 2010;468:1428–35, <http://dx.doi.org/10.1007/s11999-009-1171-5>.
14. Eck JC. Publication rates of abstracts presented at Biennial Meetings of the International Society of Arthroscopy, Knee Surgery and Orthopaedic Sports Medicine. *Knee Surg Sports Traumatol Arthrosc.* 2005;13:426–9, <http://dx.doi.org/10.1007/s00167-004-0559-8>.
15. Gavazza JB, Foulkes GD, Meals RA. Publication pattern of papers presented at the American Society for Surgery of the Hand Annual Meeting. *J Hand Surg Am.* 1996;21:742–5.
16. Hamlet WP, Fletcher A, Meals RA. Publication patterns of papers presented at the Annual Meeting of the American Academy of Orthopaedic Surgeons. *J Bone Joint Surg Am.* 1997;79:1138–43.
17. Harris IA, Mourad MS, Kadir A, Solomon MJ, Young JM. Publication bias in papers presented to the Australian Orthopaedic Association Annual Scientific Meeting. *ANZ J Surg.* 2006;76:427–31, <http://dx.doi.org/10.1111/j.1445-2197.2006.03747.x>.
18. Lloyd EW, Geller JA, Iorio R, Yoon RS, Huo MM. Publication rates of scientific presentations at the American Association of Hip and Knee Surgeons annual meetings from 1996 to 2001. *J Arthroplasty.* 2006;21:2–5, <http://dx.doi.org/10.1016/j.arth.2006.04.021>.
19. Jackson KR, Daluiski A, Kay RM. Publication of abstracts submitted to annual meeting of the Pediatric Orthopaedic Society of North America. *J Pediatr Orthop.* 2000;20:2–6.
20. Jasko JJ, Wood JH, Schwartz HS. Publication rates of abstracts presented at annual musculoskeletal tumor society meetings. *Clin Orthop Relat Res.* 2003;415:98–103, <http://dx.doi.org/10.1097/01.blo.0000093902.12372.76>.
21. Kinsella SD, Menge TJ, Andersons AF. Publication rates of podium versus poster presentations at the American Orthopaedic Society for Sports Medicine meetings: 2006–2010. *Am J Sports Med.* 2015;43:1255–9, <http://dx.doi.org/10.1177/0363546515573939>.
22. Kleine-König MT, Schulte TL, Gosheger G, Rödl R, Schiedel FM. Publication rate of abstracts presented at European Pediatric Orthopaedic Society Annual Meetings, 2006 to 2008. *J Pediatr Orthop.* 2014;34:e33–8, <http://dx.doi.org/10.1097/BPO.0000000000000140>.
23. Kleweno CP, Bryant WK, Jacir AM, Levine WN, Ahmad CS. Discrepancies and rates of publication in orthopaedic sports medicine abstracts. *Am J Sports Med.* 2008;36:1875–9, <http://dx.doi.org/10.1177/0363546508319054>.
24. Kwong Y, Kwong FN, Patel J. Publication rate of trauma abstracts presented at an international orthopaedic conference. *Injury.* 2007;38:745–9, <http://dx.doi.org/10.1016/j.injury.2006.07.002>.
25. Murray DB, Wright RW, Seiler JG 3rd, Day TE, Schwartz HS. Publication rates of abstracts presented at the 1993 annual Academy meeting. *Clin Orthop Relat Res.* 1999;359:247–53.
26. Preston CF, Bhandari M, Fulkerson E, Ginat D, Koval KJ, Egol KA. Podium versus poster publication rates at the Orthopaedic Trauma Association. *Clin Orthop Relat Res.* 2005;437:260–4, <http://dx.doi.org/10.1097/01.blo.0000167667.80325.61>.
27. Schulte TL, Trost M, Osada N, Huck K, Lange T, Gosheger G, et al. Publication rate of abstracts presented at the Annual Congress of the German Society of Orthopaedics and

- Trauma Surgery. Arch Orthop Trauma Surg. 2012;132:271–80, <http://dx.doi.org/10.1007/s00402-011-1398-y>.
28. Schulte TL, Huck K, Osada N, Trost M, Lange T, Schmidt C, et al. Publication rate of abstracts presented at the Annual Congress of the Spine Society of Europe (years 2000–2003). Eur Spine J. 2012;21:2105–12, <http://dx.doi.org/10.1007/s00586-012-2211-5>.
29. Wang JC, Yoo S, Delamarter RB. The publication rates of presentations at major Spine Speciality Society meetings (NASS, SRS, ISSLS). Spine (Phila Pa 1976). 1999;24:425–7.
30. Yoo S, Oh G, Wang JC. Publication rates of presentations made at annual meetings of the American Orthopaedic Society for Sports Medicine and the Arthroscopy Association of North America. Am J Orthop (Belle Mead NJ). 2002;31:367–9.
31. Yoon RS, Lloyd EW, McGrory B, Bal BS, Macaulay W. Studies presented in poster format at the annual meetings of the American Association of Hip and Knee Surgeons: how do they fare in the peer review process? J Arthroplasty. 2007;22:17–20, <http://dx.doi.org/10.1016/j.arth.2007.03.010>.
32. Wang JC, Yoo S, Delamarter RB. The publication rates of presentations at major Spine Speciality Society meetings (NASS, SRS ISSLS). Spine (Phila Pa 1976). 1999;24:425–7; Weber EJ, Callahan ML, Wears RL, Barton C, Young G. Unpublished research from a medical speciality meeting: why investigators fail to publish. JAMA. 1998;280:257–9.
33. Cite Factor. Available from: <http://www.citefactor.org/journal-impact-factor-list-2014.html> [accessed 13.06.17].
34. Alentorn-Geli E, Guiro P, Santana F, Torrens C. Treatment of fracture sequelae of the proximal humerus: comparison of hemiarthroplasty and reverse total shoulder arthroplasty. Arch Orthop Trauma Surg. 2014;134:1545–50, <http://dx.doi.org/10.1007/s00402-014-2074-9>.
35. Antuña S, Barco R, Martínez Diez JM, Sánchez Márquez JM. Platelet-rich fibrin in arthroscopic repair of massive rotator cuff tears: a prospective randomized pilot clinical trial. Acta Orthop Belg. 2013;79:25–30.
36. Barco R, Encinas C, Valencia M, Carrascal MT, García-Arranz M, Antuña S. Use of adipose-derived stem cells in an experimental rotator cuff fracture animal model. Rev Esp Cir Ortop Traumatol. 2015;59:3–8, <http://dx.doi.org/10.1016/j.recot.2014.07.00>.
37. Calvo E, Morcillo D, Foruria AM, Redondo-Santamaría E, Osorio-Picorne F, Caeiro JR, GEIOS-SECOT Outpatient Osteoporotic Fracture Study Group. Nondisplaced proximal humeral fractures: high incidence among outpatient-treated osteoporotic fractures and severe impact on upper extremity function and patient subjective health perception. J Shoulder Elbow Surg. 2011;20:795–801, <http://dx.doi.org/10.1016/j.jse.2010.09.008>.
38. Cervera-Irimia J, Tomé-Bermejo F, Gómez-Bermejo MA, Holgado-Moreno E, Stratenwerth EG. Treatment of comminuted olecranon fractures with olecranon plate and structural iliac crest graft. Acta Orthop Belg. 2012;78:703–7.
39. De Casas R, Lois M, Cidoncha M, Valadron M. Clinic and electromyographic results of latissimus dorsi transfer for irreparable posterosuperior rotator cuff tears. J Orthop Surg Res. 2014;9:83–96, <http://dx.doi.org/10.1186/s13018-014-0083-6>.
40. Foruria AM, Martí M, Sánchez-Sotelo J. Proximal humeral fractures treated conservatively settle during fracture healing. J Orthop Trauma. 2015;29:24–30, <http://dx.doi.org/10.1097/BOT.0000000000000244>.
41. Iftimie PP, Calmet Garcia J, de Loyola Garcia Forcada I, Gonzalez Pedrouzo JE, Giné Gomà J. Resection arthroplasty for radial head fractures: long-term follow-up. J Shoulder Elbow Surg. 2011;20:45–50, <http://dx.doi.org/10.1016/j.jse.2010.09.005>.
42. Isart A, Sánchez JF, Santana F, Puig L, Cáceres E, Torrens C. Morbidity and mortality of surgically treated proximal humerus fracture. Rev Esp Cir Ortop Traumatol. 2014;58:223–8, <http://dx.doi.org/10.1016/j.recot.2014.02.003>.
43. Martín-Fuentes AM, Cecilia-López D, Resines-Erasun C. Midterm results of unipolar modular radial head arthroplasty. Rev Esp Cir Ortop Traumatol. 2013;217–23, <http://dx.doi.org/10.1016/j.recot.2013.02.005>.
44. Rapariz JM, Martín-Martín S, Pareja-Bezares A, Ortega-Klein J. Shoulder dislocation in patients older than 60 years of age. Int J Shoulder Surg. 2010;4:88–92, <http://dx.doi.org/10.4103/0973-6042.79792>.
45. Ruiz-Moneo P, Molano-Muñoz J, Prieto E, Algorta J. Plasma rich in growth factors in arthroscopic rotator cuff repair: a randomized, double-blind controlled clinical trial. Arthroscopy. 2013;29:2–9, <http://dx.doi.org/10.1016/j.arthro.2012.08.014>.
46. Sebastiá-Forcada E, Cebrian-Gómez R, Lizaur-Utrilla A, Gil-Guillem V. Reverse shoulder arthroplasty versus hemiarthroplasty for acute proximal humeral fractures. A blinded, randomized, controlled, prospective study. J Shoulder Elbow Surg. 2014;23:1419–26, <http://dx.doi.org/10.1016/j.jse.2014.06.035>.
47. Silberberg JM, Moya-Angeler J, Martin E, Leyes M, Forriol F. Vertical versis horizontal suture configuration for the repair of isolated type II SLAP lesion through a single anterior portal: a randomized controlled trial. Arthroscopy. 2011;27:1605–13, <http://dx.doi.org/10.1016/j.arthro.2011.07.009>.
48. Torrens C, Corrales M, Melendo E, Solano A, Rodríguez-Baeza A, Cáceres E. The pectoralis major tendon as a reference for restoring humeral length and retroversion with hemiarthroplasty for fracture. J Shoulder Elbow Surg. 2008;17:947–50, <http://dx.doi.org/10.1016/j.jse.2008.05.041>.
49. Torrens C, Corrales M, González G, Solano A, Cáceres E. Cadaveric and three-dimensional computed tomography study of the morphology of the scapula with reference to reversed shoulder prosthesis. J Orthop Surg Res. 2008;3:49, <http://dx.doi.org/10.1186/1749-799X-3-49>.
50. Torrens C, Martínez-Díaz S, Ruiz A, Ginés A, Cáceres E. Assessment of radiolucent lines in cemented shoulder hemi-arthroplasties: study of concordance and reproducibility. Int Orthop. 2009;33:165–9, <http://dx.doi.org/10.1007/s00264-007-0452-4>.
51. Torrens C, Corrales M, Vilà G, Santana F, Cáceres E. Functional and quality-of-life results of displaced and nondisplaced proximal humeral fractures treated conservatively. J Orthop Trauma. 2011;25:581–7, <http://dx.doi.org/10.1097/BOT.0b013e318210ed2>.
52. Torrens C, Santana F, Picazo B, Cáceres E. Retrospective study of scapular notches in reverse shoulder arthroplasties. Am J Orthop (Belle Mead NJ). 2013;42:362–5.
53. Torrens C, Alentorn-Geli E, Sanchez JF, Isart A, Santana F. Decreased axial coracoid inclination angle is associated with rotator cuff tears. J Orthop Surg (Hong Kong). 2017;25:1–5, <http://dx.doi.org/10.1177/2309499017690329>.
54. Valencia Mora M, Morcillo Barrenechea D, Martín Ríos MD, Foruria AM, Calvo E. Clinical outcome and prognostic factors of revision arthroscopic rotator cuff repair. Knee Surg Sports Traumatol Arthrosc. 2016, <http://dx.doi.org/10.1007/s00167-016-4392-7>.
55. Preston CF, Bhandari M, Fulkerson E, Ginat D, Egol KA, Koval KJ. The consistency between scientific papers presented at the Orthopaedic Trauma Association and their subsequent full-text publication. J Orthop Trauma. 2006;20:129–33, <http://dx.doi.org/10.1097/01.bot.0000199120.45982.41>.