

Image of the month

Bouveret Syndrome: Variant of Gallstone Ileus[☆]**Síndrome de Bouveret: variante del íleo biliar**Pablo Calvo Espino,^{*} Arturo García Pavía, Mariano Artés Caselles, Víctor Sánchez Turrión

Servicio de Cirugía General y Aparato Digestivo, Hospital Universitario Puerta de Hierro Majadahonda, Majadahonda, Madrid, Spain



Fig. 1



Fig. 2

The patient is an 86-year-old woman with no relevant medical history who came to our emergency department complaining of abdominal pain, nausea and vomiting. On examination, she presented diffuse tenderness with no peritoneal signs. An abdominal CT scan demonstrated an image consistent with a 3.5 cm calculus in the distal segment of the horizontal portion of the duodenum and notable aerobilia (Figs. 1 and 2). Suprainfraumbilical midline laparotomy was performed, which revealed the stone distal to the angle of Treitz. Enterotomy was carried out with removal of the calculus.

[☆] Please cite this article as: Calvo Espino P, García Pavía A, Artés Caselles M, Sánchez Turrión V. Síndrome de Bouveret: variante del íleo biliar. Cir Esp. 2014;92:e3.

^{*} Corresponding author.

E-mail address: pcalvoespino@gmail.com (P. Calvo Espino).

2173-5077/\$ – see front matter © 2013 AEC. Published by Elsevier España, S.L. All rights reserved.