



Image of the month

Osteoarthritis Secondary to Rives–Stoppa Incisional Hernia Repair[☆]



Osteoarthritis secundaria a eventroplastia tipo Rives-Stoppa

Irene Mirón Fernández,^{*} Cristina Rodríguez Silva, José Rivas Becerra, Julio Santoyo Santoyo

Servicio de Cirugía General y del Aparato Digestivo y Trasplantes, Complejo Regional Universitario de Málaga, Málaga, Spain




Fig. 1

Clinical Case

Women of 65 years old, operated of a caesarean delivery and a hysterectomy and double adnexectomy some years later. She was operated in a programmed way of a laparotomic eventration with Rives–Stoppa technique. The patient came to Emergency after 45 days of the surgery, referring hypogastric pain, fever and functional problems with the movement of both legs. The blood test showed infection signs so a CT scan was made. It was found some lytic injuries in the pubic symphysis probably related with an osteoarthritis. The patient was in hospital for two weeks with potent antibiotics and after that she was discharged with improvement of the infection (Fig. 1). Diagnosis: Septic osteoarthritis due to Rives–Stoppa intervention.

Acknowledgments

To the Surgery and Radiodiagnostic Services for their collaboration.

[☆] Please cite this article as: Mirón Fernández I, Rodríguez Silva C, Rivas Becerra J, Santoyo Santoyo J. Osteoarthritis secundaria a eventroplastia tipo Rives-Stoppa. Cir Esp. 2020;98:43.

^{*} Corresponding author.

E-mail address: irenemironfernandez@gmail.com (I. Mirón Fernández).