

Scientific letters

CIRUGÍA ESPAÑOLA

www.elsevier.es/cirugia



Digestive symptoms and COVID-19: Importance of ruling out associated surgical pathology $\stackrel{\scriptscriptstyle{?}}{\sim}$



Sintomatología digestiva y COVID-19: importancia de descartar patología quirúrgica asociada

The Severe Acute Respiratory Syndrome-Corona Virus-2 (SARS-CoV-2) outbreak that was initially detected in China was declared a pandemic by the World Health Organization on March 11, 2020. The international underestimation of the problem, together with the high transmissibility of the virus and the lack of resources, caused an unprecedented situation worldwide. This led to the saturation of national healthcare systems, and Spain was one of the most affected countries. Some of the most common manifestations of the disease produced by the SARS-CoV-2 virus, known as COVID-19 (Coronavirus Disease 2019), include fever, dry cough, dyspnea, myalgias, fatigue, lymphopenia, elevated acute phase reactant levels and evidence of pneumonia on imaging studies.¹

Due to the saturation of the healthcare system, hospitals and medical professionals have had to reorganize and adapt to be able to provide care for the high number of infected patients.² With regard to urgent surgical pathology, there was a notable decrease in patients, while patients presented with more advanced disease.³

It is interesting to understand how the coexistence of SARS-CoV-2 infection can affect any type of acute abdominal pathology. Since many patients with COVID-19 disease present digestive symptoms,⁴ these can mask underlying surgical pathology. We present the case of a patient with SARS-CoV-2 infection and complicated acute appendicitis.

The patient is a 52-year-old woman with a history of dyslipidemia and hypertension, who came to the emergency room due to epigastric pain associated with fever, severe asthenia, ageusia, anosmia and loss of appetite over the past 7 days. Upon physical examination, she presented generalized abdominal pain, mainly in the right hemiabdomen, with no signs of generalized peritoneal irritation.

Given the epidemiological situation and compatible symptoms, a chest X-ray was performed in the emergency department as part of the pandemic protocol, which showed bilateral opacities compatible with COVID-19. Laboratory tests showed leukocytosis with neutrophilia and lymphopenia and increased lactate dehydrogenase 270 IU/L, C-reactive protein (CRP) 291.1 mg/L, gamma-glutamyl transferase 265 IU/L, ferritin (663 ng/mL) and D-dimer (2.422 ng/mL). Given the findings, a nasopharyngeal RT-PCR for SARS-CoV-2 was ordered, which was positive, and the patient was hospitalized in the Internal Medicine ward. After 2 days, and given the persistence of abdominal pain, a CT scan was ordered, which revealed complicated acute appendicitis with an inflammatory mass measuring 6.2×7.3×6 cm and radiological signs of moderate COVID-19 lung involvement (Fig. 1). Given the findings, the patient was evaluated by General Surgery (2 days after admission). The case was discussed in the Interventional Radiology service to assess the placement of a percutaneous drain, but this was ruled out because the collection was not organized.

After joint evaluation, we decided to maintain admission to the Internal Medicine ward for management of COVID-19 and conservative treatment of the plastron appendicitis.⁵ The patient received treatment with hydroxychloroquine and azithromycin for 5 days (in accordance with the COVID-19 hospital protocol at that time) and ceftriaxone for 2 days. After establishing the diagnosis of complicated acute appendicitis, broad-spectrum antibiotic treatment (piperacillin-tazobactam) was prescribed, ultimately completing a 10-day cycle.

From a respiratory standpoint, the patient required oxygen therapy with nasal cannulae at 2 L for the first 2 days; this was later able to be withdrawn, maintaining baseline oxygen saturations >95%. From the abdominal point of view, she

^{*} Please cite this article as: Vicario Bravo M, Chavarrías Torija N, Rubio-Pérez I. Sintomatología digestiva y COVID-19: importancia de descartar patología quirúrgica asociada. Cir Esp. 2021;99:385–387.



Fig. 1 – Coronal thoracoabdominal CT scans showing: (A) bilateral pulmonary involvement compatible with COVID-19; and (B) appendiceal inflammatory mass in the right iliac fossa.



Fig. 2 – Axial CT scan showing: (A) the appendiceal mass at diagnosis; and (B) practical resolution 9 days after the previous CT scan.

progressed favorably with less pain and was able to initiate oral intake progressively.

Prior to discharge, a follow-up CT scan (9 days after the initial one) showed persistence of bilateral mild/moderate pulmonary involvement (slight improvement) and a clear improvement in radiological findings compatible with complicated acute appendicitis, observing a significant decrease in the size of the collection $(1.4 \times 2 \text{ cm})$ (Fig. 2).

Likewise, follow-up lab studies showed an improvement in acute-phase reactants, although the following parameters remained high: D-dimer (870 ng/mL), gamma-glutamyl transferase (256 IU/L), lactate dehydrogenase (215 IU/L) and PCR (17.2 mg/L).

Given the good clinical–analytical–radiological evolution, the patient was discharged after 11 days of hospitalization with instructions to isolate at home, oral antibiotic therapy with amoxicillin-clavulanic acid for 5 more days, and a telephone follow-up after one month, with a good, sustained evolution. After this episode, a follow-up colonoscopy will be performed, and elective appendectomy will be considered. It is important to highlight the association found between patients who begin with complicated acute appendicitis/inflammatory mass and a possible tumor.⁶

In conclusion, the great variability of symptoms of COVID-19 disease makes it necessary to rule out underlying pathology, especially in patients with predominant abdominal symptoms. Since the availability of operating rooms has been reduced and the surgical treatment of a SARS-CoV-2-positive patient may pose an added risk,⁷ conservative treatment has been considered in those pathologies where it is safe and feasible.^{8,9} Nevertheless, patients should always be treated individually and closely monitored to detect unfavorable progression.

REFERENCES

 Rubio-Pérez I, Badia JM, Mora-Rillo M, Martín Quirós A, García Rodríguez J, Balibrea JM, con la colaboración del grupo Cirugía-AEC-COVID-19; Grupo colaborativo AEC-COVID-19 COVID-19. Key concepts for the surgeon. Cir Esp. 2020;30117– 22. <u>http://dx.doi.org/10.1016/j.ciresp.2020.04.009</u>.

Peeri NC, Shrestha N, Rahman MS, Zaki R, Tan Z, Bibi S, et al. The SARS MERS and novel coronavirus (COVID-19) epidemics, the newest and biggest global health threats: what lessons have we learned? Int J Epidemiol. 2020. <u>http://dx.doi.org/</u> <u>10.1093/ije/dyaa033</u>.

- Can-Valderrama O, Morales X, Ferrigni JC, Martín-Antona E, Turrado V, García A, et al. Reduction in emergency surgery activity during COVID-19 pandemic in three Spanish hospitals. Br J Surg. 2020. http://dx.doi.org/10.1002/bjs.11667.
- Gornet JM, Tran Minh ML, Leleu F, Hassid D. What do surgeons need to know about the digestive disorders and paraclinical abnormalities induced by COVID-19? J Visc Surg. 2020. <u>http://dx.doi.org/10.1016/j.jviscsurg.2020.04.017</u>.
- 5. Di Saverio S, Sibilio A, Giorgini E, Biscardi A, Villani S, Coccolini F, et al. The NOTA Study (Non Operative Treatment for Acute Appendicitis): prospective study on the efficacy and safety of antibiotics (amoxicillin and clavulanic acid) for treating patients with right lower quadrant abdominal pain and long-term follow-up of conservatively treated suspected appendicitis. Ann Surg. 2014;260:109–17.
- Liezten E, Grönroos JM, Mecklin J-P, Leppäniemi A, Nordtröm P, Rautio T, et al. Appendiceal neoplasm risk associated with complicated acute appendicitis – a population based study. Int J Colorectal Dis. 2019;34:39–46. <u>http://dx.doi.org/10.1007/</u> <u>s00384-018-3156-x</u>.
- Lei S, Jiang F, Su W, Chen C, Chen J, Mei W, et al. Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19 infection. E Clin Med. 2020;9:41.
- 8. Di Saverio S, Pata F, Gallo G, Carrano F, Scorza A, Sileri P, et al. Coronavirus pandemic and colorectal surgery: practical

advice based on the Italian experience. 2020. <u>http://</u><u>dx.doi.org/10.1111/codi.15056</u>.

 Balibrea JM, Badia JM, Rubio Pérez I, Martín Antona E, Álvarez Peña E, García Botella S, et al. Surgical management of patients with COVID-19 infection. Recommendations of the Spanish Association of Surgeons. Cir Esp. 2020. <u>http://</u> <u>dx.doi.org/10.1016/j.ciresp.2020.03.001</u>. S0009-739X.

Marina Vicario Bravo^{*}, Nuria Chavarrías Torija, Inés Rubio-Pérez

Servicio de Cirugía General, Hospital Universitario La Paz, Madrid, Spain

*Corresponding author. E-mail address: marinavicariobravo@gmail.com (M. Vicario Bravo).

2173-5077/

 \odot 2020 AEC. Published by Elsevier España, S.L.U. All rights reserved.

Subacute Spinal Injury: The Importance of Previous Trauma *



Lesión espinal subaguda: la importancia del antecedente traumático

Undetected spinal injuries after trauma can cause severe neurologic sequelae. After an initial trauma that is considered not serious, the appearance of neurological symptoms in the following days should alert us to the presence of a possible spinal injury. A careful medical history that includes inquiry about any possible history of trauma, together with a proper physical examination, should guide us toward rare entities that could potentially be very serious without appropriate treatment, such as this case we present.

The patient is a 40-year-old man with a history of behavioral disturbances and substance abuse, who came to the emergency department reporting progressive symptoms of hand weakness and gait disturbance over the previous 3–4 days. He had suffered moderate-energy left posterolateral blunt trauma associated with forced cervical rotation during an assault two weeks earlier, for which he did not visit the emergency department (thus, the initial injuries were not evaluated by a healthcare professional at that time). After an asymptomatic period of time, he developed progressive symptoms of predominantly distal weakness in both upper limbs. He was currently unable to open his hands and had a sensation of electrical shocks and cramping, especially in the left upper limb ipsilateral to the trauma. He also described a feeling of oppression in the form of a belt in the abdominal region and numbness in both upper extremities. The symptoms progressed and, in the 24 h prior to consultation, he began having difficulty walking and experienced a loss of strength in the lower limbs. He also reported left posterolateral cervical pain radiating to the ipsilateral upper limb.

Physical examination revealed a fever of 38 $^{\circ}$ C, with no swelling or hematoma in the area of the trauma, tetraparesis grade 4/5 (strength against resistance, without full strength), predominantly distal in the upper limbs (strength grade 2/5 in finger extension, not being able to extend the fingers against gravity) and proximal in the lower limbs, with bilateral extensor plantar response (injury to the bilateral pyramidal

^{*} Please cite this article as: Ginestal-López RC, Gómez-Iglesias P, García-Yepes M, Yus-Fuertes M, Fernández García C. Lesión espinal subaguda: la importancia del antecedente traumático. Cir Esp. 2021;99:387–389.