



## Video of the month

## Laparoscopic left hepatectomy with bile duct exploration for cholecystocholedocholithiasis and hepatolithiasis<sup>☆</sup>



### Hepatectomía izquierda con exploración de la vía biliar laparoscópica por colecistocolocolitiasis y hepatolitiasis

César Muñoz,<sup>a,b,\*</sup> Matías Orellana<sup>a</sup>

<sup>a</sup> Cirugía Digestiva, Hospital Regional de Talca, Talca, Maule, Chile

<sup>b</sup> Departamento de Cirugía, Universidad Católica del Maule, Talca, Maule, Chile




Fig. 1

Treatment of hepatolithiasis may require hepatectomy in cases of ductal stenosis and parenchymal atrophy (Fig. 1)<sup>1,2</sup>.

We present the case of a 72-year-old patient who consulted for jaundice. Ultrasound and magnetic resonance imaging of the abdomen with contrast demonstrated cholecystocholedocholithiasis associated with hepatolithiasis of the left lobe of the liver,

<sup>☆</sup> Please cite this article as: Muñoz C, Orellana M. Hepatectomía izquierda con exploración de la vía biliar laparoscópica por colecistocolocolitiasis y hepatolitiasis. Cir Esp. 2021. <https://doi.org/10.1016/j.ciresp.2020.11.022>

\* Corresponding author.

E-mail address: [cesarmunozcastro@gmail.com](mailto:cesarmunozcastro@gmail.com) (C. Muñoz).

stenosis of the hepatic duct and impacted gallstone. We performed cholecystectomy with exploration of the bile duct and primary closure to later continue with a laparoscopic left hepatectomy (Appendix B, Video).

The postoperative period was uneventful, and the patient was discharged on the fourth postoperative day.

---

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.cireng.2021.07.005>.

## REFERENCES

---

1. Jarufe N, Figueroa E, Muñoz C, Moisan F, Varas J, Valbuena JR, et al. Anatomic hepatectomy as a definitive treatment for hepatolithiasis: a cohort study. *HPB (Oxford)*. 2012;14:604–10.
2. Herman P, Perini MV, Pugliese V, Pereira JC, Machado MAC, Saad WA, et al. Does bilioenteric anastomosis impair results of liver resection in primary intrahepatic lithiasis? *World J Gastroenterol*. 2010;16:3423–6.