

Prognostic factors and analysis of mortality due to brain haemorrhages associated with vitamin K antagonist oral anticoagulants. Results from the TAC Registry*



Factores pronósticos y análisis de la mortalidad de las hemorragias cerebrales asociadas a anticoagulantes orales antagonistas de la vitamina K. Resultados del estudio TAC Registry

Dear Editor:

The article by Zapata-Wainberg et al.¹ called my attention as it makes no reference as to whether surgical treatment was used in any case. Our understanding of this condition may have improved if the benefits of surgery had been analysed. A contemporary article by Fernández-Sanz et al.² explicitly mentions that surgically treated patients were excluded.

Since the STICH study, we have questioned the usefulness of surgery for spontaneous haematomas and especially for haematomas associated with coagulation disorders. According to the clinical practice guidelines in intracerebral haemorrhage published by the Spanish Society of Neurology,³ the benefits of surgery do not outweigh the procedure's potential for harm in most cases; the guidelines provide several recommendations for surgical treatment. However, in daily clinical practice, we are often consulted about the possibility of surgical treatment that, in our opinion as experts, will be futile, even though it is difficult at times to contain the enthusiasm for treatment. In the meantime, we must acknowledge that retrospective studies such as the article in question do provide information, which is generally already known. However, there is a need for further prospective studies to establish, for example, the cases in which surgery would improve outcomes, in terms of both mortality and morbidity. Therefore, it is essential that future prospective

* Please cite this article as: Vilalta Castan J. Factores pronósticos y análisis de la mortalidad de las hemorragias cerebrales asociadas a anticoagulantes orales antagonistas de la vitamina K. Resultados del estudio TAC Registry. Neurología. 2020;35:505.

Urinary symptoms in patients with amyotrophic lateral sclerosis*



Síntomas urinarios en pacientes con esclerosis lateral amiotrófica

Dear Editor:

It was with great interest that we read the article by Martínez et al.¹ on the frequency of non-motor symptoms

* Please cite this article as: Vázquez-Costa JF and Arlandis S. Síntomas urinarios en pacientes con esclerosis lateral amiotrófica. Neurología. 2020;35:505–506.

studies include the participation of the specialties involved in this treatment: neurology, neurosurgery, neuroradiology, anaesthesiology, intensive care, etc. Despite the existence of many cerebrovascular disease registries, where the main concern of the authors often seems to be to find an attractive name, it may be more productive to focus on improving clinical history and using the minimum basic dataset (MBDS) to expand our knowledge of haemorrhagic strokes. The study by Hernández-Medrano et al.⁴ reports that the quality of the MBDS for cerebrovascular diseases guarantees the collection of valid information and that the registry of hospital discharges may be a useful tool for performing studies on this condition. The most useful parameters for determining whether a healthcare system is efficient are the destination at discharge and hospitalisation duration.

References

1. Zapata-Wainberg G, Quintas S, Ximénez-Carrillo Rico A, Benavente Fernández L, Masjuan Vallejo J, Gállego Culleré J, et al. Prognostic factors and analysis of mortality due to brain haemorrhages associated with vitamin K antagonist oral anticoagulants. Results from the TAC Registry. Neurología. 2018;33:419–26.
2. Fernández-Sanz A, Aladrén-Sangrós JA, Tejada-Meza H, Cruz-Velásquez GJ, Ángel-Ríos LF, Seral-Moral P, et al. Signos predictores de crecimiento precoz de la hemorragia intracerebral en la tomografía computarizada sin contraste y mortalidad. Rev Neurol. 2018;67:242–8.
3. Rodríguez-Yáñez M, Castellanos M, Freijo MM, López Fernández JC, Martí-Fàbregas J, Nombela F, et al. Guías de actuación clínica en la hemorragia intracerebral. Neurología. 2013;28:236–49.
4. Hernández Medrano I, Guillán M, Masjuan J, Alonso Cánovas A, Gogorcena MA. Fidabilidad del conjunto mínimo básico de datos en el diagnóstico de la enfermedad cerebrovascular. Neurología. 2017;32:74–80.

J. Vilalta Castan

Servicio de Neurocirugía, Hospital Vall d'Hebron, Barcelona, Spain

E-mail address: pelagatosiv@gmail.com

<https://doi.org/10.1016/j.nrleng.2018.12.013>

2173-5808/

© 2019 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. All rights reserved.

in patients with amyotrophic lateral sclerosis (ALS). The relatively high incidence of these symptoms is becoming increasingly apparent, and has considerable implications for prognosis and treatment.² However, the authors report a surprisingly low incidence of "urinary problems" (2%). It is unclear what type of urinary problems the authors are referring to, but a previous study by our research group³ found lower urinary tract symptoms (evaluated with standardised questionnaires) in 43.6% of a series of patients with ALS, with 26.3% of patients presenting urgency urinary incontinence; these incidence rates are similar to those reported in other studies.^{4,5} In most patients, symptoms affected both bladder filling and voiding, and were