



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Scientific and Technological Research

Are we controlling postoperative pain?☆

Jorge Enrique Machado-Alba^{a,*}, Manuel Enrique Machado-Duque^b,
Viviana Calderón Flórez^b, Alexandra Gonzalez Montoya^b, Felipe Cardona Escobar^b,
Richard Ruiz García^b, Julian Montoya Cataño^b

^a Physician, Master in Pharmacoepidemiology, Master in Pharmacology, Faculty, Health Sciences School, Pharmacoepidemiology and Pharmacovigilance Research Group, Universidad Tecnológica de Pereira – Audifarma S.A., Pereira, Colombia

^b Medical Student, Health Sciences School, Universidad Tecnológica de Pereira, Pereira, Colombia

ARTICLE INFO

Article history:

Received 19 June 2012

Accepted 23 January 2013

Available online 20 March 2013

Keywords:

Pain, Postoperative

Analgesics, Opioid

Pain

Analgesia

ABSTRACT

Introduction: Immediate postoperative pain has been underestimated and managed inadequately.

Objectives: To assess perceived pain 4 h after surgery in patients at the San Jorge University Hospital in the city of Pereira.

Materials and methods: Cross-sectional study in patients over 18 years of age was conducted between September 2nd and October 28th, 2011. Postoperative pain intensity was assessed using the Visual Analog Scale, 4 h after completing the procedure. Social, demographic, clinical and pharmacological variables were considered. The analysis was done using the SPSS 20.0 for Windows.

Results: Of the 213 postoperative patients studied, 114 (53.6%) were women and 99 (46.4%) were men, with a mean age of 47.1 ± 20.0 years. At 4 h, 51.4% of patients did not have pain control. There was a statistically significant association between lack of control and age, living in the urban area, type of surgery, non-adherence to the dose, and monotherapy analgesia.

Discussion: Inadequate pain control requires revisiting its management, ideally on the basis of clinical practice guidelines and using analgesic drugs at adequate doses and intervals.

© 2012 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L. All rights reserved.

¿Estamos controlando el dolor posquirúrgico?

RESUMEN

Introducción: El dolor en el posquirúrgico inmediato ha sido subvalorado y manejado inadecuadamente.

Objetivos: Evaluar la percepción del dolor a las 4 h del postoperatorio de pacientes del Hospital Universitario San Jorge de Pereira (Colombia).

Materiales y métodos: Estudio de corte transversal en pacientes mayores de 18 años entre el 2 de septiembre y el 28 de octubre de 2011. Se valoró la intensidad del dolor postoperatorio

Palabras clave:

Dolor posoperatorio

Analgésicos opioides

Dolor

Analgesia

☆ Please cite this article as: Machado Alba JE, et al. ¿Estamos controlando el dolor postquirúrgico? Rev Colomb Anestesiolog. 2013;41:132-8.

* Corresponding author at: Calle 105 No. 14-140, Pereira, Risaralda, Colombia.

E-mail address: machado@utp.edu.co (J.E. Machado-Alba).

mediante escala visual analógica a las 4 h del procedimiento. Se consideraron variables sociodemográficas, clínicas y farmacológicas. El análisis se hizo con SPSS 20.0 para Windows.

Resultados: Se evaluaron 213 pacientes en postoperatorio, 114 (53,6%) mujeres y 99 (46,4%) hombres, con edad promedio de $47,1 \pm 20,0$ años. El 51,4% de los pacientes no tenía controlado el dolor a las 4 h. Las variables edad, residencia urbana, tipo de cirugía, incumplimiento de la dosis y monoterapia analgésica se asociaron de manera estadísticamente significativa con la falta de control.

Discusión: El inadecuado control del dolor obliga a replantear su manejo idealmente con guías de práctica clínica y con el empleo de medicamentos analgésicos a las dosis e intervalos adecuados.

© 2012 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

According to the International Association for the Study of Pain (I.A.S.P.), pain is an unpleasant sensory and emotional experience associated with existing or potential tissue injury.¹ Dramatic progress has been made in controlling postoperative pain and there are now multiple experts and publications in this field.² Despite significant interest in improving postoperative pain management, evidence shows that world prevalence of moderately intense pain in hospitalized patients ranges between 26.0%, and 33.0%, whereas prevalence of severe pain has been estimated to be between 8.0% and 13.0%.³ Postoperative complications caused by pain in the main organ systems have been well-described. Tissue injury triggers a series of responses that may cause ventilation abnormalities (5.0–25.0% of patients), local circulation disorders, gastrointestinal and urinary disorders, and even lead to infarction or heart failure, not to mention other abnormalities in carbohydrate, lipid and protein metabolism, as well as diencephalic and cortical responses, anxiety, fear and depression, that occur when pain is not well managed.^{4–6}

It is now known that adequate control of acute postoperative pain is one of the cornerstones in achieving fast postoperative recovery. Administratively, this results in shorter hospital stays and lower costs, and from the medical standpoint, it implies reduced morbidity and mortality.^{7,8} Ever since the American Pain Society declared pain to be the “fifth vital sign”, several initiatives have been undertaken to improve its control, including the implementation of a numerical scoring scale called the Visual Analog Scale (VAS) consisting of 10 integer numbers for the subjective measurement of pain intensity.^{9,10} A pain score of 4 or more requires a comprehensive pain assessment and rapid intervention by the healthcare provider.^{11,12}

Although no drug regimen has been able to completely eliminate postoperative morbidity and mortality, adequate pain management leads to early ambulation, which, together with vomiting and ileus control, oral feeding, and preoperative antibiotic therapy, is the mainstay for comprehensive postoperative management.¹³ The pharmacological armamentarium for pain management available at the present time is quite broad and includes several groups such as opioids, analgesics, non-steroidal anti-inflammatory agents (NSAIDs), and local anesthetics. It is recommended to provide two analgesics with

a different mechanism of action in order to achieve more effective analgesia and reduce adverse reactions with the use of a lower dose of each drug.¹⁴

The goal of this study was to determine perceived pain intensity in the postoperative period by means of pain assessments at 4 h using the VAS, and to determine the social, demographic, clinical and pharmacological variables associated with pain control or lack of control in patients taken to surgery at the San Jorge University Hospital in Pereira (HUSJ), in order to optimize management.

Materials and methods

A cross-sectional study was conducted at HUSJ in a population of patients over 18 years of age undergoing surgery between 7:00 am and 6:00 pm, from September 2nd to October 28th, 2011. Assessment of postoperative pain intensity was done using the VAS in millimeters (mm), in which five categories were established. Values of 0 and 100 are absolute and represent independent categories, and the following reference values were used: 0 mm absence of pain, 1–19 mm very mild pain, 20–39 mm mild pain, 40–59 mm intermediate pain, 60–79 mm severe pain, 80–99 mm very severe pain, and 100 mm the worst possible pain; scores over 40 mm were used for undefined pain. Consequently, pain was considered to be under control when scores were lower than or equal to 39 mm.^{10,15–17}

Assessment was done 4 h after completion of the procedure with a view for assessing immediate postoperative pain management. Patients who could not take the test because of neurologic deficits, disabling motor disorders, mental retardation and severe mental diseases were excluded. The information was obtained through patient interviews by duly trained final-year medical students of Universidad Tecnológica in Pereira. Access to patient clinical records and surgical notes was also obtained by means of an informed consent. The data collection tool was developed by the researchers and included the following variables, besides the VAS:

Social, demographic and toxicological variables: Age, gender, health insurance regime (subsidized or contributive), socio-economic bracket (low, medium, high), education (primary, secondary, higher), place of residence (urban or rural), cigarette smoking, use of psychoactive substances, NSAIDs, steroids and anti-depressants.

Clinical variables: Type of surgical procedure (general, brain, urologic, plastic, otolaryngological, gynecological surgery, etc.), intra- and post-operative complications, type of anesthesia (general inhaled, intravenous, conductive, local, etc.), estimated surgical risk (high, moderate and low). For this latter variable, low risk included minimally invasive surgery with blood losses under a 200 cm³; moderate risk included moderately invasive procedures with fluid exchange and potential blood losses of up to 1000 cm³, and/or moderate mortality/morbidity; and high risk included highly invasive procedures such as radical or extensive upper abdominal, thoracic or brain surgeries with potential blood losses greater than 1000 cm³ and significant associated mortality/morbidity.

Pharmacological variables: Analgesics prescribed immediately after surgery and up to 4 h, were grouped according to pharmacological class and their use either as monotherapy or in combination, dose, dosing interval for each, drug-related adverse reactions, and use of analgesic premedication. Morphine, meperidine and fentanyl were used as strong opioids and tramadol was used as a weak opioid.

The protocol was approved by the Ethics Committee of the Health Sciences School at Universidad Tecnológica in Pereira, under the category of "Low-to-minimum risk research", in accordance with Resolution No. 008430 of 1993 of the Colombian Health Ministry, which sets forth the scientific, technical and administrative standards for health research. The analysis was conducted using the SPSS software, version 2.0 for Windows (IBM, USA). The Student t test or ANOVA were used for comparing quantitative variables, and the χ^2 test was used for comparing categorical variables. Logistic regression models were applied using pain control as the dependent variable, and those that were significant in the bivariate analysis were used as independent variables. A statistical significance level of $p < 0.05$ was established.

Results

Of a total of 213 postoperative patients who were assessed, 114 (53.6%) were women and 99 (46.4%) were men, with a mean age of 47.1 ± 20.0 years (range: 18–86 years). Table 1 summarizes the social, demographic and clinical characteristics of the patients included in the study. Pain measurements using the VAS were assessed in 213, with a mean pain score of 40.0 mm, where 111 (51.4%) patients had no pain control (VAS ≥ 40 mm) and 102 (47.2%) had pain control; moreover, it was found that 25 patients (11.7%) required analgesia during their hospital stay as a result of pain intensity. In addition, there were nine patients with no analgesic prescription, including one case of exploratory laparotomy and one C-section.

Fig. 1 shows patient distribution by pain range found on assessment, and Table 2 groups analgesics, dose and number of medications received by each patient and their associations, arranged by frequency of use for pain management, where dipirone was the most frequently used analgesic in monotherapy and in combination, followed by morphine and fentanyl.

Table 1 – Social, demographic, medical and surgical characteristics of 213 surgical patients, 2011.

Social and demographic characteristics	n = 213	(%)
Gender (males/females)	99/114	46.4/53.6
Age (mean ± SD, years)	47.1 ± 20.0	
Marital status:	44/168	20.4/77.8
Single/partner		
Health Insurance Regime:	13/197	6.0/91.2/2.8
Contributive/ Subsidized/NA		
Education: Primary/ Secondary/Higher	129/75/9	59.7/34.7/4.1
Residence: Urban/Rural	156/57	72.2/26.4
Socio-economic bracket:	127/66/13/2/1	58.8/30.6/6.0/0.9/0.5
1/2/3/4/5		
Personal history (consumption)		
Cigarette smoking	66	30.6
Psychoactive substances	13	6
Alcohol	47	21.8
Pharmacological history		
NSAIDS	46	21.3
Anti-depressants	4	1.9
Glucocorticoids	7	3.2
Type of surgery		
Plastic	10	4.7
Gynecological	27	12.7
Orthopedic	82	38.5
General	61	28.6
Urology	17	8.0
Laparoscopic	3	1.4
Neurosurgery	6	2.8
Peripheral vascular	4	1.9
Otolaryngological	3	1.4
Anesthetic premedication		
Yes/No	8/205	3.8/96.2
Type of anesthesia		
General intravenous	21	9.7
Conductive	83	39.9
General inhaled	91	42.4
Regional	7	3.2
Local	10	4.4
N/A	1	0.40

Comparison of patients with and without pain control

Table 3 shows the results of the bivariate analysis that enables a comparison of subgroups of patients with pain control and those with no pain control. It was found that marital status, health insurance regime, education, cigarette smoking, use of alcohol, psychoactive substances, NSAIDS or anti-depressants, surgical risk, use of anesthetic premedication, onset of GI bleeding, nausea and concurrent comorbidities such as diabetes mellitus, ischemic heart disease, depression, epilepsy, COPD, and renal failure were variables with no statistically significant association with the lack of pain control. It was found that gender, vomiting, age between 18 and 44, urban place of residence and the combined use of a strong opioid plus an antipyretic analgesic were variables with a statistically significant association with lack of pain control; and age between 45 and 64 years, urologic or peripheral vascular surgeries, and adherence to the dosing instructions were




Fig. 1 – Percentage of patients according to pain intensity at 4 postoperative in 213 patients of the Hospital Universitario San Jorge, Pereira, 2011.

variables with a statistically significant association with pain control.

Multivariate analysis

In the multivariate analysis, the dependent variable was the lack of pain control and the independent variables were those that showed some significant association in the bivariate analysis. It was found that having an age different from the range between 44 and 64 years (OR: 0.4; 95% CI: 0.148–0.973, $p = 0.044$), living in an urban area (OR: 2.2; 95% CI: 1.135–4.283, $p = 0.02$),

Table 2 – Drugs and regimens most commonly used in postoperative patients, 2011.

Pharmacological variables	4 h	
<i>Number of drugs per patient</i>	<i>n = 213</i>	<i>%</i>
None	9	4.2
1	90	42.2
2	75	35.2
3	29	13.6
4	10	4.6
<i>Analgesics used</i>	<i>n = 352</i>	
Dipirone	128	36.4
Morphine	80	22.7
Fentanyl	79	22.4
Diclofenac	38	10.8
Remifentanyl	7	2.0
Tramadol	4	1.1
Lidocaine	3	0.9
Acetaminophen	1	0.3
<i>Most frequent regimens</i>		
Antipyretic analgesic alone	45	21.1
Strong opioid + antipyretic analgesic	63	29.6
Strong opioid	42	19.7
NSAID + antipyretic analgesic	10	4.7
Tramadol + antipyretic analgesic	2	0.9
Strong opioid + antipyretic analgesic + NSAID	12	5.6
Others	30	14.1

and having undergone a procedure other than urologic surgery (OR: 0.05; 95% CI: 0.006–0.4, $p = 0.005$) were the independent variables with a statistically significant association with the lack of pain control.

Discussion

The importance of adequate management of postoperative pain and the need to set up specialized multi-disciplinary centers with anesthetists, general practitioners and nurses trained in the use of analgesic drugs, and the provision of rigorous pain monitoring and control have been demonstrated in the world literature.^{14,18} This study found a high prevalence of uncontrolled postoperative pain at 4 h, a result that was very different from that found by a meta-analysis of more than 20,000 patients around the world of 11% of cases with postoperative pain, but very similar to Spanish and Colombian publications that report low pain control ranging between 40.0% and 69.3% of postoperative patients assessed.^{3,19–24} The wide use of drugs as monotherapy, contrary to the ASA guidelines that propose the use of 2 medications with a different mechanism of action at appropriate doses and dosing schedules, may account for the low levels of pain control.^{14,25,26}

The use of dipirone, alone or in combination with opioids, and its association with the lack of pain control is consistent with the results of similar studies conducted in Colombia in which monotherapy was the most widely used with inadequate pain control.^{19,23} Results of Spanish studies show that the most frequent prescriptions are opioids alone or in association with NSAIDs.^{20,27} Almost all the patients who required rescue analgesia continued to experience pain, which is evidence of the failure to use more effective analgesia or analgesics at higher doses in patients with very intense pain perception. Some of the international studies have also shown the ineffectiveness of rescue analgesia.^{19,20}

Opioid administration under a suboptimal regimen may be due to the lack of knowledge of the pharmacokinetics of these drugs, and to the fear of adverse reactions. This practice already reported by other authors does not achieve pain relief and maintains the risk of adverse reactions, including delirium in the elderly.^{20,25,28} It is worth noting that the finding of patients with no analgesic prescription is inconsistent with similar studies in which all the patients received analgesic management. This may be an indication of the indifference of healthcare personnel regarding pain, and lack of knowledge of patient rights.^{19–22,25,26,29,30}

Differences in terms of pain perception and control have already been reported in other studies, and it has been found that males in Spain report more intense pain.^{23,29} Living in an urban area was associated with lack of pain control when compared with the rural area, something that is not reported in the literature in association with postoperative pain. This relationship that may have cultural components should be explored in more depth.

The specialties performing surgery most frequently at HUSJ were orthopedics, general surgery, obstetrics and gynecology, and it was in those procedures where uncontrolled pain was also more prevalent. Considering that there was

Table 3 – Bivariate analysis of pain control at 4 h versus the main social, demographic, pharmacological and clinical variables of surgical patients, 2011.

Characteristics	Pain control at 4 h		No pain control at 4 h		p* value	RR	95%** CI Lower-upper
	Number	%	Number	%			
<i>Gender</i>							
Male	57	57.6	42	42.4	0.005	0.460	0.267-0.795
Female	45	39.5	69	60.5			
<i>Ages</i>							
Young adult (18-44)	36	36.4	63	63.6	0.03	0.442	0.255-0.764
Middle age (45-64)	39	62.9	23	37.1	0.03	2.449	1.335-4.495
Elderly (65+)	25	48.1	27	51.9	0.769	1.099	0.586-2.060
<i>Required rescue analgesia</i>							
Yes	2	8.0	23	92.0	<0.001	0.079	0.018-0.345
No	100	53.2	88	46.8			
<i>Vomiting</i>							
Yes	8	27.6	21	72.4	0.023	0.377	0.159-0.894
No	94	51.1	90	48.9			
<i>Place of residence</i>							
Urban	68	43.6	88	56.4	0.028	0.505	0.273-0.935
Rural	34	59.6	23	40.4			
<i>Surgical specialty</i>							
Urologic	16	94.1	1	5.9	<0.001	21.0	2.735-161.627
Peripheral vascular	4	100	0	0	0.033	0.462	0.400-0.534
General	25	40.3	37	59.7	0.197	0.676	0.372-1.228
Orthopedics	32	39.0	50	61.0	0.059	0.585	0.335-1.023
Obstetrics and gynecology	10	34.5	19	65.5	0.14	0.543	0.240-1.231
<i>Type of anesthesia</i>							
Conductive	37	44.0	47	56.0	0.456	0.811	0.468-1.406
General intravenous	10	45.5	12	54.5	0.861	0.924	0.381-2.239
General inhaled	47	51.1	45	48.9	0.327	1.310	0.763-2.251
<i>Adherence to analgesic dose</i>							
Yes	59	48.8	62	51.2	0.609	1.151	0.671-1.972
No	43	45.3	52	54.7			
<i>Adherence to dosing schedule</i>							
Yes	11	73.3	4	26.7	0.036	3.324	1.024-10.792
No	88	44.5	110	55.5			
<i>Analgesic regimens</i>							
Antipyretic analgesic alone	24	53.3	21	46.7	0.356	1.363	0.705-2.632
NSAID + antipyretic analgesic	6	60.0	4	40.0	0.407	1.719	0.471-6.272
Strong opioid + antipyretic analgesic	23	36.5	40	63.5	0.043	0.539	0.295-0.984
Strong opioid + antipyretic analgesic + NSAID	5	41.7	7	58.3	0.692	0.788	0.242-2.564
Strong opioid	22	52.4	20	47.6	0.456	1.293	0.658-2.538

* Based on Chi square test.

** 95% confidence Interval, lower-upper limit.

greater evidence of uncontrolled pain in orthopedic, general surgery and obstetric and gynecological procedures, they are shown to involve determining factors in pain perception such as extensive tissue damage and the involvement of several systems. It has already been reported that the type of intervention, the surgical technique and the anesthetic management are major determining factors of pain intensity and duration.^{19,31} In the analysis by type of surgery, it was found that urologic and peripheral vascular surgeries were associated with improved pain control, contrary to the results found by another Colombian study in

which those procedures were associated with lack of pain control.¹⁹

The main limitations found in this study are due to the fact that, added to the lack of entries in some clinical records, the VAS measures only one dimension as it examines the sensory component only, excluding the patient's affective and cognitive components.²⁰ The wide variety of regimens and drugs used for pain management imposes a limitation on the ability to interpret the results, and highlights the importance of incorporating effective and easy-to-use postoperative pain management guidelines.²⁶

We consider that the lack of postoperative pain control is evident at HUSJ. It is an issue associated with the female gender, the onset of vomiting, age between 18 and 44 years, living in the urban area of the municipality of Pereira or the department of Risaralda, and the combined use of a strong opioid plus an antipyretic analgesic, prescribed at inadequate intervals.

There is a need to revisit pain management in this hospital and make adjustments on the basis of international guidelines, or internally developed guidelines, with a view at ensuring adequate control of acute postoperative pain. This may be accomplished by focusing on the importance of postoperative pain control, and making appropriate use of analgesic medications, in doses and schedules tailored to the needs of individual patients. It would also be important to consider creating an acute pain unit, which has proven to be highly effective in comprehensive patient management and control.^{3,14,18,32} The use of clinical practice guidelines for adequate analgesic management has been shown to reduce pain and its complications significantly.²⁶

Funding

This research has received funds by the Universidad Tecnológica de Pereira.

Conflicts of interest

The authors declare not to have any conflicts of interest.

REFERENCES

- International Association for the Study of Pain [sede web]. Seattle: Merskey H, Bogduk N; 1994. IASP taxonomy [accessed on 16.02.12]. Available at: <http://www.iasp-pain.org/Content/NavigationMenu/GeneralResourceLinks/PainDefinitions/default.htm>
- Carr DB, Goudas LC. Acute pain. *Lancet*. 1999;353:2051-8.
- Dolin SJ, Cashman JN, Bland JM. Effectiveness of acute postoperative pain management. I. Evidence from published data. *Br J Anaesth*. 2002;89:409-23.
- Muñoz-Blanco F, Salmerón J, Santiago J, Marcote C. Complicaciones del dolor postoperatorio. *Rev Soc Esp Dolor*. 2001;8:194-211.
- Martinez-Vazquez de Castro J, Torres LM. Prevalencia del dolor postoperatorio. Alteraciones Fisiopatológicas y sus repercusiones. *Rev Soc Esp Dolor*. 2000;7:465-76.
- Finkel DM, Schlegel HR. El dolor postoperatorio. Conceptos básicos y fundamentos para un tratamiento adecuado. *Rev Hosp Gen Agudos Dr J M Ramos Mejía*. 2003;8:1-17.
- Reyes Fierro A, de la Gala García F. Dolor postoperatorio: analgesia multimodal. *Patología del Aparato Locomotor*. 2004;2:176-88.
- Brandsborg B. Pain following hysterectomy: epidemiological and clinical aspects. *Dan Med J*. 2012;59:B4374.
- Campbell J. Pain: the fifth vital sign. In: Presidential address presented at the meeting of the American Pain Society. 1995.
- Gkotsi A, Petsas D, Sakalis V, Fotas A, Triantafyllidis A, Vouros I, et al. Pain point system scale (PPSS): a method for postoperative pain estimation in retrospective studies. *J Pain Res*. 2012;5:503-10.
- Craine M, Kerns RD. Pain management improvement strategies in the Veteran's Health Administration. *APS Bull*. 2003. Available at: <http://www.ampainsoc.org/pub/bulletin/sep03/article1.htm> [accessed on 16.02.12].
- Veterans Health Administration. Pain as the 5th vital sign toolkit. October 2000, revised edition. Geriatrics and Extended Care Strategic Healthcare Group, National Pain Management Coordinating Committee; 2005. Available at: <http://www.va.gov/oa/pocketcard/pain.asp> [accessed on 16.02.12].
- Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth*. 1997;78:606-17.
- American Society of Anesthesiologists Task Force on Acute Pain Management. Practice guidelines for acute pain management in the perioperative setting: an updated report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology*. 2004;100:1573-81.
- DeLoach LJ, Higgins MS, Caplan AB, Stiff JL. The visual analog scale in the immediate postoperative period: intrasubject variability and correlation with a numeric scale. *Anesth Analg*. 1998;86:102-6.
- Dexter F, Chestnut DH. Analysis of statistical tests to compare visual analog scale measurements among groups. *Anesthesiology*. 1995;82:896-902.
- Myles PS, Troedel S, Boquest M, Reeves M. The pain visual analog scale: is it linear or nonlinear? *Anesth Analg*. 1999;89:1517-20.
- Robaina F. ¿Por qué las Unidades del Dolor deben ser multidisciplinarias? *Rev Soc Esp Dolor*. 2005;12:137-40.
- Cardona E, Castaño M, Builes A, Castro G. Manejo del dolor postquirúrgico en el Hospital Universitario San Vicente de Paul de Medellín. *Rev Col Anest*. 2003;31:111-6.
- López S, Méndez H, Real J, Gordo F, Fernández DL. Manejo de la analgesia postoperatoria en las primeras 24 horas en un Hospital de segundo nivel. *Rev Soc Esp Dolor*. 2006;13:18-23.
- Gallego JI, Rodríguez de la Torre MR, Vazquez-Guerrero JC, Gil M. Estimation of the prevalence and severity of postoperative pain and relation with patient satisfaction. *Rev Soc Esp Dolor*. 2004;11:197-202.
- Cadauid A, Mendoza J, Gómez N, Berrío M. Prevalencia de dolor agudo posoperatorio y calidad de la recuperación en el Hospital Universitario San Vicente de Paúl, Medellín, Colombia, 2007. *IATREIA*. 2009;22:11-5.
- Machado-Alba J, Quintero A, Mena M, Castaño C, Lopez E, Marin D, et al. Evaluación del manejo del dolor postquirúrgico en pacientes adultos de una clínica de tercer nivel de Pereira, Colombia. *Rev Investigaciones Andina*. 2012;14:547-59.
- Allegrí M, Clark MR, De Andrés J, Jensen TS. Acute and chronic pain: where we are and where we have to go. *Minerva Anestesiol*. 2012;78:222-35.
- Rocchi A, Chung F, Forte L. Canadian survey of postsurgical pain and pain medication experiences. *Can J Anaesth*. 2002;49:1053-6.
- Usichenko TI, Röttenbacher I, Kohlmann T, Jülich A, Lange J, Mustea A, et al. Implementation of the quality management system improves postoperative pain treatment: a prospective pre-/post-interventional questionnaire study. *Br J Anaesth*. 2013;110:87-95.
- Moreno-Azcoitia M, De Andrés J, Torres L, Vidal M. Estudio Observacional sobre el dolor postoperatorio leve o moderado desde el punto de vista del anestesiólogo en España. *PATHOS. Rev Soc Esp Dolor*. 2007;14:550-67.
- Morrison RS, Magaziner J, Gilbert M, Koval KJ, McLaughlin MA, Orosz G, et al. Relationship between pain and opioid analgesics on the development of delirium following hip fracture. *J Gerontol A Biol Sci Med Sci*. 2003;58:76-81.

29. Valentín López B, García Caballero J, Muñoz Ramón JM, Aparicio Grande P, Díez Sebastián J, Criado Jiménez A. Postoperative pain management in a tertiary care hospital: initial situation prior to starting a quality assurance program. *Rev Esp Anestesiología Reanimación*. 2006;53:408-18.
30. Brennan F, Cousins MJ. Pain relief as a human right. *Rev Soc Esp Dolor*. 2005;12:17-23.
31. Soler E, Faus MT, Montaner MC, Morales F, Martínez-Pons V. Prevalencia, tratamiento y factores determinantes del dolor postoperatorio en un servicio de cirugía general y aparato digestivo. *Rev Soc Esp Dolor*. 2001;8:317-26.
32. Manion SC, Brennan TJ. Thoracic epidural analgesia and acute pain management. *Anesthesiology*. 2011;115:181-8.