



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Scientific and Technological Research

Temperature management during the perioperative period and frequency of inadvertent hypothermia in a general hospital[☆]

Caridad Greta Castillo Monzón^{a,*}, César Augusto Candia Arana^a,
Hugo Antonio Marroquín Valz^b, Fernando Aguilar Rodríguez^a,
Jairo Julián Benavides Mejía^c, José Antonio Alvarez Gómez^a

^a Especialista en Anestesiología y Reanimación, Servicio de Anestesiología y Reanimación, Complejo Hospitalario Universitario de Cartagena-Murcia, Spain

^b Médico de Urgencias, Servicio de Urgencias, Magister en Administración, Servicio de Urgencias, Complejo Hospitalario Universitario de Cartagena-Murcia, Spain

^c Médico Residente de tercer año de Anestesiología, Servicio de Anestesiología y Reanimación, Complejo Hospitalario Universitario de Cartagena-Murcia, Spain

ARTICLE INFO

Article history:

Received 20 November 2012

Accepted 14 February 2013

Available online 12 April 2013

Keywords:

Anesthesia

Hypothermia

Body temperature regulation

Perioperative period

ABSTRACT

Introduction: Perioperative hypothermia is associated with increased morbidity and mortality. Consequently, surgical patient temperature should be the fundamental concern but, nonetheless, it is still the least valued physiological parameter.

Objectives: To assess temperature management during the perioperative period and determine the frequency of inadvertent hypothermia and related factors.

Materials and methods: Prospective observational study in adult patients scheduled for surgical procedure with anesthesia time ≥ 30 min. Hypothermia is defined as a forehead skin temperature $\leq 35.9^\circ\text{C}$. The null hypothesis of no difference between patients with normothermia and hypothermia was proposed. Comparison of quantitative variables was analyzed with the Student "t" test, and the Chi square was used for the qualitative variables. The analysis was followed by a logistic regression analysis.

Results: We included 167 consecutive patients; intraoperative monitoring of temperature was used in 10% of patients, and the use of warm intravenous fluids and forced air heating in 78% and 63%, respectively. The frequency of inadvertent hypothermia was 56.29%, associated with age ≥ 65 years, female gender and BMI ≥ 30 kg/m². This last variable might have been influenced by the method of temperature measurement.

Conclusion: Warming measures without temperature monitoring do not result in the desired effect. The high frequency of inadvertent hypothermia requires action guidelines for prevention and management, especially in high-risk patients who, in this study, were patients ≥ 65 years of age and females.

© 2012 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L. All rights reserved.

[☆] Please cite this article as: Castillo Monzón CG, et al. Manejo de la temperatura en el perioperatorio y frecuencia de hipotermia inadvertida en un hospital general. Rev Colomb Anestesiolog. 2013;41:97-103.

* Corresponding author at: Calle Vicenza No. 29, Polígono Santa Ana, CP: 30310, Cartagena, Murcia, Spain.

E-mail address: gretacastillo9@hotmail.com (C.G. Castillo Monzón).

Manejo de la temperatura en el perioperatorio y frecuencia de hipotermia inadvertida en un hospital general

R E S U M E N

Palabras clave:

Anestesia
Hipotermia
Regulación de la temperatura corporal
Periodo perioperatorio

Introducción: La hipotermia perioperatoria está asociada con mayor morbimortalidad, por lo que la temperatura del paciente quirúrgico debería ser una preocupación fundamental; sin embargo, es el parámetro fisiológico menos valorado.

Objetivos: Evaluar el manejo de la temperatura en el perioperatorio, determinar la frecuencia de hipotermia inadvertida y los factores relacionados.

Material y métodos: Estudio prospectivo observacional en pacientes adultos programados para procedimiento quirúrgico con tiempo ≥ 30 min de anestesia. La hipotermia se definió como una temperatura de la piel de la frente $\leq 35,9^\circ\text{C}$. Se planteó la hipótesis nula de no diferencia entre los pacientes con normotermia e hipotermia. La comparación de las variables cuantitativas fue analizada con la prueba t de Student y las cualitativas con la prueba del Chi cuadrado, y después se realizó un análisis de regresión logística.

Resultados: Se incluyeron 167 pacientes consecutivos; la monitorización intraoperatoria de la temperatura se usó en el 10% de los pacientes, el uso de líquidos intravenosos tibios y calentamiento con aire forzado en el 78 y el 63%, respectivamente. La frecuencia de hipotermia inadvertida fue del 56,29%, asociada a edad ≥ 65 años, sexo femenino e índice de masa corporal $\geq 30\text{ kg/m}^2$. Esta última variable podría estar influenciada por el método de medición de la temperatura.

Conclusiones: Las medidas de calentamiento sin monitorización de la temperatura no tienen el efecto esperado. La frecuencia elevada de hipotermia inadvertida hace necesaria una guía de actuación de prevención y manejo en especial en pacientes de riesgo, que en este estudio fueron edad ≥ 65 años y sexo femenino.

© 2012 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

There is evidence that hypothermia is associated with systemic complications¹⁻⁶ and alters the pharmacokinetics and pharmacodynamics of anesthetic agents.⁷⁻¹¹ The most frequent perioperative thermoregulation alteration is inadvertent hypothermia.¹² The reported incidence varies greatly between 6% and 90%¹³⁻¹⁷ depending on the type of surgery, and it is associated with a high potential for complications,¹ including increased blood loss,^{2,3} morbid cardiac events,⁴ compromised healing and surgical wound infection,^{5,6} and higher mortality.¹⁸

Intraoperative temperature monitoring became popular starting in the 1960s, and even more than 50 years later this physiological parameter is still not monitored rigorously or managed by the anesthetist, despite the knowledge that adequate treatment improves the final outcome for the surgical patient.^{19,20}

Few recommendations are available regarding temperature. The 2007 guidelines of the American College of Cardiology on perioperative cardiovascular care and assessment for non-cardiac surgery recommend maintenance of perioperative normothermia on the basis of Class I (level B) evidence.²¹ The guidelines of the American Society of Anesthesiologists (ASA)²² mention temperature very briefly: "temperature must be assessed periodically during recovery from anesthesia". In England, the National Institute for Health and Clinical Excellence (NICE), published in 2008 some guidelines for the management of inadvertent perioperative

hypothermia, including recommendations for its adequate management throughout the whole preoperative, intraoperative and postoperative period.²³

The objective of the study was to assess temperature management during the perioperative period and determine the frequency of inadvertent hypothermia and associated factors.

Materials and methods

The protocol for this prospective observational study was approved by the Ethics Review Board of the University Hospital Complex in Cartagena-Murcia, Spain. Adult patients coming for any type of elective surgery under different anesthetic techniques lasting more than 30 min were included consecutively in the study. Obstetric and pediatric patients were excluded.

The following were the data collected in the study: sex, age in years, weight in kilograms, height in centimeters, body mass index (BMI), ASA classification, surgical specialty, anesthesia time and type, use of temperature monitoring, use of warm intravenous fluids and of forced-air warming systems during surgery, and clinical manifestations of hypothermia in the recovery room. Patients were divided into two age groups – under 65 years of age and 65 or more. BMI was classified as lower than 30 kg/m^2 and equal to or greater than 30 kg/m^2 . Forehead skin temperature was recorded as soon as patients were brought to the recovery room and 1 h later, considering that patient stay in the recovery room is usually longer than 1 h but shorter than 2 h. Using or not using intraoperative

temperature monitoring, and the techniques used to maintain temperature were left to the discretion of the anesthetist, and that information was gathered verbally from the anesthesia team upon arrival of the patient to the recovery room. During the study, it was decided not to inform the anesthetists about the follow-up conducted in the recovery room in order not to induce changes in monitoring behavior or in their strategies for managing temperature. Clinical manifestations of hypothermia during the stay in the recovery room were recorded.

Reusable or disposable sensors of the Ohmeda Aestiva 3000 anesthesia machine were used in those cases where intraoperative temperature was measured, with inferior or nasopharyngeal recording. The Bair Huggers 750 and 775 units were used in those cases where forced-air warming (blanket/mattress) was used. Intravenous fluids were warmed in thermostatic baths for water (Precisterm P Selecta) at a temperature of 40 °C.

Temperature was measured in the recovery room at 5 cm from the forehead skin surface using a PCE-FIT 10 (PCE Deutschland GmbH, accuracy ± 0.2 °C in the 36–39 °C range, and ± 0.3 °C in the 32–35.9 °C range, measurement range 32–42.4 °C), which is the device available in this area. The device was maintained and calibrated according to the manufacturer's instructions in order to obtain a temperature reading equivalent to central temperature.

Hypothermia was defined as a temperature equal to or lower than 35.9 °C at three levels: mild hypothermia, 35–35.9 °C; moderate hypothermia, 34–34.9 °C; and severe hypothermia, ≤ 33.9 °C.¹⁷

The sample was selected on a convenience basis. For the statistical tests, the null hypothesis of no difference between normothermic patients and patients with hypothermia on arrival to the recovery room was proposed. Quantitative variables were compared using the Student "t" test, and qualitative variables were compared using the Chi square test. After the comparison, a multivariate analysis (binary logistic regression) was applied, including variables for which a *p* value equal to or lower than 0.08 was obtained; moreover, polytomic variables were converted to the dichotomic form. Data were analyzed using the statistical package SPSS, version 12 (SPSS Inc., Chicago, IL) and an Excel worksheet, version 12 (Microsoft Corporation). All tests with a *p* < 0.05 were considered.

Results

Data were obtained from 200 consecutive patients. Of those, 33 were excluded because of incomplete information, for a final number of 167 included in the statistical analysis.

Hypothermia was observed in 56.29% of the patients (94/167), in 41.32% at the time of arrival at the recovery room, and in 14.97% 1 h into their stay in this area. Table 1 shows the course of the temperature. Of the patients who presented with hypothermia on arrival to the recovery room, 68 (40.72%) had mild hypothermia and 1 (0.6%), an 89-year-old patient, had moderate hypothermia. The age range was 17–89 years. When patient characteristics were compared in the bivariate analysis, significant (*p* < 0.02) differences were found between the normothermia and hypothermia groups in terms of age,

Table 1 – Presence of inadvertent hypothermia.

<i>Temperature assessment upon arrival at the recovery room</i>		
Hypothermia	69	41.32%
Normothermia	98	58.68%
Total patients	167	100.00%
<i>Course of temperature 1 h after arrival at the recovery room</i>		
Maintained hypothermia	41	24.55%
Temperature normalization	28	16.77%
New cases of hypothermia	25	14.97%
Maintained normothermia	73	43.71%
Total patients	167	100.00%
<i>Total number of cases with hypothermia^a</i>	94	56.29%

^a Hypothermia on arrival to recovery plus new cases within the first hour.

sex, obesity (BMI ≥ 30) and ASA classification. For the logistic regression, the surgical specialty and the type of anesthesia were also included because of a *p* = 0.08 (Table 2).

The following were the independent factors related to hypothermia resulting from the binary logistic regression analysis: age group greater or equal to 65 years, BMI greater or equal to 30 kg/m², and female gender, all with a *p* < 0.03 (Table 3).

Regarding intraoperative temperature management, it was found that temperature was monitored in 10% of patients and that the warming methods used were warm intravenous fluids in 78% and a forced-air warming system in 63% of patients, with no statistically significant difference between normothermic patients and those with hypothermia (Table 2). No patient with neuroaxial anesthesia was monitored intraoperatively. No relationship was found between temperature management measures and the presence of risk factors for hypothermia such as ASA and/or age extremes, or the use of warm fluids and/or warming with forced air.

During their stay in the recovery room, 9% (15/167) of the patients reported feeling cold and/or were found to be shivering, and they were managed using forced-air warming blankets. This method was used in one patient who was found to be hypothermic but with no clinical manifestations.

Discussion

This study found a high percentage (56.29%) of inadvertent hypothermia, a figure which is within the wide range of incidence variation reported in the literature between 6% and 90%.^{13–17} Of the variables considered in the study, only age ≥ 65 years, female gender and BMI ≥ 30 were found to be associated with hypothermia.

The bivariate analysis did not find a relationship between hypothermia and the time and type of anesthesia, the surgical specialty, intraoperative temperature monitoring, the use of warm intravenous fluids and the use of forced-air warming systems; additionally, the ASA classification was excluded as a result of the logistic regression. It is important to mention that some of those factors are considered in the recommendations for the prevention of inadvertent hypothermia.²³

It is known that body temperature is not homogenous and that central temperature is the best indicator for thermal

Table 2 – Characteristics of the subjects studied.

Variable	Normothermia n=98	Hypothermia n=96	P value
Age	50.3 ± 17.3	60.6 ± 16.3	<0.01
BMI (kg/m ²)	27.9 ± 5.9	30.3 ± 6.3	<0.02
Anesthesia time (min)	110.4 ± 57.7	124.6 ± 69.9	0.15
Temperature (°C) ^a	36.3 ± 0.25	35.6 ± 0.23	<0.01
Age group			<0.01
<65 years	71 (72%)	35 (51%)	
≥65 years	27 (28%)	34 (49%)	
Gender			<0.02
Male	46 (47%)	20 (29%)	
Female	52 (53%)	49 (71%)	
Obesity (BMI ≥ 30)			<0.02
Yes	23 (24%)	29 (42%)	
No	75 (77%)	40 (58%)	
ASA classification			<0.02
ASA I	23 (24%)	6 (9%)	
ASA II	55 (56%)	36 (52%)	
ASA III	19 (19%)	24 (35%)	
ASA IV	1 (1%)	3 (4%)	
Surgical specialty			0.08
Vascular surgery	8 (8%)	3 (4%)	
General surgery	24 (24%)	21 (30%)	
Maxillo-facial surgery	8 (8%)	2 (3%)	
Plastic surgery	5 (5%)	3 (4%)	
Gynecology	19 (19%)	10 (14%)	
Otolaryngology	7 (7%)	2 (3%)	
Traumatology	16 (16%)	24 (35%)	
Urology	11 (11%)	4 (6%)	
Type of anesthesia			0.08
General	60 (61%)	32 (46%)	
Neuroaxial	30 (31%)	26 (38%)	
Combined	5 (5%)	10 (14%)	
Sedation	3 (3%)	1 (1%)	
Intraoperative temperature monitoring			0.96
Yes	7 (10%)	10 (10%)	
No	63 (90%)	88 (90%)	
Use of warm intravenous fluids			0.30
Yes	51 (74%)	63 (64%)	
No	18 (26%)	35 (36%)	
Use of blanket/thermal mattress			0.65
Yes	42 (61%)	63 (64%)	
No	27 (39%)	35 (36%)	

^a On arrival at the recovery room. Values of the quantitative variables are given as mean ± standard deviation. Qualitative variables are expressed in absolute numbers and percentage.

status in humans. Temperature determination in the pulmonary artery is the gold standard, but it has the disadvantage of being invasive. Intraoperatively, acceptable semi-invasive monitoring sites are the nasopharynx, the esophagus and the

urinary bladder.¹⁹ In the systematic review of the literature, non-invasive oral measurement is valid and safe for central temperature determination,¹⁹ which would make it the best alternative in the conscious patient. Langham et al.²⁴

Table 3 – Logistic regression analysis.

Variable	OR	95% Confidence interval		P value
		Upper limit	Lower limit	
Age ≥ 65 years	2.588	1.238	5.411	0.012
BMI ≥ 30	2.414	1.163	5.012	0.018
Female gender	2.166	1.084	4.328	0.029

found that electronic oral temperature measurement was the most adequate for use in the postoperative period, followed by axillary temperature. Höcker et al.²⁵ showed that sublingual temperature measurement is a good practical method for monitoring perioperative temperature in both anesthetized and conscious patients.

In this study, an infrared skin thermometer was used to measure forehead temperature, because it was the device available to us. It is known that temperature in peripheral tissues depends on exposure, central temperature and vasomotor thermoregulation.¹² Axillary and skin temperature is prone to artifacts,²⁶ which is why it might not be the best option. Unlike what some authors suggest, the measurement was not adjusted with the central temperature²⁷⁻²⁹ (skin temperature 0.7°C lower than central temperature) because the equipment had been calibrated for first use with a central temperature measurement in accordance with the manufacturer's recommendation, and this calibration is stable for periods of one to three years.

Age over 60 years, female gender and high-level spinal block have been reported as risk factors for perioperative hypothermia, on the basis of weak evidence (level B, Class IIa or IIb). Risk factors with insufficient evidence (level C, Class IIa or IIb) include BMI under the normal value, normal BMI, length of the procedure, uncovered surgical area, duration of anesthesia and diabetes mellitus.³⁰ Among the factors associated with hypothermia found in this study, the variable of a BMI ≥ 30 did not correlate with the published data. It has been reported that greater body weight protects against central hypothermia.^{31,32} Fat in obese individuals has conductivity, which reduces heat loss through the skin and minimizes hypothermia.³² Moreover, the vasoconstriction threshold at low ambient temperature is high in obese patients.³³ In view of the above, the result found in this study might be related to the measurement method, given that the reduced loss of heat through the skin would be particularly intensified in obese patients.

On the other hand, NICE²³ has defined high-risk patients as those with two or more of the following factors: ASA greater than I, preoperative temperature below 36°C, combined anesthesia, intermediate or major surgery, and patients with cardiovascular history. They recommend measuring temperature 1 h before induction, every 30 min intraoperatively, and postoperatively upon arrival at the recovery room every 15 min until it reaches 36°C, and then every hour until it reaches 36.5°C.

In this study, no patient was pre-warmed. Pre-warming is a current recommendation,^{15,23,34,35} and it attenuates substantially the initial drop in temperature in the anesthetized patient, as it prevents redistribution heat losses. Until this technique is implemented, active intraoperative warming will continue to be the primary strategy to fight hypothermia. We may state that the absence of pre-warming and the low percentage of use of intraoperative temperature monitoring may explain the lack of a statistically significant difference between hypothermic and normothermic patients despite the use of warm intravascular fluids and of a forced-air warming system. When no pre-warming is used, intraoperative warming techniques, including the use of forced-air warming, fail to eliminate the initial drop in temperature.³⁴

Inadvertent hypothermia must be prevented. It is easier to maintain intraoperative normothermia than to rewarm patients in the postoperative period.³⁶ Intraoperatively, the patient is vasodilated and thermal transfer is easier than when the patient is in vasoconstriction, as is the case in the postoperative period. Peripheral vasoconstriction limits the flow of heat toward the peripheral compartment, increasing the gradient due to the accumulation of heat generated by tissue metabolism in the central compartment.³⁷

The two most important mechanisms responsible for heat loss in the operating room are radiation and convection, in that order. Radiation accounts for 60% of the losses, which is the reason why a relative humidity of >45% and a temperature ranging between 21 and 14°C must be maintained in the operating room for adult patients, and between 24 and 26°C for pediatric patients. Regarding this issue, the NICE guidelines state: "temperature in the operating room must be at least 21°C while the patient is exposed". ASPAN recommends maintaining operating room temperature between 20 and 25°C (Class I, Level C). Operating room temperature and humidity were not recorded in this study, and that may be an important factor in the occurrence of hypothermia associated with the type of surgery.²⁹

Forced-air warming, available since 1980, works on the principle of hot air infusion that escapes through small orifices pointing at the patient. It has been shown to be the only efficient method for maintaining temperature and warming the patients in the perioperative period.³⁸⁻⁴⁰ The efficacy of the system is enhanced by covering the blanket with a cotton sheet, and it has the advantage of being flexible, enabling optimal coverage of the skin surface, regardless of positioning. Reported complications with the Bair Hugger systems are rare, with one report of a third degree burn⁴¹ and one case of thermal softening of the tracheal tube.⁴²

It has been found that the administration of warm fluids and line warming are equally effective for preventing perioperative hypothermia.⁴³ Fluid warming does not warm the patient, but rather minimizes the incidence of perioperative hypothermia.⁴⁴ In machines that allow line warming, fluids are warmed to 38°C, but they have to be warmed to 41°C when cabinets are used; in both cases, when they reach the patient their temperature is 37°C. In this study, no patient received line-warmed fluids even though the resource was available. The ATLS (Advanced Trauma Life Support) manual of the American College of Surgeons recommends microwave warming of resuscitation fluids to 39°C. The 500 ml bags may be warmed at 400 W for 100 s or 800 W for 50 s. Consequently, an inexpensive alternative is to warm the fluids with microwaves.⁴⁵

The systematic implementation of perioperative temperature management is amply justified. The evidence supports starting active warming before the operation and temperature monitoring throughout the perioperative period in order to prevent hypothermia. The use of warming methods is supported by the evidence, but it is optimized only when temperature is monitored, considering that it is impossible to manage temperature if it is not measured.⁴⁶

In conclusion, warming measures without temperature monitoring fail to reduce the presence of hypothermia, contrary to what may be expected. Given the high incidence of

inadvertent hypothermia despite the availability of adequate resources for monitoring and managing temperature, there is a need to standardize and implement action guidelines to prevent and manage its occurrence including, among other measures, pre-warming and temperature monitoring before, during and after anesthesia for all patients, in particular in the groups at risk which, in this study, were patients ≥ 65 years and of the female gender.

Funding

Author's own resources

Conflict of interests

The authors have no conflict of interests to declare.

REFERENCES

- Roth JV. Some unanswered questions about temperature management. *Anesth Analg.* 2009;109:1695-9.
- Rajagopalan S, Mascha E, Na J, Sessler DI. The effects of mild perioperative hypothermia on blood loss and transfusion requirement. *Anesthesiology.* 2008;108:71-7.
- Tse EY, Cheung WY, Ng KF, Luk KD. Reducing perioperative blood loss and allogeneic blood transfusion in patients undergoing major spine surgery. *J Bone Joint Surg Am.* 2011;93:1268-77.
- Frank SM, Fleisher LA, Breslow MJ, Higgins MS, Olson KF, Kelly S, et al. Perioperative maintenance of normothermia reduces the incidence of morbid cardiac events. A randomized clinical trial. *JAMA.* 1997;277:1127-34.
- Melling AC, Ali B, Scott EM, Leaper DJ. Effects of preoperative warming on the incidence of wound infection after clean surgery: a randomised controlled trial. *Lancet.* 2001;358:876-80.
- Mauermann WJ, Nemergut EC. The Anesthesiologist's role in the prevention of surgical site infections. *Anesthesiology.* 2006;105:413-21.
- Heier T, Caldwell JE. Impact of hypothermia on the response to neuromuscular blocking drugs. *Anesthesiology.* 2006;104:1070-80.
- Heier T, Caldwell JE, Sessler DI, Miller RD. Mild intraoperative hypothermia increases duration of action and spontaneous recovery of vecuronium blockade during nitrous oxide-isoflurane anesthesia in humans. *Anesthesiology.* 1991;74:815-9.
- Leslie K, Sessler DI, Bjorksten AR, Moayeri A. Mild hypothermia alters propofol pharmacokinetics and increases the duration of action of atracurium. *Anesth Analg.* 1995;80:1007-14.
- Vitez TS, White PF, Eger E. Effects of hypothermia on halothane MAC and isoflurane MAC in the rat. *Anesthesiology.* 1974;41:80-1.
- Sessler DI. Complications and treatment of mild hypothermia. *Anesthesiology.* 2001;95:531-43.
- Sessler DI. Temperature monitoring and perioperative thermoregulation. *Anesthesiology.* 2008;109:318-38 [review].
- Harper CM, Andrzejowski JC, Alexander R. Nice and warm. *Br J Anaesth.* 2008;101:293-5.
- Knaepel A. Inadvertent perioperative hypothermia: a literatura review. *J Perioper Pract.* 2012;22:86-90.
- Moola S, Lockwood C. Effectiveness of strategies for the management and/or prevention of hypothermia within the adult perioperative environment. *Int J Evid Based Healthc.* 2011;9:337-45.
- Ross-Anderson DJ. A NICE idea or a high price to pay? Local assessment of national guidelines. *Anaesthesia.* 2009;64:330-1.
- De Mattia AL, Barbosa MH, De Mattia Rocha A, Farias HL, Santos CA, Santos DM. Hipotermia en pacientes en el período perioperatorio. *Rev Esc Enferm USP.* 2012;46:60-6.
- Karalapillai D, Story DA, Calzavacca P, Licari E, Liu YL, Hart GK. Inadvertent hypothermia and mortality in postoperative intensive care patients: retrospective audit of 5050 patients. *Anaesthesia.* 2009;64:968-72.
- Torossian A. Thermal management during anaesthesia and thermoregulation standards for the prevention of inadvertent perioperative hypothermia. *Best Pract Res Clin Anaesthesiol.* 2008;22:659-68.
- Hannenberg AA, Sessler DI. Improving perioperative temperature management. *Anesth Analg.* 2008;107:1454-7.
- Fleisher LA, Beckman JA, Brown KA, Calkins H, Chaikof E, Fleischmann KE, et al., ACC/AHA. Guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery. *J Am Coll Cardiol.* 2007;50:1707-32.
- American Society of Anesthesiologist Task Force on Postanesthetic Care. Practice guidelines for postanesthetic care: a report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology.* 2002;96:742-52.
- National Institute for Health and Clinical Evidence. Clinical practice guidelines: The management of inadvertent perioperative hypothermia in adults. Available from: www.nice.org.uk/CG65
- Langham GE, Maheshwari A, Contrera K, You J, Mascha E, Sessler DI. Noninvasive temperature monitoring in postanesthesia care units. *Anesthesiology.* 2009;111:90-6.
- Höcker J, Bein B, Böhm R, Steinfath M, Scholz J, Horn EP. Correlation, accuracy, precision and practicability of perioperative measurement of sublingual temperature in comparison with tympanic membrane temperature in awake and anaesthetized patients. *Eur J Anaesthesiol.* 2012;29:70-4.
- de Silva A. Anesthetic monitoring. In: *Basic of anesthesia.* 6th ed. Philadelphia: Elsevier Saunders Inc.; 2011. p. 817.
- American Society of Anesthesiologists. Standards for basic anesthetic monitoring; 2010. <http://asahq.org/For-Healthcare-Professionals/~media/For%20Members/documents/Standards%20Guidelines%20Stmnts/Basic%20Anesthetic%20Monitoring%202011.ashx> [accessed 13.01.11].
- Manuel Blanco-Pajón. ¿Es recomendable el monitoreo de la temperatura en los pacientes bajo anestesia? Implicaciones clínicas y anestésicas. *Rev Mex Anestesiología.* 2010;33:S70-5.
- Modell JG, Katholi CR, Kumaramangalam SM, Hudson EC, Graham D. Unreliability of the infrared tympanic thermometer in clinical practice: a comparative study with oral mercury and oral electronic thermometers. *South Med J.* 1988;91:649-54.
- Hooper VD, Chard R, Clifford T, Fetzer S, Fossum S, Godden B, et al. ASPAN's evidence-based clinical practice guideline for the promotion of perioperative normothermia: second edition. *J Perianesth Nurs.* 2010;25:346-65.
- Kongsayreepong S, Chaibundit C, Chadpaibool J, Komoltri C, Suraseranivongse S, Suwannanonda P, et al. Predictor of core hypothermia and the surgical intensive care unit. *Anesth Analg.* 2003;96:826-33.
- Fernandes LA, Braz LG, Koga FA, Kakuda CM, Módolo NS, de Carvalho LR, et al. Comparison of peri-operative core temperature in obese and non-obese patients. *Anaesthesia.* 2012;67:1364-9.

33. Kasai T, Hirose M, Matsukawa T, Takamata A, Tanaka Y. The vasoconstriction threshold is increased in obese patients during general anaesthesia. *Acta Anaesthesiol Scand*. 2003;47:588-92.
34. Hart SR, Bordes B, Hart J, Corsino D, Harmon D. Unintended perioperative hypothermia. *Ochsner J*. 2011;11:259-70.
35. Andrzejowski J, Hoyle J, Eapen G, Turnbull D. Effect of prewarming on post-induction core temperature and the incidence of inadvertent perioperative hypothermia in patients undergoing general anaesthesia. *Br J Anaesth*. 2008;101:627-31.
36. Díaz M, Becker DE. Thermoregulation: physiological and clinical considerations during sedation and general anesthesia. *Anesth Prog*. 2010;57:25-32.
37. Matsukawa T, Sessler DI, Sessler AM, Schroeder M, Ozaki M, Kurz A, et al. Heat flow and distribution during induction of general anesthesia. *Anesthesiology*. 1995;82:662-73.
38. Torrie JJ, Yip P, Robinson E. Comparison of forced-air warming and radiant heating during transurethral prostatic resection under spinal anaesthesia. *Anaesth Intensive Care*. 2005;33:733-8.
39. Leung KK, Lai A, Wu A. A randomised controlled trial of the electric heating pad vs forced-air warming for preventing hypothermia during laparotomy. *Anaesthesia*. 2007;62:605-8.
40. Fanelli A, Danelli G, Ghisi D, Ortu A, Moschini E, Fanelli G. The efficacy of a resistive heating under-patient blanket versus a forced-air warming system: a randomized controlled trial. *Anesth Analg*. 2009;108:199-201.
41. Truell KD, Bakerman PR, Teodori MF, Maze A. Third-degree burns due to intraoperative use of a Bair Hugger warming device. *Ann Thorac Surg*. 2000;69:1933-4.
42. Ayala JL, Coe A. Thermal softening of tracheal tubes: an unrecognized hazard of the Bair Hugger active patient warming system. *Br J Anaesth*. 1997;79:543-5.
43. Andrzejowski JC, Turnbull D, Nandakumar A, Gowthaman S, Eapen G. A randomised single blinded study of the administration of pre-warmed fluid vs. active fluid warming on the incidence of peri-operative hypothermia in short surgical procedures. *Anesthesia*. 2010;65:942-5.
44. Insler SR, Sessler DI. Perioperative thermoregulation and temperature monitoring. *Anesthesiol Clin*. 2006;24:823-37.
45. Lindhoff GA, Mac G, Palmer JH. An assessment of the thermal safety of microwave warming of crystalloid fluids. *Anaesthesia*. 2000;55:251-4.
46. Nierman DM. Tools that we use: if you can't measure it, you can't manage it. *Crit Care Med*. 2007;35:312-3.