



Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

www.revcolanest.com.co



Review

Post cardiac arrest syndrome[☆]



José Ricardo Navarro-Vargas^{a,*}, José Luis Díaz^b

^a Associate Professor, School of Medicine, Universidad Nacional de Colombia, Colombia

^b Assistant Professor, Department of Anesthesia and Critical Care, Mayo Clinic, FL, USA

ARTICLE INFO

Article history:

Received 29 July 2013

Accepted 20 January 2014

Available online 28 February 2014

Keywords:

Heart arrest

Cardiopulmonary resuscitation

Percutaneous coronary intervention

Hypothermia induced

Blood Circulation

ABSTRACT

Background: Resuscitation from cardiac arrest with global ischemia restores spontaneous circulation in some patients; however, survival depends on many factors associated with post cardiac arrest syndrome. During the last ten years, the understanding and control of these factors have improved the prognosis in a subgroup of patients.

Objective: To describe the pathophysiology and current management of the post cardiac arrest syndrome (PCAS).

Methodology: Narrative review of the literature using Medline via PubMed and Clinical Trials, using the terms MeSH cardiac arrest – Cardiopulmonary Resuscitation and (no term MeSH) Post cardiac arrest syndrome.

Results: Clinical trials have established a set of management protocols and guidelines based on therapeutic objectives with survival rates exceeding 50% of the cardiac arrest victims.

Conclusions: The management of this syndrome has actually strengthened the last link in the survival chain by standardizing the evaluation and selection of cardiac arrest victims via a therapeutic hypothermia protocol and early percutaneous coronary intervention.

© 2013 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L. All rights reserved.

Síndrome posparo cardiaco

RESUMEN

Antecedentes: La reanimación en el paro cardiaco con isquemia global logra restablecer la circulación espontánea en algunos pacientes; sin embargo, la sobrevida depende de muchos factores que explican el síndrome posparo cardiaco. El entendimiento y el control de estos factores durante la última década han logrado mejorar el pronóstico en un subgrupo de pacientes.

Objetivo: Describir la fisiopatología y el manejo actual del síndrome posparo cardiaco.

Metodología: Revisión narrativa de la literatura a través de las bases electrónicas de Medline vía PubMed y Ensayos Clínicos usando los términos MeSH Cardiac arrest – Cardiopulmonary Resuscitation y (el término no MeSH) Post cardiac arrest syndrome.

Palabras clave:

Paro cardiaco

Resucitación cardiopulmonar

Intervención coronaria percutánea

Hipotermia inducida

Circulación sanguínea

[☆] Please cite this article as: Navarro-Vargas JR, Díaz JL. Síndrome posparo cardiaco. Rev Colomb Anestesiolog. 2014;42:107-113.

* Corresponding author at: Universidad Nacional de Colombia, Calle 42 22 29, Bogotá, DC 0571, Colombia. Tel.: +57 3406593.

E-mail address: jnavarrovargas@hotmail.com (J.R. Navarro-Vargas).

Resultados: Los estudios clínicos han establecido una serie de protocolos y guías de manejo basadas en objetivos terapéuticos con tasas de supervivencia que superan el 50% de las víctimas de paro cardíaco.

Conclusiones: Actualmente el manejo de este síndrome ha fortalecido el último eslabón de la cadena de supervivencia al estandarizar la evaluación y la selección de víctimas de paro cardíaco con un protocolo de hipotermia terapéutica e intervención coronaria percutánea precoz.

© 2013 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

In 1972 the Russian pathophysiologist Vladimir Negovsky described the syndrome as “a post-resuscitation disease”.¹ However, since it involves a series of uncontrolled events, the International Liaison Committee on Resuscitation (ILCOR) adopted the term post cardiac arrest syndrome.²

The incidence rate due to all cardiac causes is 460,000 deaths/year.^{3,4} Prospective trials refer to 350,000 coronary disease-related deaths/year; this is 1–2/1000 people for the American population.⁵ There are survival reports of patients with extra-hospital cardiac arrest of 23.8% at the time of admission and of 7.6% at the time of discharge.⁶

Biological death depends on the cardiac arrest mechanism, on the underlying disease and on the delay in starting the resuscitation maneuvers (CPR). A poor neurological prognosis after 4–6 min of an unattended arrest is irreversible,⁷ and hence resuscitation must be a constant mission.⁸

When the arrest mechanism is an asystole or a pulseless electrical activity (PEA), the progression to neurological injury is faster and leads to a worse prognosis.⁷

Mild therapeutic hypothermia (32–34 °C) is the gold standard in post-arrest care.⁹

A narrative review of the literature using Medline via PubMed and Clinical Trials using the terms MeSH cardiac arrest – Cardiopulmonary Resuscitation and no term MeSH Post cardiac arrest syndrome was carried out.

Clinical evolution

Once the patient recovers spontaneous cardiac circulation, a cascade of events develops, mainly characterized by anoxic brain injury, post cardiac arrest myocardial dysfunction, “ischemic/reperfusion” systemic response, and the typical pathology of the triggering cause of the cardiac arrest. The clinical evolution shall be dependent on clinical conditions such as the patient’s co-morbidities, the duration of the ischemic lesion and the cause that triggered the cardiac arrest.¹⁰

Pathophysiology

Oxygen deficiency and generalized acidosis develop during cardiac arrest. If the victim is resuscitated using CPR/defibrillation maneuvers, with resumption of spontaneous circulation, PCSS develops, which is characterized by a systemic inflammatory response of the immune system and of

coagulation.¹¹ Cell damage seems to affect the enzyme calpain and peroxidation caused by oxygen free radicals that begin to develop during the phase of global ischemia and perpetuates during reperfusion.¹²

The main cause of extra-hospital cardiac arrest in the adult is acute myocardial infarction.¹³ There are many other pathologies leading to multisystem failure and subsequent cardiac arrest in the hospitalized patient.¹³

Treatment

According to the ILCOR^{2,10} document, the PCAS classification follows physiological criteria in five phases:

1. Immediate care: the initial 20 min following the patient’s spontaneous recovery of circulation.
2. Early phase: from 20 min to 6–12 h, when critical protective and therapeutic measures are required for a successful outcome.
3. Intermediate phase: from 6–12 h to 72 h a close surveillance and ICU treatment are required consistent with the therapeutic objectives.
4. Recovery phase: comprises the patient’s condition after the initial 72 h when there is a clearer diagnosis and a more predictable result.
5. Rehabilitation phase: focuses on the patient’s complete recovery. Any electrolytic abnormalities shall be corrected during phases 1 and 2, in addition to providing inotropic support and optimized oxygenation.¹⁴

Goal-targeted therapy

Ventilation support

1. Normocapnia (PaCO₂ between 40 and 45 mmHg). However, arterial gasometry should be properly interpreted in patients undergoing therapeutic hypothermia. When a patient reaches a central body temperature of close to 33 °C, the actual PaCO₂ may be up to 7 mmHg below the value in the arterial gases machine.¹⁵ Hyperventilation has been associated with a drop in coronary perfusion and venous return, in addition to cerebral vasoconstriction.¹⁶
2. Normoxia. Both, hypoxia and hyperoxia (PaO₂ > 300 mmHg) may result in secondary neurological injury. Using an inspired oxygen fraction to maintain the arterial saturation between 95% and 99% or a PaO₂ > 100 mmHg is

considered very reasonable.¹⁷ However, other authors have not achieved similar results.¹⁸

Hemodynamic support

Systemic perfusion and in particular cerebral perfusion have prognostic implications. Cardiac arrest patients experience a loss of cerebral self-regulation. Based on positron emission tomography studies, the suggestion is to maintain a mean pressure between 80 and 100 mmHg because this is the range in which perfusion matches the cerebral metabolic activity.¹⁹ During the first 48–72 h following the cardiac arrest, vasopressors and inotropic agents are usually needed. Transthoracic echocardiography may assist in accomplishing this goal. Apparently there is no difference in terms of the clinical prognosis regardless of the vasopressor (norepinephrine versus dopamine). Monitoring the mixed venous saturation may help in the interpretation of pharmacological interventions such as the introduction of inotropic agents.²⁰

Hypothermia has shown huge benefits since 2002,^{21,22} however, the procedure is not standardized in every country. There is an Italian study in ICUs indicating that only 16% used the therapeutic hypothermia protocol.²³

Hypothermia Management Scheme²⁴

1. The induction phase (body temperature between 32 °C and 34 °C). It should be started early to prevent neuroexcitotoxicity.²⁵ It has been shown that the benefit of hypothermia will not be achieved if it is introduced after 6 h of the spontaneous return of circulation.²⁶ Both PRINCE (Pre-ROSC Intra-nasal Cooling Effectiveness)²⁷ and Nagao²⁸ trials showed that initiating hypothermia before the patient recovers spontaneous cardiac circulation during CPR improves the neurological results and protects the myocardium from reperfusion injuries. Therapeutic hypothermia may not last less than 6 h and it should be continued for 12–24 h.²⁹

Some recommend starting a rapid saline solution infusion at 4 °C to accomplish these goals.³⁰ The rate at which the body temperature drops is close to 1 °C in 15 min, when administering 1 liter of cold saline solution. This method is thought to be comparable to or even better than the use of intravascular catheters.³¹ The infusion of 30 ml/kg of saline solution at 4 °C lowers the body temperature at a rate of over 2 °C/h.^{30,32} Chills are one of the side effects of cooling down because of the rise in metabolic oxygen consumption; the recommendation to prevent chills includes administering magnesium sulfate (5 g/5 h), sedation and proper analgesia, in addition to the occasional administration of muscle relaxants.^{13,24} A very common intravenous sedation regime is titrating with a continuous infusion of Propofol (up to 50 µg/kg/min) and Fentanyl. Hypothermia between 32 °C and 34 °C results in a drop in cardiac output (between 25% and 40%) at the expense of a heart rate decrease. Arrhythmias occur at much lower temperatures than these ranges or as a result of electrolyte imbalances.²⁴ The most destabilizing factors during this phase are hypovolemia and electrolyte imbalances

(hypokalemia and hypomagnesemia). Increased diuresis results in a hydro-electrolytic imbalance, hemoconcentration and a rise in blood viscosity.³³

2. The maintenance phase should be free of any temperature variations beyond 0.2–0.5 degrees centigrade of the temperature achieved. Be very careful with the metabolic requirements of the patient since these are reduced in up to 50% and monitor coagulation, although different trials do not show a significant risk of bleeding.^{24,34} Hypothermia extends the half-life of all drugs. The administration of muscle relaxants may suppress chills and hence prevent rises in body temperature. However, muscle relaxants may mask seizures (that are present in up to 44% of the cases).³⁵ The therapeutic approach includes the use of multiple anti-convulsive agents as is the case in status epilepticus (valproic acid, phenytoin, midazolam, phenobarbital and propofol) because seizures tend to be refractory.³⁶
3. The premise during the warm-up phase is to slowly recover the temperature at a rate of 0.2–0.3 degrees centigrade. It is recommended to start warming-up after 12–24 h of introducing the hypothermia until the temperature is normalized.^{13,24} It is usual to experience hyperkalemia, cerebral edema, and seizures during this phase; likewise, ruling out any risk of infection is a priority since one of the effects of hypothermia is the inhibition of the immune response. The use of antibiotic prophylaxis is only indicated for high-risk patients or in prolonged hypothermia (over 48 h).³⁷ The level of coagulopathy is considered to be mild when compared against that of cardiac arrest victims who are maintained at a normal temperature.³⁸ Increased infection rates have been described associated with therapeutic hypothermia for 24 h; however, it has not been associated with higher mortality.^{35,39} There can also be bradycardia, prolongation of the QTc segment, hyperglycemia, diuresis with subsequent hypokalemia, hypomagnesemia and hypophosphatemia.³⁹

Hypothermia protection

During the first phase, when the patient recovers spontaneous circulation, hypothermia reduces the metabolic consumption of oxygen and glucose.⁴⁰

During the second phase, hypothermia reduces the occurrence of excitatory amino acids, particularly glutamate; these amino acids are responsible for the activation of the cytotoxic cascade, the formation of reactive oxygen species (ROS) and nitric oxide.

During the third phase, hypothermia preserves the integrity of cell membranes interfering with the action of calpain and thus preventing the occurrence of cerebral edema, neuronal death and blood brain barrier injury.^{41–43}

The benefits of therapeutic hypothermia have been very well explained in a recent meta-analysis. Patients treated with therapeutic hypothermia exhibited improved neurological function (RR 1.55; 95 CI 1.22–1.96) and had a higher probability of survival at discharge (RR 1.35; 95 CI 1.10–1.65) compared to patients not treated with hypothermia.⁴⁴

The international agencies (AHA – ERC) have accepted and promoted the indication of hypothermia for managing the post cardiac arrest syndrome, not only in defibrillator-amenable patients, but also in cases of worse AESP prognosis and asystole in which the most important prognostic factor is the *initiation of therapy*.^{16,29}

Indications for therapeutic hypothermia^{16,45}

Return to spontaneous cardiac circulation following cardiac arrest (any type of rhythm)

Coma

Over 18 years old

Absolute contraindications

Non-compressible active bleeding

Do not resuscitate medical order (DNR)

The hypothermia protocol must be interrupted in the presence of sepsis or pneumonia, refractory hemodynamic instability and severe refractory arrhythmia.

Hypothermia in patients with a FV cardiac arrest has an NNT of 6 (confidence interval of 4-13). Despite the consistency of the trials in terms of the rhythm of presentation of the arrest, cooling method used, time to reach therapeutic hypothermia and duration thereof, no follow-up studies have been made beyond one year to establish the final neurological involvement.⁴⁶

Hypothermia should not be delayed, even when the patient requires transluminal percutaneous coronary angioplasty.⁴⁷

The best site to monitor the central temperature is through the pulmonary artery catheter or the central venous line; however, other sites are recommended such as the esophagus and the bladder, because they are easy to place and have minimal effects.^{13,48} The bladder temperature may be misinterpreted in the presence of oliguria.

When should you do percutaneous coronary intervention (PCI) and thrombolysis

Ask about any history of coronary heart disease, prior chest pain symptoms and initial arrest rhythm (if the QRS tracing in the ECG is widened, it is most likely of coronary etiology); however, the literature has not found a positive predictive value for coronary obstruction of these two parameters.⁴⁹

Thrombolysis is only recommended in isolated cases of pulmonary embolism and acute myocardial infarction with ST elevation (IAMST) and a 3-h therapeutic window or in the case of IAMST, with no other option for invasive coronary intervention.⁵⁰

In cardiac arrest patients with no evident extra cardiac cause or who present ST elevation in the ECG or sudden left branch blockade, an early angiographic exploration is recommended, followed by percutaneous coronary intervention.⁵⁰ In a report written by Spaulding et al.,⁵¹ 49% of all cardiac arrest survivors who underwent routine coronary angiography showed a severe coronary obstruction.

Further randomized clinical trials are needed showing the positive impact of emergency percutaneous coronary intervention in this syndrome. As of now, the studies reporting benefits with the use of PCI-associated therapeutic

hypothermia exhibit various biases and undetermined confounding factors.⁵²

The similarity of the cardiohemic involvement of PCAS³³ to septic shock has led to the search of alternative aggressive management options with quite successful results such as circulatory support with extracorporeal oxygenation⁵³ and renal replacement therapy with high volume hemofiltration.⁵⁴

In refractory cardiac arrest patients, the alternative to use venous-arterial extracorporeal membrane oxygenation and PCI has been suggested.⁵²

Recommendation for neurological follow-up of the patient

The brain is the key organ in resuscitation, hence the high importance given to therapeutic hypothermia.⁵⁵ There is an interesting article on the “self-fulfilling prophecy”⁵⁶ that states that no negative prior judgments should be made in a patient who has overcome the arrest, regardless of the cause because by not doing what should be done, the patient will not survive, giving the false impression that what is appropriate is not to start any medical intervention under these circumstances. However, there is always room for uncertainty in every prognosis.⁵⁶

The prognosis after the clinical and paraclinical exams at different points in the evolution must have a 72-h (3 days) period of observation before any decision is made.⁵⁷ The American Academy of Neurology suggests that in the absence of brain death, findings of the clinical examination such as absence of corneal or pupillary reflexes, of motor response or extended absence of motor response, lead to a high rate or poor prognosis and consequently low false positives.⁵⁸

Efforts have been made to identify serum markers for predicting the development of multiple organ failure in PCAS patients. Soluble Ang1 factor stabilizes the endothelium Ang2 levels, and a higher proportion of Ang2/Ang1 is predictive of multiple organ failure and poor prognosis.⁵⁹

High procalcitonin levels are shown to be associated with PCAS and adverse neurological prognosis rather than to early infection in patients with anoxic encephalopathy.⁶⁰ Among the clinical factors themselves, a research comparing two hospitals in distant geographical regions showed that the pre- and intra-cardiac arrest conditions determine the severity of the PCAS.⁶¹

Prognostic factors¹³

- Patient history: old age – diabetes – sepsis – CVA
- Cardiac arrest and resuscitation conditions: time elapsed since the cardiac arrest and the start of CPR – CPR quality – asystole as arrest rhythm – implementation of a protocol for managing the PCAS.⁶²
- Later evolution in accordance with the neurological examination, neurophysiological exams, neuroimaging, biochemical markers and intracranial Doppler.

The two scales used to establish the degree of disability and the quality of life are the Glasgow prognostic scale¹³ and the Glasgow-Pittsburgh¹³ scale for categorizing brain performance.

What is currently being done?

A large international prospective trial was completed coordinated from Sweden⁶³ – the TTM trial (Target Temperature Management, identified as NCT01020916), whose results were published late in 2013. The purpose of the trial was to establish the level of evidence of hypothermia for the management of PCAS with optimal control to avoid hyperthermia in both groups; n: 939 adult patients. Comparison: controlled hypothermia (33 °C) versus patients maintained with a temperature of 36 °C. Period of time during which the patients were evaluated: up to 180 days. The results obtained are controversial with regard to the optimal threshold of hypothermia. The conclusion of the trial was based on a strict temperature control (below 36 °C) to prevent hyperthermia, which is considered a determining factor for poor prognosis.⁶⁴ Emphasis was also placed on the establishment of standardized protocols, even for each subgroup of patients who may benefit from individualized treatment.⁶⁵

The recommendation is to optimize the perfusion of vital organs in addition to the brain, since the prevalence of multiple extra-cerebral organ failure impacts mortality, particularly the involvement of the cardiovascular (depends on the fine vasopressor support) and respiratory system (by providing improved oxygenation). An independent association has been identified between these two organs and their involvement has been linked to a higher hospital mortality in PCAS.⁶⁶

Conclusion

PCAS is an entity that has been recognized for the last 40 years; however, the treatment for PCAS has only been established in an organized manner in the last decade.

The pathophysiology includes a serious involvement of the brain and leads to myocardial dysfunction and systemic inflammation. Failure to intervene early on results in irreversible brain injury, vegetative state and death.

Funding

Authors.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. Negovsky VA. The second step in resuscitation: the treatment of the "post-resuscitation disease". *Resuscitation*. 1972;1:1-7.
2. Neumar RW, Nolan JP, Adrie C, Aibiki M, Berg RA, et al. A consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation, Resuscitation Council of Asia, and The Resuscitation Council of Southern Africa); The American Heart Association Emergency Cardiovascular Care Committee; The Council on Cardiovascular Surgery and Anesthesia; The Council on Cardiopulmonary, Perioperative, and Critical Care; The Council on Clinical Cardiology; and The Stroke Council: Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. *Circulation*. 2008;118:2452-83.
3. Cobb LA, Fahrenbruch CE, Olsufka M, Copass MK. Changing incidence of out-of-hospital ventricular fibrillation, 1980-2000. *J Am Med Assoc*. 2002;288:3008.
4. Rosamond W, Flegal K, Furie K, Go A, Greenlund K, Haase N, et al. American Heart Association Statistics Committee and Stroke Statistics Subcommittee: Heart Disease and Stroke Statistics—2008 update: a report from the AHA Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2008;117:e25.
5. Myerburg RJ. Sudden cardiac death: exploring the limits of our knowledge. *J Cardiovasc Electrophysiol*. 2001;12:369.
6. Sasson C, Rogers MA, Dahl J, Kellermann AL. Predictors of survival from out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes*. 2010;3:63-81.
7. Myerburg RJ, Castellanos A. Cardiac arrest and sudden cardiac death. In: Bonow: Braunwald' heart disease. A textbook of cardiovascular medicine. 9th ed. Saunders; 2012. p. 845-84.
8. Navarro-Vargas JR. Sobre la reanimación: una misión permanente. *Rev Colomb Anestesiología*. 2008;36:9-10.
9. Derwall M, Fries M. Advances in brain resuscitation: beyond hypothermia. *Crit Care Clin*. 2012;28:271-81.
10. Stub D, Bernard S, Duffy SJ, Kaye D. Post cardiac arrest syndrome. *Circulation*. 2011;123:1428-35.
11. Laver S, Farrow C, Turner D, Nolan J. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med*. 2004;30:2126-8.
12. Globus MY, Busto R, Lin B, Schnipper H, Ginsberg MD. Detection of free radical activity during transient global ischemia and recirculation: effects of intraschemic brain temperature modulation. *J Neurochem*. 1995;65:1250-6.
13. Martín-Hernández H, López-Messa JB, Pérez-Vela JL, Molina-Latorre R, Cárdenas-Cruz A. Manejo del Síndrome posparada cardiaca. Documento de consenso. *Med Intensiva*. 2010;34:107-26.
14. Binks A, Nolan JP. Post-cardiac arrest síndrome. *Minerva Anestesiología*. 2010;76:362-8.
15. Sitzwohl C, Kettner SC, Reinprecht A, Dietrich W, Klimscha W, Fridrich P, et al. The arterial to end-tidal carbon dioxide gradient increases with uncorrected but not with temperature-corrected PaCO₂ determination during mild to moderate hypothermia. *Anesth Analg*. 1998;86:1131.
16. Peberdy MA, Callaway CW, Neumar RW, Geocadin RG, Zimmerman JL, Donnino M, et al. Part 9: post-cardiac arrest care: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010;122 18 Suppl 3:S768.
17. Janz DR, Hollenbeck RD, Pollock JS, McPherson JA, Rice TW. Hyperoxia is associated with increased mortality in patients treated with mild therapeutic hypothermia after sudden cardiac arrest. *Crit Care Med*. 2012;40:3135.
18. Bellomo R, Bailey M, Eastwood GM, Nichol A, Pilcher D, Hart GK, et al., Study of Oxygen in Critical Care (SOCC) Group. Arterial hyperoxia and in-hospital mortality after resuscitation from cardiac arrest. *Crit Care*. 2011;15:R90.
19. Schaafsma A, de Jong BM, Bams JL, Haaxma-Reiche H, Pruim J, Zijlstra JG. Cerebral perfusion and metabolism in resuscitated patients with severe post-hypoxic encephalopathy. *J Neurol Sci*. 2003;210:23.
20. Gaieski DF, Band RA, Abella BS, Neumar RW, Fuchs BD, Kolansky DM, et al. Early goal-directed hemodynamic optimization combined with therapeutic hypothermia in

- comatose survivors of out-of-hospital cardiac arrest. *Resuscitation*. 2009;80:418.
21. Walters JH, Morley PT, Nolan JP. The role of hypothermia in post-cardiac arrest patients with return of spontaneous circulation: a systematic review. *Resuscitation*. 2011;82:508-16.
 22. Holtzer M, Cerchiari E, Martens P, Roine R, Sterz F, Eisenburger P, et al. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. The Hypothermia After Cardiac Arrest Study Group. *N Engl J Med*. 2002;346:549-56.
 23. Bianchin A, Pellizzato N, Martano L, Castioni CA. Therapeutic hypothermia in Italian intensive care units: a national survey. *Minerva Anestesiologica*. 2009;75:357-62.
 24. Schaller B, Graf R. Hypothermia and stroke: the pathophysiological background. *Pathophysiology*. 2003;10:7-35.
 25. Kuboyama K, Safar P, Radovsky A, Tisherman SA, Stezoski SW, Alexander H, et al. Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: a prospective, randomized study. *Crit Care Med*. 1993;21:1348-58.
 26. Wolff B, Machill K, Schumacher D, Schulzki I, Werner D. Early achievement of mild therapeutic hypothermia and the neurologic outcome after cardiac arrest. *Int J Cardiol*. 2009;133:223.
 27. Elste T, Roessler M, Fritz H, Durnez P, Busch HJ, Inderbitzen B, et al. Study PRINCE: Pre-ROSC Intranasal Cooling Effectiveness. Intra-Arrest transnasal evaporative cooling: a randomized, prehospital, multicenter. *Circulation*. 2010;122:729-36.
 28. Nagao K. Therapeutic hypothermia following resuscitation. *Curr Opin Crit Care*. 2012;18:239-45.
 29. Bernard S. Hypothermia after cardiac arrest: expanding the therapeutic scope. *Crit Care Med*. 2009;37:227-33.
 30. Kim F, Olsufka H, Carlbom D, Deen S, Longstreth WT, Hanrahan M, et al. Pilot randomized clinical trial of prehospital induction of mild hypothermia in out-of-hospital cardiac arrest patients with a rapid infusion of 4 degrees C normal saline. *Circulation*. 2007;115:3064-70.
 31. Al-Senani FM, Graffagnino C, Grotta JC, Saiki R, Wood D, Chung W, et al. A prospective, multicenter pilot study to evaluate the feasibility and safety of using the CoolGard System and Icy catheter following cardiac arrest. *Resuscitation*. 2004;62:143.
 32. Tømte Ø, Drægni T, Mangschau A, Jacobsen D, Auestad B, Sunde K. A comparison of intravascular and surface cooling techniques in comatose cardiac arrest survivors. *Crit Care Med*. 2011;39:443.
 33. Adrie C, Monchin M, Cariou A, Dhainaut J-F, Spalding C. Postresuscitation disease after cardiac arrest: a sepsis-like syndrome? *Curr Opin Crit Care*. 2004;10:208-12.
 34. Valeri CR, MacGregor H, Cassidy G, Tinney R, Pompei F. Effects of temperature on bleeding time and clotting time in normal male and female volunteers. *Crit Care Med*. 1995;23:698-704.
 35. Nielsen N, Sunde K, Hovdenes J, Riker RR, Rubertsson S, Ståmmet P, et al. Hypothermia Network. Adverse events and their relation to mortality in out-of-hospital cardiac arrest patients treated with therapeutic hypothermia. *Crit Care Med*. 2011;39:57.
 36. Agarwal P, Kumar N, Chandra R, Gupta G, Antony AR, Garg N. Randomized study of intravenous valproate and phenytoin in status epilepticus. *Seizure*. 2007;16:527.
 37. Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med*. 2002;346:557-63.
 38. Perbet S, Mongardon N, Dumas F, Bruel C, Lemiale V, Mourvillier B, et al. Early-onset pneumonia after cardiac arrest: characteristics, risk factors and influence on prognosis. *Am J Respir Crit Care Med*. 2011;184:1048-54.
 39. Polderman KH. Is therapeutic hypothermia immunosuppressive? *Crit Care*. 2012;16 Suppl 2:A8.
 40. Nolan JP, Morley PT, Van den Hoek TL, Hickey RW, Kloock WG, Billi J, et al. Therapeutic hypothermia after cardiac arrest: an advisory statement by the advance life support task force of the International Liaison Committee on Resuscitation. *Circulation*. 2003;108:118-21.
 41. Safar P. Cerebral resuscitation after cardiac arrest: a review. *Circulation*. 1986;74:138-53.
 42. Sunde K, Pytte M, Jacobsen D, Mangschau A. Implementation of standardised treatment protocol for postresuscitation care after out-of-hospital cardiac arrest. *Resuscitation*. 2007;73:29-39.
 43. Beccaria PF, Turi S, Cristofolini M, Leggieri C, Vinciguerra F, Zangrillo A. Postcardiac arrest therapeutic hypothermia in adult patients, state of art and practical considerations. *HSR Proc Intensive Care Cardiovasc Anesth*. 2010;2:93-103.
 44. Arrich J, Holzer M, Havel C, Müllner M, Herkner H. Hypothermia for neuroprotection in adults after cardiopulmonary resuscitation. *Cochrane Database Syst Rev*. 2012;9:CD004128.
 45. Hassani H, Meyer S. Hypothermia for neuroprotection in adults after cardiopulmonary resuscitation. *Cochrane for clinicians*, accessed from: <http://www.aafp.org/afp/2010/0901/p477.pdf>. Accessed february 8th, 2014.
 46. Holzer M, Bernard SA, Hachimi-Idrissi S, Roine RO, Sterz F, Müllner M, on behalf of the Collaborative Group on Induced Hypothermia for Neuroprotection After Cardiac Arrest: hypothermia for neuroprotection after cardiac arrest: systemic review and individual patient data meta-analysis. *Crit Care Med*. 2005;33:414-8.
 47. Batista LM, Lima FO, Januzzi JL, Donahue V, Snyderman C, Greer DM, et al. Feasibility and safety of combined percutaneous coronary intervention and therapeutic hypothermia following in cardiac arrest. *Resuscitation*. 2010;81:398-403.
 48. Polderman KH, Herold I. Therapeutic hypothermia and controlled normothermia in the intensive care unit: practical considerations, side effects, and cooling methods. *Crit Care Med*. 2009;37:1101-20.
 49. Dumas F, Cariou A, Manzo-Silberman S, Grimaldi D, Vivien B, Rosencher J, et al. Immediate percutaneous coronary intervention is associated with better survival after out-of-hospital cardiac arrest: insights from the PROCAT (Parisian Region Out of hospital Cardiac Arrest) registry. *Circ Cardiovasc Interv*. 2010;3:200-7.
 50. Charalampopoulos A, Nikolaou N. Emerging pharmaceutical therapies in cardiopulmonary resuscitation and post-resuscitation syndrome. *Resuscitation*. 2011;82:371-7.
 51. Spaulding CM, Joly LM, Rosenberg A, Monchi M, Weber SM, Dhainaut JF, Carli P, et al. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med*. 1997;336:1629-33.
 52. Kagawa E, Keigo D, Yasuki K. Percutaneous coronary intervention for postcardiac arrest syndrome. *Curr Opin Cardiol*. 2013;28:689-94.
 53. Megarbane B, Leprince P, Deye N, Resiere D, Guerrier G, Rettab S, et al. Emergency feasibility in medical intensive care unit of extracorporeal life support for refractory cardiac arrest. *Intensive Care Med*. 2007;33:758-64.
 54. Laurent I, Adrie C, Vinsonneau C, Cariou A, Chiche JD, Ohanessian A, et al. High volume hemofiltration after out-of-hospital cardiac arrest: a randomized study. *J Am Coll Cardiol*. 2005;46:432-7.
 55. Mongardon N, Dumas F, Ricome S, Grimaldi D, Hissem T, Pène F, et al. Postcardiac arrest syndrome: from immediate

- resuscitation to long-term outcome. *Ann Intensive Care*. 2011;1:45-56.
56. Hemphill JC, White DB. Clinical nihilism in neuroemergencies. *Emerg Med Clin N Am*. 2009;27: 27-37.
 57. Levy DE, Caronna JJ, Singer BH, Lapinsky RH, Frydman H, Pulm F, et al. Predicting outcome from hypoxic-ischemic coma. *J Am Med Assoc*. 1985;253:1420-6.
 58. Wijdicks EFM, Hijdra A, Young GB, Bassetti CL, Wiebe S. Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2006;67:203-10.
 59. Wada T, Jesmin S, Gando S, Yanagida Y, Mizugaky A, Sultana SN, et al. Angiogenic factors and their soluble receptors predict organ dysfunction and mortality in post-cardiac arrest symbol. *J Inflamm (Lond)*. 2013. <http://www.journal.inflammation.com/content/10/1/6>. Accessed february 8th, 2014.
 60. Annborn M, Dankiewicz J, Erlinge D, Hertel S, Rundgren M, Smith JG, et al. Procalcitonin after cardiac arrest – an indicator of severity illness, ischemia-reperfusion injury and outcome. *Resuscitation*. 2013;84:782-7.
 61. Donnino MW, Miller JD, Bivens M, Cocchi MN, Saliccioli J, Farris S, et al. A pilot study examining the severity and outcome of the Post-cardiac arrest syndrome: a comparative analysis of two geographically distinct hospitals. *Circulation*. 2012;126:1478-83.
 62. Kellum MJ, Kennedy KW, Barney R, Keilhauer FA, Bellino M, Zuercher M, Ewy GA, et al. Cardiocerebral resuscitation improves neurologically intact survival of patients with out-of-hospital cardiac arrest. *Ann Emerg Med*. 2008;52:244-52.
 63. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, et al. Target temperature management after out-of-hospital cardiac arrest – a randomized, parallel-group, assessor-blinded clinical trial-rationale and design. *Am Heart J*. 2012;163:541-8 [Clinicaltrials.gov identifier: NCT01020916].
 64. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, et al. Targeted temperature management at 33 °C versus 36 °C after cardiac arrest. *N Engl J Med*. 2013;369:2197-206.
 65. Rittenberger JC, Callaway CW. Temperature management and modern post-cardiac arrest care. *N Engl J Med*. 2013;369:1557-8 [Editorial].
 66. Roberts BW, Kilgannon JH, Chansky ME, Mittal N, Wooden J, Parrillo JE, et al. Multiple organ dysfunction after return of spontaneous circulation in postcardiac arrest syndrome. *Crit Care Med*. 2013;41:1492-501.