



Review

Anaesthesia and the elderly patient, seeking better neurological outcomes[☆]



CrossMark

Cristina Alexandra Benavides-Caro*

A Anaesthetist Physician, Specialist in Bioethics, Adjunct Associate Professor in Neuroanaesthesia, Fundación Universitaria Sanitas, Bogotá, DC, Colombia

ARTICLE INFO

Article history:

Received 21 April 2015

Accepted 2 February 2016

Available online 3 April 2016

Keywords:

Aged

Anesthesia

Postoperative complications

Anesthesia, conduction

General surgery

ABSTRACT

Introduction: Elderly patients in need of surgery under anaesthesia present the challenge of maintaining their especially delicate balance under stress conditions.

Objective: This review of the literature aims to bring the anaesthesiologist closer to this group of patients, examining some of the physiological changes that occur with ageing, the effects that anaesthesia might have on their postoperative neurological state, the monitoring options, and the most frequent neurological complications.

Materials and methods: A search was conducted in the LILACS, MEDLINE and GOOGLE SCHOLAR databases using the terms anaesthesia, elderly, outcomes and neurologic, together with an ambispective snowball search from 2000 until today.

Conclusion: Institutions and staff responsible for the care of elderly patients must have the training and knowledge necessary for providing comprehensive treatment to this group of patients, in order to help them maintain their independence and physical and mental health which are so important during the final stage of life.

© 2016 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. All rights reserved.

Anestesia y paciente anciano, en busca de mejores desenlaces neurológicos

R E S U M E N

Palabras clave:

Anciano

Anestesia

Complicaciones posoperatorias

Anestesia de conducción

Cirugía general

Introducción: El paciente anciano llevado a cirugía y anestesia es un desafío al intentar mantener su delicado equilibrio bajo condiciones de estrés.

Objetivos: Esta revisión de la literatura tiene como objetivo aproximar al anestesiólogo a este grupo de pacientes al examinar algunos de los cambios fisiológicos que se presentan con el envejecimiento, los efectos que la anestesia puede tener sobre el estado neurológico postoperatorio; las opciones de monitorización y las complicaciones neurológicas más frecuentes.

* Please cite this article as: Benavides-Caro CA. Anestesia y paciente anciano, en busca de mejores desenlaces neurológicos. Rev Colomb Anestesiol. 2016;44:128-133.

* Correspondence to: Calle 23 # 66 – 46, consultorio 403-404, Colombia.

E-mail address: alexb36@gmail.com

Materiales y Métodos: Para este artículo se realizó una búsqueda en las bases de datos LILACS, MEDLINE Y GOOGLE SCHOLAR con los siguientes términos anestesia, anciano, desenlaces y neurológico, que se complementó con una búsqueda en bola de nieve ambispectiva a partir del año 2000 hasta la actualidad.

Conclusión: Las instituciones y personal a cargo del paciente anciano deben contar con la preparación y conocimiento necesarios para el manejo de este grupo de pacientes, con el fin de mantener la independencia y salud física y mental tan importantes en la etapa final de la vida.

© 2016 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

With improved medical care and health conditions, average life expectancy has increased throughout the world.¹ This increase in longevity results in a growing number of patients over 65 that are taken to surgery.^{2,3}

When managing this group of patients it is important to remember that the definitions of ageing and elderly are in constant flux and depend on the environment in which the elderly individual lives. The characteristics used as the basis to define what an elderly individual is may be associated with multiple social factors, including the age at which he or she stops being fully active at work or in society, is forced to change roles, or has lost functional capacities.³ The definition of the healthy elderly individual must also be considered, including several concepts such as independence for activities of daily living, wellness, social participation and good quality of life.⁴ It is generally accepted that an elderly patient is 65 or older.⁵

Elderly patients are complex when it comes to the approach to their management, and it is important to take into account that outcome goals are different than for the rest of the population due to the functional implications that a surgery may have for this group of people.⁶ An important percentage of patients over 65 years of age exhibit preclinical stages of Alzheimer's disease, Parkinson's disease, Lewy body dementia or cerebrovascular disease⁷; and this, together with the physiological changes and the effects of surgical stress and anaesthesia, increases the risk of developing postoperative complications when compared with middle aged individuals.⁸

Methods

For this review, a search was conducted in the LILACS, MEDLINE and GOOGLE SCHOLAR databases, using the terms anaesthesia, elderly, outcomes and neurological, together with an ambispective snowball search between 2000 and today.

Most frequent postoperative complications in elderly patients

The frequency of complications in elderly patients is very high. In the United States, 30% of the patients living in retirement

homes who were taken to colectomy died within the first three months after surgery, and 40% of those who survived showed significant functional decline⁹; the incidence of complications increases with age, and 20% of patients over 80 years of age taken to surgery had a complication, and the presence of a complication increased mortality, with a range between 4% and 26%.¹⁰

The above implies that, in the elderly, the healthcare team is mandated to have a very clear indication for surgery and to give priority to the patient's decision regarding his/her disease and the management options, over those of the family. Close to three-fourths of critically ill patients living in the United States prefer not to undergo any intervention that may entail the risk of decline or loss of their ability to function.⁹

Two of the most frequent postoperative neurological complications with a greater impact on patient outcome are delirium and cognitive dysfunction.

Postoperative delirium is defined as an acute disturbance of consciousness with signs of inattention, disorientation and memory abnormalities that fluctuate over time.¹¹ It is independently associated with increased mortality, length of stay, functional decline, and overall costs for the institutions. It usually manifests between the first and third postoperative days, with an incidence ranging between 5% and 15%, and may be as high as 33–62% in some groups such as that of hip fractures. The exact cause of this syndrome is yet unknown but it appears to be influenced by several combined factors such as the use of benzodiazepines, opioids or steroid medications, the level of schooling, the type of surgery, the depth of anaesthesia, and comorbidities such as terminal kidney disease or diabetes.¹²⁻¹⁴

Postoperative cognitive dysfunction is a term used to describe a syndrome characterized by consciousness, attention, perception, thinking, memory, behaviour and emotional disturbances that manifest after surgery. This complication appears within days or weeks, as early as within the first 7 days or as late as within the first 3 months. Incidence is highly variable and it is highly significant in cardiac surgery, ranging between 30% and 80%, and in major non-cardiac surgery, ranging between 25% for early stage and 9.9% for late stage. The causes of postoperative cognitive dysfunction are not clear and it appears to be the result of a combination of factors.^{15,16} In their study, Monk et al. found age to be an independent risk variable for postoperative cognitive dysfunction and also found a correlation between mortality within the first year after surgery and postoperative cognitive dysfunction. The

presence of late cognitive dysfunction is an important predictor of mortality.¹⁷

There are various tests that can be used easily and quickly during the consultation and after surgery in order to identify these two syndromes¹⁸; unfortunately, the staff responsible for the patients are not adequately trained to make the assessment, and reports on the frequency of missed delirium range between 33% and 66%.¹⁹

Pathophysiology of neurologic complications

In elderly patients, depending on their preoperative reserve and comorbidities, normal response to surgical stress may be altered, leading to dysfunction of haemodynamic, endocrine and immune responses.²⁰ Because of changes in their physiology, elderly patients are more prone to neurological complications. These changes include reduced cortical thickness of the brain, in particular in the prefrontal and parietal areas which are critical for memory and orientation, reduced synaptic junctions and dendritic trees in the pyramidal cells of the prefrontal area which is apparently the area with the highest cortical integration with thalamic nuclei; reduced number of neurotransmitters and neuroreceptors, which has been shown to be a risk factor for depression and neurodegenerative diseases like Alzheimer's; and increased susceptibility to the effects of oxidative stress and to the side effects of inflammatory processes.²¹

There is a relationship between ageing and the development of chronic diseases such as Type 2 diabetes, cancer, heart and renal diseases, all of them associated with postoperative complications.²² In elderly and middle-aged patients with diabetes and arterial hypertension, there is a deterioration of the periventricular and frontal white matter.²³

Associated with these changes are social factors such as abandonment and isolation, which have a powerful impact on the overall surgical outcome. These factors may mean that an important percentage of elderly individuals may present with malnutrition, inadequate disease management, and cognitive alterations.^{24,25}

Frailty

The effect of frailty must be considered in elderly patients. Frailty is defined as a state of diminished physiological reserve and vulnerability to the effects of stress; when frail patients are exposed to stressful events, they exhibit disproportionate decompensation. A frail patient taken to surgery has increased postoperative adverse effects, prolonged hospital stay, functional decline, disability and increased mortality.²⁶

The pathophysiology of frailty has not been well established but it appears to be related to inflammation, excess oxidative stress and a combination of immune and physiological alterations. Frail patients appear to have dysregulation of the immune, hormonal and endocrine systems, with higher levels of inflammatory cytokines as well as of inflammation biomarkers such as interleukin 6 and C reactive protein.^{26,27} These markers have also been associated with the presence of delirium and cognitive dysfunction and are measured more

frequently in elderly and frail patients.^{28,29} Leung found a relation between delirium and frailty in elderly patients during the early postoperative period.³⁰ The incidence of frailty increases with age, and frailty and age have been found to be independent factors associated with postoperative complications.³¹ In the population of elderly surgical patients, a significant incidence is found in 41.8–50.3% of cases, highlighting the vulnerability of this population during surgery.³² For Colombia, there is no accurate figure of the percentage of frail patients taken to surgery. Curcio, in a study conducted in a rural population over 60 years of age, found 15.2% of frail patients, a figure that is quite similar to that found in other studies conducted in Latin America.³³

Management of frail patients requires the integration of the entire treating team as well as institutional preparedness, including includes information and training on the implications and risks of this syndrome.³⁴

Effects of anaesthesia

A relationship has been found between anaesthetics, surgery and cognitive dysfunction in several in vitro and animal studies that have shown effects on neuronal development. It has been found that volatile and intravenous agents used for general anaesthesia appear to have a negative effect on neuronal structures, such as altered axonal growth and formation of neuronal networks, deranged axonal growth, apoptosis, telomere abnormalities, and alterations in critical areas for memory.³⁵

The study on late postoperative cognitive dysfunction, ISPOCD1, concluded unequivocally that anaesthesia and surgery produce long-term cognitive decline in elderly patients and that the risk increases with age, although it was unable to explain the pathophysiological causes and the anaesthetic or surgical events that may trigger this complication.¹⁶

In a review on anaesthesia and neurotoxicity, Lin et al. did not find an age of greater susceptibility to the deleterious neurological effects of anaesthesia, although they did find that the most affected neuronal groups change as age increases.³⁶ In a study that assessed the effects of desflurane on cognitive performance in rats, Callaway et al. found alterations in memory and in the ability to learn, and observed that those alterations were age- and dose-dependent and manifested in adult rats but not so in young individuals.³⁷

On the other hand, a relationship has been found between general anaesthesia administration and Alzheimer's disease. There are two histopathological markers in this disease, namely the extracellular amyloid plaques consisting of the beta amyloid peptide, and neurofibrillary interneuronal tangles made of an aberrant hyperphosphorylated Tau protein. Studies have shown that anaesthesia plays an important role in accelerating the pathogenesis of the Tau protein, which is influenced by hypothermia. Although this is not the definitive factor for the presentation of the aberrant form – equally demonstrated in normothermia – it does produce a marked increase in protein hyperphosphorylation. Although the relationship between hypothermia, increase in aberrant Tau protein and postoperative cognitive dysfunction is not

clear, studies in mice have found memory disorders in animals exposed to hypothermia and general anaesthesia.³⁸

Regional anaesthesia

Several studies on the incidence of postoperative neurological complications in elderly patients have not found differences in terms of the percentage of occurrence or the use of regional or general anaesthesia. Some studies do not even show differences in cardiovascular complications, the use of the intensive care unit, or the overall costs of hospitalization in elderly patients with the use of one or the other type of anaesthesia.^{39,40}

In a meta-analysis of 18 randomized studies on delirium and cognitive dysfunction, Bryson did not find significant differences between the use of regional or general anaesthesia, underlining the difficulty of preventing these two events because of the challenge of managing some of the patient-related risk factors.⁴¹

White, in an observational study of 65,535 patients, did not find any differences in mortality in patients taken to surgery for hip fracture repair under regional or general anaesthesia, and the only difference in outcome was increased mortality within the first 24 h associated with the use of cemented prostheses.⁴²

In analysing the physiological effects of regional anaesthesia, particularly in elderly patients, it may be that the effect is more severe because of depleted compensatory mechanisms as a result of chronic illnesses like hypertension, atherosclerosis, coronary heart disease, cardiac hypertrophy or dilatation, reduced ejection fraction, besides the physiological changes associated with age *per se*.^{7,43,44}

There is also the dose-effect relationship of medications in the central or peripheral nervous system as a result of the deterioration of myelin bridges and reduced amount of cerebrospinal fluid.⁴⁵

Intraoperative neurological monitoring devices

Devices for measuring the depth of anaesthesia have been developed over the past few years. Based on the measurement of electroencephalographic changes, they include the bispectral index monitor, the entropy module and the narcotrend monitors, which apply Fourier's theorem to transform the electroencephalographic signal into an interpretable numerical readout. Although studies are still needed to confirm some of the advantages attributed to these devices,⁴⁶ there is evidence already of the benefits of their use, particularly in patients with neurological or vital risk, as is the case with elderly patients.

From the anaesthetic perspective, these devices allow for a tighter control of the effects of anaesthesia on the brain, as they allow observation of signals that appear to be related with patient outcomes.

Sessler showed an increase in mortality and length of stay in patients with bispectral index values (BIS) under 45, low anaesthetic concentration, low mean alveolar concentration (MAC) and low arterial pressure.⁴⁷ In an earlier study, Monk

had already shown an increase in mortality one year following the surgery, finding a correlation between comorbidities, and low BIS and intraoperative blood pressure values.⁴⁸ The association between mortality and low BIS values could be secondary to an intrinsic patient sensitivity to the anaesthetic agents, where an "overdose" may occur without adequate dose measurement in the brain, increasing postoperative complications and short-term mortality.⁴⁹ BIS monitoring to avoid values under 40 for more than 5 min could be useful for patient management.⁵⁰

Using near-infrared spectroscopy (NIRS), the brain oximeter measures light-wave absorption and release at a specific frequency with a venous/arterial proportion of 70% and 30%, respectively, to determine the content of oxygenated haemoglobin.⁵¹

Oximetry values are relative as they depend on the patient's basal state, and have to be individualized for each case.

Cerebral oximetry helps recognize the basal level of brain oxygenation and identify changes in brain perfusion resulting from changes in arterial pressure and oxygenation during surgery, thus optimizing potential sources of neurological risk.⁵² Hoppenstein analyzed the changes in regional blood flow measured with cerebral oximetry during anaesthesia in patients with femoral neck fractures randomized to general or spinal anaesthesia, and found that desaturation percentages were higher in patients who received spinal anaesthesia compared to those who received general anaesthesia. The author concludes that the observation may depend on the patient's prior pathophysiological state and the ability to respond to changes in regional blood flow.⁵³ In a study in patients under general anaesthesia, Burkhardt observed a lower autoregulation capability in elderly patients compared to younger patients, using cerebral oximetry values as markers of autoregulation effectiveness.⁵⁴ In geriatric patients taken to surgery under general anaesthesia for hip fracture repair, Papadopoulos observed a relationship between brain desaturation and increased postoperative cognitive dysfunction.⁵⁵ Casati also found a relationship between cerebral desaturation and postoperative cognitive disorders, and also prolonged length of stay.⁵⁶ The effect of cerebral desaturation as a measurement of brain autoregulation may influence outcome, reflecting integrated information regarding oxygen supply and consumption in the brain; cerebral oximetry is a promising technology considering that it monitors essential and relevant physiology.⁵⁷

Conclusions

Elderly patients require a different approach based on the understanding of the physiological, anatomical, social and vital differences that occur with old age, in order to offer better therapeutic options that are sometimes far from science and closer to the form of vital support derived from listening and understanding.

The adequate treatment of elderly patients leads to the reduction of complications that usually result in higher overall costs for the health system and for society.

Programmes focused on elderly patient care, supported by social programmes that promote maintenance of a healthy

and functional life, should be set up at least in high complexity centres.

Conflict of interest

The author has worked as speaker and consultant for Medtronic.

Funding

The author's own resources.

REFERENCES

1. United Nations, Department of Economic and Social Affairs, Population Division, United Nations, New York 2010. World population ageing. New York: Population Division, United Nations; 2010.
2. López JH, López R. Sobre la muerte: a quien pueda interesar. *Rev Colomb Anestesiol.* 2012;40:195-8.
3. Ward SA, Parikh S, Workman B. Health perspectives: international epidemiology of ageing. *Best Pract Res Clin Anaesthesiol.* 2011;25:305-17.
4. Fuchs J, Scheidt-Nave C, Hinrichs T, Mergenthaler A, Stein J, Riedel-Heller SG, et al. Indicators for healthy ageing – a debate. *Int J Environ Res Public Health.* 2013;10:6630-44.
5. Zhuravleva T, Alperin B, Haring A, Rentz D. Age-related decline in bottom-up and selective attention in the very old. *J Clin Neurophysiol.* 2014;261-71.
6. Peden CJ, Grocott MPW. National research strategies: what outcomes are important in peri-operative elderly care? *Anaesthesia.* 2014;69:61-9.
7. Yang R, Wolfson M, Lewis MC. Unique aspects of the elderly surgical population: an anesthesiologist's perspective. *Geriatr Orthop Surg Rehabil.* 2011;56-64.
8. Silbert B, Evered L, Scott DA. Cognitive decline in the elderly: is anaesthesia implicated? *Best Pract Res Clin Anaesthesiol.* 2011;25:379-93.
9. Glance LG, Osler TM, Neuman MD. Redesigning surgical decision making for high-risk patients. *N Engl J Med.* 2014;370:1379-81.
10. Deiner S, Jeffrey H. Long-term outcomes in elderly surgical patients. *Mt Sinai J Med.* 2013;79:95-106.
11. Guenther U, Radtke FM. Delirium in the postanaesthesia period. *Curr Opin Anaesthesiol.* 2011;24:670-5.
12. McDaniel M, Brudney C. Postoperative delirium: etiology and management. *Curr Opin Crit Care.* 2012;18:372-6.
13. Alcover L, Badenes R, Montero MJ, Soro M, Belda FJ. Postoperative delirium and cognitive dysfunction. *Trends Anaesth Crit Care.* 2013;3:199-204.
14. Monk TG, Price CC. Postoperative cognitive disorders. *Curr Opin Crit Care.* 2011;17:376-81.
15. Grape S, Ravussin P, Rossi A, Kern C, Steiner LA. Postoperative cognitive dysfunction. *Trends Anaesth Crit Care.* 2012;2:98-103.
16. Moller J, Cluitmans P, Rasmussen L, Houw P, Rasmussen H, Canet J, et al. Long-term postoperative cognitive dysfunction in the elderly: ISPOCD1 study. *Lancet.* 1998;351:857-61.
17. Monk TG, Weldon BC, Garvan CW, Dede DE, Van der Aa MT, Heilman KM, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology.* 2008;108:18-30.
18. Long L, Shapiro W, Leung J. A brief review of practical preoperative cognitive screening tools. *Can J Anaesth.* 2012;59:798-804.
19. Gurlit S, Möllmann M. How to prevent perioperative delirium in the elderly? *Z Gerontol Geriatr.* 2008;41:447-52.
20. Kim S, Brooks AK. Preoperative assessment of the older surgical patient: honing in on geriatric syndromes. *Clin Interv Aging.* 2015;10:13-27.
21. Brown EN, Purdon PL. The aging brain and anesthesia. *Curr Opin Anaesthesiol.* 2013;26:414-9.
22. Fontana L. Modulating human aging and age-associated diseases. *Biochim Biophys Acta.* 2009;1790:1133-8.
23. Yau PL, Hempel R, Tarsi A, Convit A. Cerebral white matter and retinal arterial health in hypertension and type 2 diabetes mellitus. *Int J Hypertens.* 2013;3296-302.
24. Stone P, Doherty P. Anaesthesia for elderly patients. *Anaesth Intensive Care Med.* 2007;8:361-4.
25. Miyawaki CE. Association of social isolation and health across different racial and ethnic groups of older Americans. *Ageing Soc.* 2015;35:2201-28.
26. Afilalo J, Alexander KP, Mack MJ, Maurer MS, Green P, Allen LA, et al. Frailty assessment in the cardiovascular care of older adults. *J Am Coll Cardiol.* 2014;63:747-62.
27. Strøm C, Rasmussen LS, Sieber FE. Should general anaesthesia be avoided in the elderly? *Anaesthesia.* 2014;69:35-44.
28. Marcantonio ER, Rudolph JL, Culley D, Crosby G, Alsop D, Inouye SK. Serum biomarkers for delirium. *J Gerontol A Biol Sci Med Sci.* 2006;61:1281-6.
29. van den Boogaard M, Kox M, Quinn KL, van Achterberg T, van der Hoeven JG, Schoonhoven L, et al. Biomarkers associated with delirium in critically ill patients and their relation with long-term subjective cognitive dysfunction; indications for different pathways governing delirium in inflamed and noninflamed patients. *Crit Care.* 2012;15:1-9.
30. Leung JM, Tsai TL, Sands LP. Preoperative frailty in older surgical patients is associated with early postoperative delirium. *Anesth Analg.* 2011;112:1199-201.
31. Hubbard RE, Story DA. Patient frailty: the elephant in the operating room. *Anaesthesia.* 2014;69:26-34.
32. Partridge JSL, Harari D, Dhesi JK. Frailty in the older surgical patient: a review. *Age Ageing.* 2012;41:142-7.
33. Curcio C-L, Henao G-M, Gomez F. Frailty among rural elderly adults. *BMC Geriatr.* 2014;10:2.
34. McMillan GJ, Hubbard RE. Frailty in older inpatients: what physicians need to know. *QJM.* 2012;105:1059-65.
35. Jevtic-Todorovic V, Absalom AR, Blomgren K, Brambrink A, Crosby G, Culley DJ, et al. Anaesthetic neurotoxicity and neuroplasticity: an expert group report and statement based on the BJA Salzburg Seminar. *Br J Anaesth.* 2013;111:143-51.
36. Lin EP, Soriano SG, Loepke AW. Anesthetic neurotoxicity. *Anesthesiol Clin.* 2014;32:133-55.
37. Callaway JK, Jones NC, Royse AG, Royse CF. Memory impairment in rats after desflurane anesthesia is age and dose dependent. *J Alzheimer's Dis.* 2015;44:995-1005.
38. Whittington RA, Bretteville A, Dickler MF, Planell E. Anesthesia and tau pathology. *Prog Neuropsychopharmacol Biol Psychiatry.* 2013;47:147-55.
39. Slor CJ, De Jonghe JP, Vreeswijk R, Groot E, Ploeg TV, Van Gool WA, et al. Anesthesia and postoperative delirium in older adults undergoing hip surgery. *J Am Geriatr Soc.* 2011;59:1313-9.
40. Le-Wendling L, Bihorac A, Baslanti TO, Lucas S, Sadashivan K, Wendling A, et al. Regional anesthesia as compared to general anesthesia for surgery in geriatric patients with hip fracture: does it decrease morbidity, mortality and healthcare costs? Results of a single centered study. *Pain Med.* 2012;13:948-56.
41. Bryson GL, Wyand A. Evidence-based clinical update: general anesthesia and the risk of delirium and postoperative cognitive dysfunction. *Can J Anaesth.* 2006;53:669-77.

42. White SM, Moppett IK, Griffiths R. Outcome by mode of anaesthesia for hip fracture surgery. An observational audit of 65 535 patients in a national dataset. *Anaesthesia*. 2014;69:224–30.
43. Halaszynski TM. Pain management in the elderly and cognitively impaired patient: the role of regional anaesthesia and analgesia. *Curr Opin Anaesthesiol*. 2009;22:594–9.
44. Inés M, Valencia B. Envejecimiento de la población: un reto para la salud pública. *Rev Colomb Anestesiol*. 2012;40:192–4.
45. Veering B. Regional anaesthesia in the elderly. *Eur Soc Anaesthesiol*. 2011;18–21.
46. Ellerkmann RK, Soehle M, Kreuer S. Brain monitoring revisited: what is it all about? *Best Pract Res Clin Anaesthesiol*. 2013;27:225–33.
47. Sessler DI, Sigl JC, Kelley SD, Chamoun NG, Manberg PJ, Saager L, et al. Hospital stay and mortality are increased in patients having a “triple low” of low blood pressure, low bispectral index, and low minimum alveolar concentration of volatile anesthesia. *Anesthesiology*. 2012;116:1195–203.
48. Monk TG, Saini V, Weldon BC, Sigl JC. Anesthetic management and one-year mortality after noncardiac surgery. *Anesth Analg*. 2005;100:4–10.
49. Leslie K, Short TG. Low bispectral index values and death: the unresolved causality dilemma. *Anesth Analg*. 2011;113:660–3.
50. Leslie K, Myles PS, Forbes A, Chan MTV. The effect of bispectral index monitoring on long-term survival in the B-aware trial. *Anesth Analg*. 2010;110:816–22.
51. Murkin JM, Arango M. Near-infrared spectroscopy as an index of brain and tissue oxygenation. *Br J Anaesth*. 2009;103:i3–13.
52. Grocott HP. Avoid hypotension and hypoxia: an old anesthetic adage with renewed relevance from cerebral oximetry monitoring. *Can J Anaesth*. 2011;58:697–702.
53. Hoppenstein D, Zohar E, Ramaty E, Shabat S, Fredman B. The effects of general vs spinal anesthesia on frontal cerebral oxygen saturation in geriatric patients undergoing emergency surgical fixation of the neck of femur. *J Clin Anesth*. 2005;17:431–8.
54. Burkhart CS, Rossi A, Dell-Kuster S, Gamberini M, Möckli A, Siegemund M, et al. Effect of age on intraoperative cerebrovascular autoregulation and near-infrared spectroscopy-derived cerebral oxygenation. *Br J Anaesth*. 2011;107:742–8.
55. Papadopoulos G, Karanikolas M, Liarmakopoulou A, Papathanakos G, Korre M, Beris A. Cerebral oximetry and cognitive dysfunction in elderly patients undergoing surgery for hip fractures: a prospective observational study. *Open Orthop J*. 2012;6:400–5.
56. Casati A, Fanelli G, Pietropaoli P, Proietti R, Tufano R, Montanini S, et al. Monitoring cerebral oxygen saturation in elderly patients undergoing general abdominal surgery: a prospective cohort study. *Eur J Anaesthesiol*. 2007;24:59–65.
57. Meng L, Gelb W. Oximetría cerebral: tres preguntas esenciales. *Rev Colomb Anestesiol*. 2015;3:52–6.