



### Case report

# Anesthesia for bariatric surgery in a patient with Prader-Willi syndrome: Case report<sup>☆</sup>



Camilo Rada-Ortega<sup>a,\*</sup>, Carlos Fernando Gómez-Ramírez<sup>b</sup>

<sup>a</sup> MD, Postgraduate student in Anesthesiology, Universidad CES, Medellín, Colombia

<sup>b</sup> MD, Anesthesiologist, Hospital Manuel Uribe Ángel, Envigado, Colombia

#### ARTICLE INFO

##### Article history:

Received 5 March 2014

Accepted 26 February 2016

Available online 6 April 2016

##### Keywords:

Anesthesia

Obesity

Bariatric surgery

Sleep apnea, Obstructive

Prader-Willi syndrome

#### ABSTRACT

Prader-Willi syndrome is a genetic disorder characterized by hypotonia, obesity, short stature, mental retardation, hyperphagia, hypogonadism and low life expectancy.

We describe the case of a 31-year-old female patient with Prader-Willi syndrome scheduled for bariatric surgery. Anesthetic considerations are reviewed highlighting perioperative complications associated with this syndrome.

© 2016 Published by Elsevier España, S.L.U. on behalf of Sociedad Colombiana de Anestesiología y Reanimación.

### Anestesia para cirugía bariátrica en paciente con síndrome de Prader-Willy: reporte de un caso

#### RESUMEN

El síndrome de Prader-Willi es un desorden genético caracterizado por hipotonía, obesidad, baja estatura, retraso mental, hiperfagia, hipogonadismo y expectativa de vida reducida.

Describimos el caso de una paciente de 31 años con antecedente de síndrome de Prader-Willi, programada para realización de cirugía bariátrica. Se revisan las consideraciones anestésicas, haciendo énfasis en las complicaciones perioperatorias secundarias a este síndrome.

© 2016 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Colombiana de Anestesiología y Reanimación.

##### Palabras clave:

Anestesia

Obesidad

Cirugía bariátrica

Apnea del sueño obstructiva

Síndrome de Prader Willi

<sup>☆</sup> Please cite this article as: Rada-Ortega C, Gómez-Ramírez CF. Anestesia para cirugía bariátrica en paciente con síndrome de Prader-Willy: reporte de un caso. Rev Colomb Anestesiolog. 2016;44:255-258.

\* Corresponding author at: Carrera 40 A No. 11 B-54, Medellín, Colombia.

E-mail address: [drcroe@gmail.com](mailto:drcroe@gmail.com) (C. Rada-Ortega).

## Introduction

Prader-Willi syndrome is a genetic disorder characterized by hypotonia, obesity, short stature, mental retardation, hyperphagia, hypogonadism and low life expectancy.<sup>1</sup>

It was described initially by JL Down in 1887 in a patient diagnosed with "polysarcia". In 1956, Prader, Labhart and Willi described nine other cases and gave the syndrome its name. In 1980, Ledbetter discovered the existence of a microdeletion of the 15q11-q13 region and three years later, Butler and Nichols observed this genomic imprinting in patients with Prader-Willi syndrome.<sup>2</sup>

There are few reported cases in the literature on the anesthetic management of adult patients with Prader-Willi syndrome with low life expectancy as a consequence of morbid obesity.<sup>3-5</sup> Furthermore, these patients are predisposed to sudden death from respiratory diseases or in the postoperative period.<sup>6-10</sup>

We present the anesthetic management of a 31-year-old patient diagnosed with Prader-Willi syndrome who was scheduled to undergo bariatric surgery.

## Case description

The patient was admitted to surgery at the Manuel Uribe Ángel Hospital in Envigado, Antioquia, Colombia. The patient was 31 years old and female with a background of Prader-Willi syndrome associated with hypertension, extreme obesity (BMI 54), mental retardation, hypothyroidism and obstructive sleep apnea syndrome. She was admitted to undergo a gastric bypass.

The patient is considered to be a NYHA functional class III, ASA 4.

The physical exam revealed central cyanosis and blood pressure of 140/80 mmHg, heart rate of 68 bpm, respiratory rate of 19 breaths/min, Mallampati score of II, oral aperture above 4 cm, cervical perimeter below 40 cm, 35° extension of the upper neck and thyromental distance above 6 cm.

Rhythmic cardiac sounds were auscultated, revealing P2 > A2 along with a protomesosystolic murmur at the pulmonary and tricuspid focuses, III/VI. There was a diminished bilateral vesicular murmur.

Oxygen saturation oscillated between 80 and 90% through the nasal canal at 3 l/min.

The patient did not resist any of the examiner's maneuvers and collaborated during the whole evaluation.

Presurgical exams revealed: hemoglobin 12.2 g/dl, hematocrite 41%, 371 000 platelets/mL, TSH 7.92 mIU/L, HA1G 6.75%, arterial blood gases (FiO<sub>2</sub>) 0.21: pH 7.38, PCO<sub>2</sub> 54 mmHg, PO<sub>2</sub> 25 mmHg, HCO<sub>3</sub> 29 mEq/l. Echocardiogram (TTE): FE: 60%, PSAP 52 mmHg. There were slightly insufficient tricuspid and pulmonary valves, but normal cavity size.

In the operating room, a 16G cannula was placed on 2 cephalic veins; a rapid sequence induction was performed with previous oxygenation during 5 min until a maximum saturation of 95% was reached, followed by a 100 mcg bolus intravenous administration of remifentanyl, 60 mg lidocaine, 150 mg propofol, 100 mg succinylcholine, laryngoscopy with

valve curve #3 (Cormack II visualization), introduction of 8.0 mm orotracheal tube, maintenance with 3% sevoflurane, flow of oxygen gases at 0.3 l/min and air at 0.5 l/min, and finally an intravenous infusion of remifentanyl at 0.1 mcg/kg/min.

The anesthesia machine was programmed for volume, with a tidal volume of 7 ml/kg, PEEP 5 cm H<sub>2</sub>O, showing peak pressures higher than 35 cm H<sub>2</sub>O with plateau pressures lower than 30 cm H<sub>2</sub>O. The bilateral position of the orotracheal tube was immediately verified through auscultation to rule out obstruction or kinking. The monitor showed a progressive increase in the level of CO<sub>2</sub> at the end of the expiration without presenting inclination in the curve of the capnogram in phase 2. 6 mg cisatracurium was administered intravenously, with significant improvement in the respective values.

During the intraoperative period the surgical team identified severe cardiomegaly associated with hepatomegaly, which led to the decision to perform gastric sleeve surgery to then proceed with a bypass in the second surgical period.

At the end of the procedure, 0.6 mg of hydromorphone was administered intravenously.

There were no episodes of desaturation, severe hypotension/hypertension or arrhythmias. She was extubated after finishing surgery and transferred to the special care unit and supplemented with 50% oxygen at 10 l/min with the Ventury system. No reintubation was required.

## Discussion

Prader-Willi syndrome is considered to be the leading cause of obesity associated with genetic syndromes and has a prevalence of approximately 1/25000.<sup>11</sup> The partial deletion of the long arm of the paternal chromosome 15 is a common marker of this disease,<sup>12</sup> though it can also present as a maternal uniparental disomy.<sup>13</sup>

The annual mortality rate was 3% for all ages, but the rate increases to 7% over the age of 30.<sup>11</sup> Patients generally suffer an early death due to secondary complications related to obesity, such as diabetes mellitus, hypertension, obstructive apnea-hypopnea, cardiovascular disease and respiratory failure.<sup>14</sup>

The clinical course for Prader-Willi syndrome is usually divided in two phases.<sup>15</sup> The first phase occurs during the neonatal and lactation periods and is characterized by marked hypotonia, difficulties with suction, persistent cough, crying and episodes of asphyxia.<sup>16,17</sup> The second phase, between the ages of 2-5, is characterized by hypogonadism, mental retardation and obesity related to hyperphagia, most likely due to a hypothalamic defect in the satiety center<sup>18</sup> or an innate failure to metabolize lipids and carbohydrates.<sup>19</sup>

As was previously mentioned, there are few available reports on the anesthetic management of patients with Prader-Willi syndrome, and in the case at hand the patient who was scheduled for a low-risk cardiovascular procedure was in her fourth decade of life with extreme obesity, obstructive sleep apnea syndrome and chronic hypoxia.

Alterations in body temperature (hyperthermia and hypothermia), intraoperative arrhythmias, cor pulmonale and clinical consequences of obesity (decrease in functional

residual capacity) are particular problems during the perioperative period.<sup>20-24</sup>

During the intraoperative period the only difficulty with our patient was the elevated peak pressure values accompanied by normal plateau pressure levels. An elevated peak pressure can be accompanied by a concomitant elevation of plateau pressure, but when this association is not present, we are faced with an increase in the resistance of the airways (decrease of dynamic pulmonary distensibility), causing us to rule out bronchospasms, presence of secretions, obstruction, kinking or biting of the orotracheal tube.<sup>25</sup>

It should be noted that excessive intra-abdominal pressure decreases the pulmonary volumes in morbidly obese patients under sedation or anesthesia, notably in terms of residual functional capacity, while the alveolo-arterial oxygenation gradient is increased along with respiratory resistance.<sup>26</sup> Our patient's condition improved with the administration of a neuromuscular blocker. This technique is similar to that performed in patients with intra-abdominal hypertension syndrome, in which neuromuscular blockers are used to lower intra-abdominal and airway pressure.<sup>27</sup>

There was also no difficulty in performing the laryngoscopy, despite reports of difficult airways in these patients.<sup>28</sup>

A transfer to the special care unit was considered since these patients have an increased risk of postoperative hypoxia attributed to the altered response in consciousness level due to changes in oxygen and CO<sub>2</sub> in blood pressure.<sup>29</sup>

Another issue is the documented presence of an oppositional and aggressive personality type associated with this syndrome,<sup>30</sup> although in our case the patient cooperated at all times during her hospital stay and did not require sedatives such as benzodiazepine which can possibly cause episodes of desaturation and apnea.

In conclusion, the approach that an anesthesiologist takes with a Prader-Willi patient should include all the associated comorbidities, especially secondary ones like morbid obesity, taking into account the appropriate postoperative monitoring to avoid the onset of respiratory complications.

## Funding

Authors' own resources.

## Conflict of interest

None declared.

## REFERENCES

- Holm VA, Cassidy SB, Butler MG, Hanchett JM, Greenswag LR, Whitman BY, et al. Prader-Willi syndrome: consensus diagnostic criteria. *Pediatrics*. 1993;91:398-402.
- Zapico M. Aspectos endocrinológicos del síndrome de Prader-Willi. *Rev Med Chil*. 2005;132:427-31.
- Cho EC, Jee SE, Jang Y, Park SS, Kim JT, Song HK. Prader-Willi syndrome: a case report. *Korean J Anesthesiol*. 1999;36:1091-4.
- Cavaliere F, Cormaci S, Cormaci M, Alberti A, Colabucci F. General anesthesia in Prader-Willi syndrome. *Minerva Anesthesiol*. 1996;62:327-32.
- Sloan TB, Kaye CI. Rumination risk of aspiration of gastric contents in the Prader-Willi syndrome. *Anesth Analg*. 1991;73:492-5.
- Stevenson DA, Anaya TM, Clayton-Smith J, Hall BD, Van Allen MI, Zori RT, et al. Unexpected death and critical illness in Prader-Willi syndrome: report of ten individuals. *Am J Med Genet A*. 2004;124:158-64.
- Nagai T, Obata K, Tonoki H, Temma S, Murakami N, Katada Y, et al. Cause of sudden, unexpected death of Prader-Willi syndrome patients with or without growth hormone treatment. *Am J Med Genet A*. 2005;136:45-8.
- Nordmann Y, Eiholzer U, l'Allemand D, Mirjanic S, Markwalder C. Sudden death of an infant with Prader-Willi syndrome - not a unique case? *Biol Neonate*. 2002;82:139-41.
- Schrander-Stumpel CT, Curfs LM, Sastrowijoto P, Cassidy SB, Schrander JJ, Fryns JP. Prader-Willi syndrome: causes of death in an international series of 27 cases. *Am J Med Genet A*. 2004;124:333-8.
- Oiglane E, Ounap K, Bartsch O, Rein R, Talvik T. Sudden death of a girl with Prader-Willi syndrome. *Genet Couns*. 2002;13:459-64.
- Whittington JE, Holland AJ, Webb T, Butler J, Clarke D, Boer H. Population prevalence and estimated birth incidence and mortality rate for people with Prader-Willi syndrome in one UK Health Region. *J Med Genet*. 2001;38:792-8.
- Ledbetter DH, Riccardi VM, Airhart SD, Strobel RJ, Keenan BS, Crawford JD. Deletions of chromosome 15 as a cause of the Prader-Willi syndrome. *N Engl J Med*. 1981;304:325-30.
- Mascari MJ, Gottlieb W, Rogan PK, Butler M, Waller D, Armour J, et al. The frequency of uniparental disomy in Prader-Willi syndrome. *N Engl J Med*. 1992;326:1599-607.
- Cassidy SB, Driscoll DJ. Prader-Willi syndrome. *Eur J Hum Genet*. 2009;17:3-13.
- Zellweger H, Schneider HJ. Syndrome of hypotonia hypopontia hypogonadism obesity (HHHO) or Prader-Willi syndrome. *Am J Dis Child*. 1968;115:588-98.
- Cohen MM Jr, Gorlin RJ. The Prader-Willi syndrome. *Am J Dis Child*. 1969;117:213-8.
- Landwirth J, Schwartz AH, Grunt JA. Prader-Willi syndrome. *Am J Dis Child*. 1968;116:211-7.
- Uehling D. Cryptorchidism in the Prader-Willi syndrome. *J Urol*. 1980;124:103-4.
- Palmer SK, Atlee JL. Anesthetic management of the Prader-Willi syndrome. *Anesthesiology*. 1976;44:61-3.
- Yamashita M, Koishi K, Yamaya R, Tsubo T, Matsuki A, Oyama T. Anaesthetic considerations in the Prader-Willi syndrome: report of four cases. *Can Anaesth Soc J*. 1983;30:179-84.
- Mayhew JF, Taylor B. Anaesthetic considerations in the Prader-Willi syndrome. *Can Anaesth Soc J*. 1983;30:565-6.
- Milliken RA, Weintraub DM. Cardiac abnormalities during anesthesia in a child with Prader-Willi syndrome. *Anesthesiology*. 1975;43:590-2.
- Yukioka H, Kitamura E, Nagata N, Fujimori M. Prader-Willi syndrome and anesthetic management. *Jpn J Anesth*. 1979;28:518-21.
- Noguchi I, Ebihara M, Yamaya R, Tsubo T, Matsuki A, Oyama T. Experience of anesthesia for a patient with Prader-Willi syndrome. *J Jpn Dent Soc Anesth*. 1980;8:56-62.
- Cover T, Tung Niu N. Differential diagnosis of high peak airway pressures. *Dimens Crit Care Nurs*. 2015;34:19-23.
- Pelosi P, Croci M, Ravagnan I, Cerisara M, Vicardi P, Lissoni A, et al. Respiratory system mechanics in sedated, paralyzed, morbidly obese patients. *J Appl Physiol*. 1997;82:811-8.

27. Rivera R, Rivera JS. Bloqueantes neuromusculares: en pro del uso adecuado. *Rev Colomb Anesthesiol.* 2011;39:352-7.
28. Lirk P, Keller C, Colvin J, Rieder J, Wulf K. Anaesthetic management of the Prader-Willi syndrome. *Eur J Anaesthesiol.* 2004;21:831-3.
29. Nixon GM, Brouillette RT. Sleep and breathing in Prader-Willi syndrome. *Pediatr Pulmonol.* 2002;34:209-17.
30. Rittinger O. Clinical aspects and genetics of Prader-Willi syndrome. *Klin Padiatr.* 2001;213:91-8.