



# Revista Colombiana de Anestesiología

Colombian Journal of Anesthesiology

[www.revcolanest.com.co](http://www.revcolanest.com.co)



## Essay

# Monitoring of national surgical care indicators in the Peruvian health system<sup>☆</sup>



Carlos Javier Shiraishi-Zapata <sup>a,b,\*</sup>

<sup>a</sup> Anaesthesia Service, Hospital ESSALUD Talara, Piura, Peru

<sup>b</sup> "Teléforo León Velasco" Anaesthesia Service, Hospital ESSALUD José Cayetano Heredia, Piura, Peru

### ARTICLE INFO

#### Article history:

Received 25 January 2017

Accepted 3 April 2017

Available online 4 July 2017

#### Keywords:

Operating Rooms

Mortality

Surgeons

Anesthesiologists

Health Surveillance

### ABSTRACT

Surgical care is an integral component of healthcare services in all countries, and its quality is evaluated through monitoring of national indicators of safe surgical and anaesthesia care. In Peru, monitoring is only partial because of the lack of data regarding some of the indicators. However, there is a need for comprehensive monitoring in order to gain knowledge of the progress towards the healthcare goals proposed by the World Health Organisation and The Lancet Commission Global Surgery.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Monitorización de indicadores nacionales de atención quirúrgica en el sistema sanitario peruano

### RESUMEN

La atención quirúrgica es un componente integral de los sistemas sanitarios en todos los países, y la valoración de su calidad se realiza a través de la monitorización de indicadores nacionales de atención segura quirúrgica y anestésica. En Perú, esta monitorización se realiza de forma parcial, pues no hay información respecto a algunos indicadores. Sin embargo, debería ser integral para conocer el progreso de los objetivos sanitarios propuestos por la Organización Mundial de la Salud y la Comisión Lancet en Cirugía Global.

© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

#### Palabras clave:

Quirófanos

Mortalidad

Cirujanos

Anestesiólogos

Vigilancia Sanitaria

\* Please cite this article as: Shiraishi-Zapata CJ. Monitorización de indicadores nacionales de atención quirúrgica en el sistema sanitario peruano. Rev Colomb Anestesiol. 2017;45:210-215.

\* Correspondence to: Servicio de Centro Quirúrgico y Anestesiología Hospital ESSALUD Talara, Avenida Panamericana s/n Talara, Piura, Peru.

E-mail address: [cshiraza@alumnes.ub.edu](mailto:cshiraza@alumnes.ub.edu)

2256-2087/© 2017 Sociedad Colombiana de Anestesiología y Reanimación. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Surveillance in public health has used standardised measurements designed to quantify the cost of disease in a population, monitor mortality rates, and provide guidance to health systems. Maternal mortality (MM), infant mortality (IM) and life expectancy (LE) rates have been important indicators for measuring the effectiveness of healthcare services over a considerable time period. However, as a result of the increase in LE in the world, as well as the drop in MM and IM rates, other measurements such as surgical care have gained importance when it comes to evaluating the health system.<sup>1</sup>

Surgical services are integral components of the public health system because they are part of the continuous primary care process. Although they have been recognised as a costly component, surgeries are performed in both affluent and precarious economic contexts alike.<sup>2,3</sup> In 2012, 312.9 million surgical procedures were carried out in the world<sup>4</sup>; however, information regarding frequency and safety of this care was severely limited by the lack of national data, considering that 70% of countries had no information on their national surgery volumes, and practically none attempted to value the allocation of surgical resources or the results.<sup>1</sup>

Peru was not an exception, because up until recently it did not monitor the main indicators on preparation, provision and effect of safe and affordable surgical and anaesthesia care for the population.<sup>1,5</sup> Some of the characteristics of the Peruvian health system are described below, followed by current considerations regarding monitoring of indicators and, finally, proposed measurements for comprehensive monitoring.

## The Peruvian health system

A mixed system, it comprises two large sub-sectors, public and private, with different assets and sources of funding coming from tax collection (Ministry of Health – MINSA – with its offering of Comprehensive Health Insurance – SIS), social security contributions through taxation on active worker salaries (Social Health Insurance – EsSalud), and premium payments for private insurance. This results in a fragmented system with several weaknesses such as network overlap, inability to provide comprehensive care, and absence of complementary services and continuity of care.<sup>6,7</sup>

## Timely access to essential surgery

Timely surgical care is the ability to receive rapid and appropriate care from a healthcare institution (primary care level hospital) within a two-hour period.<sup>5,8</sup> A segment of the Peruvian population lives at high altitude in the Andean region where access is difficult, but travel time to the closest healthcare institution (IPRESS) went down

from 39 to 42 min (walking), and from 27 to 23 in motor vehicles, between 2011 and 2014.<sup>9</sup> However, these time periods were estimated for the total number of healthcare institutions, including many facilities without surgical care capability.

## National surgical volume

According to the National Health Superintendency (SUSALUD), 613,396 surgical procedures (major and minor)<sup>10</sup> were performed in 31,151,643 inhabitants, for a rate of 1969 surgeries for every 100,000 inhabitants in 2015. The Lancet Commission on Global Surgery (LCGS) proposed that, by 2030, countries should be performing at least 5000 major procedures for every 100,000 inhabitants per year, given that it is the volume that correlates with several desirable health results such as a LE of 74–75 years, a MM rate of less than 100 women for every 100,000 live births, and the 143 million additional surgeries required worldwide in order to save lives and prevent disability. Moreover, this figure could be used for benchmarking surgical services and as a target for monitoring the execution of surgical care plans at a national level.<sup>5,11,12</sup>

## Number of operating rooms

The total number of Peruvian healthcare institutions (IPRESS) that provide surgical care are already registered in an electronic database called the National Registry of Healthcare Institutions (RENIPRESS) belonging to SUSALUD, although there is no information on the number of operating rooms that are operational in each surgical service. This means that there are no data regarding the volume of surgical patients per operating room per year.<sup>13</sup> Only ESSALUD reported running 250 operating rooms at a national level, covering 10,754,665 affiliates in 2015.<sup>14,15</sup>

## Density of surgical specialists

In 2014, there were 1382 specialists accredited as anaesthetists and 8804 accredited as surgeons (all surgical specialties).<sup>16</sup> However, in the Medical College of Peru (MCP) database, licensure figures for 2016 were 1706 anaesthetists, 6765 surgeons and 3055 obstetricians and gynaecologists,<sup>17</sup> accounting for a rate of 36.6 specialists for every 100,000 inhabitants in a total population of 31,488,625 inhabitants.<sup>18</sup> But this rate might be higher considering that many physicians graduating from residency programmes practice in different specialties without having yet obtained their professional degree, which is a pre-requisite for inclusion in the national MCP database of specialists.

There is a correlation between this density figure and the MM rate given that, for every 10 increment units there is a 13.1% (95% CI 11.3–14.8) reduction in the MM rate.<sup>19</sup> For this reason, the LCGS has emphasised the importance of achieving

**Table 1 – Monitoring of national safe surgery indicators in three countries with different THEPC.**

Country	THEPC (in \$int)	Classification by THEPC	Measurement period	Surgical volume	Rate of procedures per operating room	Surgical mortality rate on post-operative day 1	30-day post-operative mortality rate	Benefits of monitoring
El Salvador	\$565	Intermediate	2009–2010	172,972 per year	1,197	0.42%	1.58%	<ul style="list-style-type: none"> <li>- Provided an estimate of the surgical capacity and mortality in the country.</li> <li>- Contributed with a starting point for understanding how surgical services are provided to the national population.</li> </ul>
United Kingdom	\$3,377	High	2005–2009	5.4 millions (period)/1,156,443 (2009)	1,571	0.0011%	0.012%	<ul style="list-style-type: none"> <li>- Provided an important information summary about surgical performance, allowing for a statistical control of the process.</li> <li>- Assessed the service performances in order to help monitor demand and benchmark standards, and to guide service rationalisation and identify gaps in resource distribution, safety and quality.</li> </ul>
Liberia	\$98	Very low	2009–2010	7,654 (2010)–330.1 <sup>a</sup>	589 (51–1,464)	145 <sup>a,b</sup>	0.1% 1.4%	<ul style="list-style-type: none"> <li>- Recognised a significant surgical volume and low density and quality of infrastructure and adequately trained staff.</li> <li>- Provided information on the crisis or surgical and anaesthetic services in Africa.</li> <li>- Distances covered by patients in order to get to the hospitals was 30 km.</li> <li>- Revealed that surgical mortality rates were still a problem and that improvements were required in order to improve post-operative survival.</li> </ul>

Source: Adapted from Molina, et al.<sup>31</sup>, Agha<sup>32</sup>, Knowlton, et al.<sup>33</sup>, Organización Mundial de la Salud<sup>34</sup> y Weiser, et al.<sup>35</sup>.

<sup>a</sup> Rates per 100,000 habitantes.

<sup>b</sup> Intra-operative mortality rate.

a minimum density of 20 specialists for every 100,000 inhabitants by 2030.<sup>20</sup>

### **Protection against impoverishment and catastrophic healthcare expenditures**

The percentage of the Peruvian population with access to coverage, as well as the total health expenditure per capita (THEPC) increased steadily during the time period between 2004 and 2014 (from 37.3 to 69%, and from int. \$282.22 to int. \$656.18, respectively).<sup>10,21,22</sup> However, a segment of the population (27% or nearly 8 million, including more than 1.7 million in a situation of poverty) has no coverage whatsoever and, consequently, no way to access healthcare services. This creates a risk of impoverishment and catastrophic spending associated with surgical care. The Peruvian population exposed to the risk of such expenditures is 36.1% and 29%, respectively, according to a research based on stochastic and surgical cost models.<sup>23,24</sup>

This indicator is subject to continuous monitoring by the National Statistics and Information Technology Institute (INEI) as part of the sustainable development goals, as a way to evaluate the goal on universal health coverage, including protection against financial risks.<sup>25</sup>

### **Problems associated with absent indicators**

Absent indicators include timely access to essential surgery, number of operating rooms, and mortality rates during hospitalisation (with reasonable correlation with 30-day mortality),<sup>5,26</sup> and they preclude comprehensive monitoring and knowledge of the magnitude of the work performed in surgery and anaesthesia. Likewise, absent indicators do not allow to quantify the impact of the national regulations on safe and standardised anaesthesia care (including the World Health Organisation surgical safety checklist).<sup>27,28</sup> Consequently, it is not possible to address deficiencies in peri-operative outcomes<sup>29</sup> or guide national surgical care policies and programmes.<sup>30</sup> In contrast, Table 1 summarises the benefits derived from this form of monitoring in three countries with different THEPC and surgical volumes (El Salvador, United Kingdom and Liberia).<sup>31-35</sup>

The WHO has emphasised that national monitoring of these indicators must be implemented urgently in all member countries because it will enhance the understanding of unmet needs, improve overall surgical and anaesthetic capacity in the context of universal health coverage, and will help implement reliable steps regarding access to essential surgery and anaesthesia.<sup>3</sup>

### **Proposals for achieving comprehensive indicator monitoring**

Because of its fragmentation, the Peruvian health system will require a number of actions to unify information coming from the different sub-sectors in order to consolidate

single national indicators. Proposed measures include the following.

First of all, strengthening tools for automated collection of standardised indicator data. Several are already available in various health sub-sectors, but SUNASA has a suitable software for entering standardised data through RENIPRESS,<sup>13</sup> which displays open information regarding infrastructure, equipment, human resources and organisation (including availability of surgical services) of healthcare institutions (IPRESS) of all the sub-sectors. It also supports IPRESS registration, categorisation, information updates and withdrawals. Consequently, it could consolidate data regarding indicators pertaining to such things as active operating rooms and post-operative mortality in each Peruvian healthcare institution.<sup>36</sup>

Second, collection of health information by means of population-based mechanisms (census). The INEI has scheduled a geo-referenced population census in 2017 using a mapping process of the total number of Peruvian households. This census will allow to obtain updated information of population segments with health coverage.<sup>37,38</sup> It could also estimate the time required to access a healthcare institution with surgical services by means of geospatial mapping, as has been done indirectly in other countries through multiple interviews with hospital representatives.<sup>8</sup>

Equally important is continuous coordination between MCP, scientific societies of surgical specialties, the National Council (CONAREME) and medical residency programmes in order to acquire accurate consensus data regarding the existing number of surgical specialists in Peru. These open data will serve to compare regional realities within the national context, identifying regions with a deficit of specialists in order to inform plans for corrective action.

Moreover, it would be desirable to develop a national plan for safe surgery in order to coordinate policies in accordance with the national and supranational regulations regarding staffing, infrastructure, provision of care, financing and use of the information.<sup>5</sup>

Finally, a national observatory consisting of representatives of the health sub-sectors would provide continuous monitoring of surgical and anaesthesia care indicators.

### **Funding**

The author did not receive sponsorship to carry out this article.

### **Conflict of interest**

The author declares having no conflicts of interest.

### **REFERENCES**

- Weiser TG, Makary MA, Haynes AB, Dziekan G, Berry WR, Gawande AA. Safe Surgery Saves Lives Measurement and Study Groups. Standardised metrics for global surgical surveillance. Lancet. 2009;374:1113-7.

2. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet.* 2008;372:139-44.
3. World Health Organization. Sixty-Eighth World Health Assembly. Resolutions, decisions and annexes. WHA68.15: strengthening emergency and essential surgical care and anaesthesia as a component of universal health coverage. Ginebra [Internet]; 2015 [Cited 2016 May 22] Available from: [http://apps.who.int/gb/ebwha/pdf\\_files/WHA68-REC1/A68\\_R1.REC1-en.pdf#page=27](http://apps.who.int/gb/ebwha/pdf_files/WHA68-REC1/A68_R1.REC1-en.pdf#page=27)
4. Weiser TG, Haynes AB, Molina G, Lipsitz SR, Esquivel MM, Uribe-Leitz T, et al. Estimate of the global volume of surgery in 2012: an assessment supporting improved health outcomes. *Lancet.* 2015;385 Suppl. 2:S11.
5. Meara JG, Leather AJ, Hagander L, Alkire BC, Alonso N, Ameh EA, et al. Global surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *Lancet.* 2015;386:569-624.
6. Ministerio de Salud. Dirección General de Epidemiología. Análisis de la Situación de Salud en Perú; 2013 [Cited 2017 Mar 01] Available from: <http://www.dge.gob.pe/portal/docs/intsan/asis2012.pdf>
7. Alcalde-Rabanal JE, Lazo-González O, Nigenda G. Sistema de salud de Perú. *Salud Publica Mex.* 2011;53 Suppl. 2: S243-54.
8. Raykar NP, Bowder AN, Liu C, Vega M, Kim JH, Boye G, et al. Geospatial mapping to estimate timely access to surgical care in nine low-income and middle-income countries. *Lancet.* 2015;385 Suppl. 2:S16.
9. INEI. Compendio Estadístico del Perú 2015 [Internet]; 2015 [Cited 2016 Aug 14] Available from: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Lib1253/compendio2015.html](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1253/compendio2015.html)
10. SUSALUD. Datos abiertos: Producción Asistencial en Intervenciones Quirúrgicas de las IPRESS [Internet]. 2017 [Cited 2017 Apr 10] Available from: <http://datos.susalud.gob.pe/dataset/consulta-h-produccion-asistencial-en-intervenciones-quirurgicas-de-las-ipress>
11. Uribe-Leitz T, Esquivel MM, Molina G, Lipsitz SR, Verguet S, Rose J, et al. Projections to achieve minimum surgical rate threshold: an observational study. *Lancet.* 2015;385 Suppl. 2:S14.
12. Esquivel MM, Molina G, Uribe-Leitz T, Lipsitz SR, Rose J, Bickler SW, et al. Proposed minimun rates of surgery to support desirable health outcomes: an observational study based on four strategies. *Lancet.* 2015;385 Suppl. 2: S12.
13. SUSALUD. RENIPRESS [Internet]. 2017 [Cited 2016 Nov 29] Available from: <http://datos.susalud.gob.pe/dataset/registro-de-ipress-renipress>
14. ESSALUD. Oficina de Relaciones Institucionales. Estadística Institucional: Memoria Anual 2015 de ESSALUD [Internet]. 2017 [Cited 2016 Nov 03] Available from: <http://www.essalud.gob.pe/memoria-institucional/>
15. ESSALUD. Oficina de Relaciones Institucionales. Estadística Institucional: ESSALUD: Principales Indicadores de Salud, 1990-2015 [Internet]. 2017 [Cited 2016 Nov 3]. Available from: [http://www.essalud.gob.pe/downloads/series\\_estadisticas\\_1990\\_2015\\_VF.pdf](http://www.essalud.gob.pe/downloads/series_estadisticas_1990_2015_VF.pdf)
16. Holmer H, Lantz A, Kunjumen T, Finlayson S, Hoyler M, Siyam A, et al. Global distribution of surgeons, anaesthesiologists, and obstetricians. *Lancet Glob Health.* 2015; Suppl. 2: S9-11.
17. Colegio Médico del Perú. Conoce a tu médico [Internet]. 2017 [Cited 2016 Sep 09] Available from: <http://cmp.org.pe/servicios/conoce-a-tu-medico/>
18. INEI. Índice Temático: Población [Internet]. 2017 [Cited 2016 Sep 10] Available from: <https://www.inei.gob.pe/estadisticas/indice-tematico/population/>
19. Holmer H, Shrime MG, Riesel JN, Meara JG, Hagander L. Towards closing the gap of the global surgeon, anaesthesiologist, and obstetrician workforce: thresholds and projections towards 2030. *Lancet.* 2015;385 Suppl. 2:S40.
20. Daniels KM, Riesel JN, Meara JG. The scale-up of the surgical workforce. *Lancet.* 2015;385 Suppl. 2:S41.
21. World Health Organization. Global health expenditure database. NHA indicators. 2017 [Cited 2016 Sep 05] Available from: <http://apps.who.int/nha/database/ViewData/Indicators/en>
22. Ministerio de Salud. Avances de la Reforma de Salud. Reporte Informativo 2013-2014. Oficina General de Comunicaciones; 2015 [Cited 2016 Nov 05] Available from: <http://bvs.minsa.gob.pe/local/minsa/3240.pdf>
23. Ministerio de Salud. Boletín N°6: Reforma es más salud; 2015 [Cited 2016 Nov 08] Available from: <http://bvs.minsa.gob.pe/local/MINSA/3440.pdf>
24. Shrime MG, Dare A, Alkire BC, Meara JG. A global country-level comparison of the financial burden of surgery. *Br J Surg.* 2016;103:1453-61.
25. INEI. Perú: Sistema de Monitoreo y Seguimiento de los Indicadores de Objetivos de Desarrollo Sostenible. 2017 [Cited 2016 Mar 07] Available from: <http://ods.inei.gob.pe/ods/inicio.html>
26. Ariyaratnam R, Palmqvist CL, Hider P, Laing GL, Stupart D, Wilson L, et al. Towards a standard approach to measurement and reporting of peri-operative mortality rate as a global indicator of surgery. *Surgery.* 2015;158:17-26.
27. Ministerio de Salud. Oficina General de Comunicaciones. Normas Legales. RM 486-2005. NT N° 030-MINSA/DGSP-V.01: "Norma Técnica de los Servicios de Anestesiología"; 2005, 2017 [Cited 2017 Mar 09] Available from: <http://www.dgiem.gob.pe/norma-tecnica-de-los-servicios-de-anestesiologia/>
28. Ministerio de Salud. Oficina General de Comunicaciones. Normas Legales. NT N° 089-MINSA/DGSP-V.01: "Norma Técnica de Salud para la Atención Anestesiológica"; 2011 [Cited 2017 Mar 09] Available from: [ftp://ftp2.minsa.gob.pe/normaslegales/2011/RM022-2011-MINSA\(A\).PDF](ftp://ftp2.minsa.gob.pe/normaslegales/2011/RM022-2011-MINSA(A).PDF)
29. Weiser TG, Makasa EM, Gelb AW. Improving perioperative outcomes in low-resource countries: it can't be fixed without data. *Can J Anesth.* 2015;62:1239-43.
30. Steffner KR, McQueen KA, Gelb AW. Patient safety challenges in low-income and middle-income countries. *Curr Opin Anaesthesiol.* 2014;27:623-9.
31. Molina G, Funk LM, Rodriguez V, Lipsitz SR, Gawande A. Evaluation of surgical care in El Salvador using the WHO surgical vital statistics. *World J Surg.* 2013;37:1227-35.
32. Agha R. Towards national surgical surveillance in the UK – a pilot study. *PLoS One.* 2012;7:e47969.
33. Knowlton LM, Chackungal S, Dahn B, LeBrun D, Nickerson J, McQueen K. Liberian surgical and anesthesia infrastructure: a survey of county hospitals. *World J Surg.* 2013;37:721-9.
34. OMS [Internet]. 2017 [Cited 2017 Mar 22] Available from: <http://www.who.int/countries/es/>
35. Weiser TG, Haynes AB, Molina G, Lipsitz SR, Esquivel MM, Uribe-Leitz T, et al. Size and distribution of the global volume of surgery in 2012. *Bull World Health Organ.* 2016;94, 201-209F.

36. Ministerio de Salud. Oficina General de Estadística e Informática. RENIPRESS. 2017 [Cited 2017 Mar 03] Available from: [http://www.minsa.gob.pe/portalweb/02estadistica/estadistica\\_2.asp?sub5=2](http://www.minsa.gob.pe/portalweb/02estadistica/estadistica_2.asp?sub5=2)
37. INEI. Conoce la cédula censal. 2017 [Cited 2017 Mar 03] Available from: [http://www.censos2017.pe/?page\\_id=1927](http://www.censos2017.pe/?page_id=1927)
38. Mayo, Adriana: “Censo del 2017 ubicará cada vivienda con tecnología precisa”. Diario La República. 23 May 2016. [Cited 2017 Mar 07] Available from: <http://larepublica.pe/impresa/sociedad/769713-censo-del-2017-ubicara-cada-vivienda-con-tecnologia-precisa>