



IMAGE OF THE MONTH

Gastric heterotopia of the rectum

Heterotopía gástrica en el recto


Eduardo Dantas^{a,*}, Diva Yamaguti^b, Kendi Yamazaki^c^a Gastroenterology Department, Centro Hospitalar de Setúbal, Setúbal, Portugal^b Pathology Department, Fleury Medicina Diagnóstica, São Paulo, Brazil^c Gastrointestinal Endoscopy Unit, University of São Paulo Medical School, São Paulo, Brazil

Figure 1 Colonoscopy showing a 12mm non-granular flat lesion in the rectum, at 3 cm from the anal verge.

A 44-year-old male was referred for colonoscopy due to family history of colorectal cancer. Colonoscopy revealed a laterally spreading, non-granular flat type lesion (Paris classification 0-IIa), measuring 12 mm, localized in the rectum

at 3 cm from the anal verge (Fig. 1), which was removed en bloc by endoscopic mucosal resection (EMR). Histologic evaluation showed fragments of fundic gastric mucosa along with unaltered colonic mucosa (Fig. 2).

Gastric heterotopia (GHT) has been reported in various locations along the gastrointestinal tract. GHT in the colon is very rare, and most cases are located in the rectum.¹ Clinical presentation is variable, ranging from asymptomatic to the most common symptom of painless rectal bleeding. Endoscopically, rectal GHT may present as diverticula, ulcer, polyp or flat/depressed lesion mimicking early cancer. Histopathological examination is the gold standard for diagnosis. Fundic-type mucosa is the most common histologic subtype described.² *Helicobacter pylori* may colonize gastric mucosa regardless of its location. GHT has an associated risk of malignant transformation, as the first case of colonic adenocarcinoma arising from GHT has already been described.³ Endoscopic removal either by EMR or endoscopic submucosal dissection (ESD) or, alternatively, surgical excision, are considered the treatment of choice.

* Corresponding author.

E-mail address: dantas16@gmail.com (E. Dantas).




Figure 2 Biopsies showing fragments of fundic gastric mucosa along with unaltered colonic mucosa.

References

1. Yang Z, Yates T, Jhala NC. Gastric heterotopia in the rectum. *Clin Gastroenterol Hepatol*. 2010;8:A24.
2. Fang Y, Chen L, Chen DF, Ren WY, Shen CF, Xu Y, et al. Prevalence, histologic and clinical characteristics of heterotopic gastric mucosa in Chinese patients. *World J Gastroenterol*. 2014;20:17588–21759.
3. Ko H, Park SY, Cha EJ, Sohn JS. Colonic adenocarcinoma arising from gastric heterotopia: a case study. *Korean J Pathol*. 2013;47:289–92.