

IMAGE OF THE MONTH

Acute abdomen secondary to torsion and infarction of a wandering spleen



Síndrome abdominal agudo secundario a torsión e infarto de bazo errante

Miguel Angel Perez-Rosillo*, Maria Gomez-Huertas, Angela Salmeron-Ruiz, Antonio Jesus Lainez-Ramos-Bossini

Virgen de las Nieves University Hospital, Granada, Spain

An 18-year-old male presented to the emergency department with a chief complaint of lower abdominal pain of severe intensity for 8 h. His medical history revealed a left diaphragmatic hernia repair two years earlier. Physical examination was unremarkable except for inconclusive

Blumberg's sign. Blood tests showed high levels of C-reactive protein [266 mg/dL] and leukocytosis [15,130 cells/mm³].

Abdominal ultrasound and subsequent CT scan (Fig. 1) revealed that the spleen was not located in the left upper quadrant. A low-enhancing, comma-shaped mass was




Figure 1 Contrast-enhanced abdominal CT in portal (A–C) and delayed (D) phases. Sagittal (A) and coronal (B) reformatting. Hypogastric mass corresponding to enlarged spleen. Inflammatory changes and peri-splenic fluid (green arrows) can be seen. In the coronal view, torsion of the splenic vessels with abnormal surrounding fat is shown (blue arrow). No significant differences in the enhancement pattern between the portal and delayed phases are observed (asterisk in C and D).

* Corresponding author.

E-mail address: maprosillo@gmail.com (M.A. Perez-Rosillo).




Figure 2 Emergent splenectomy through midline laparotomy (A). The spleen showed an ischaemic appearance and torsion of the pedicle was found. (B–D) Surgical specimen consisting of a congestive and enlarged spleen (B, C) with torsion of its pedicle (D).

observed in the hypogastric region showing twisting of its pedicle (“swirl” sign). These findings were consistent with torsion of a wandering spleen with established infarction, thus emergent splenectomy was performed (Fig. 2). The patient evolved favourably and has remained asymptomatic to date.

Wandering spleen consists of the ectopic location of the spleen.¹ Although patients are usually asymptomatic, its torsion may lead to an acute abdomen. Contrast-enhanced CT scan is a cornerstone diagnostic exam in the emergency setting and typical findings include absence or abnormal rotation of the spleen in the left upper quadrant and a “comma” shaped mass in the abdomen.² Treatment requires emergent splenopexy or splenectomy based on the presence of infarction, either through laparotomy or laparoscopy.³

References

1. Rodríguez Vargas D, Parada Blázquez MJ, Vargas Serrano B. Diagnostic imaging of abnormalities in the number and location of the spleen. *Radiologia*. 2019;61:26–34, <http://dx.doi.org/10.1016/j.rx.2018.07.002>.
2. Reisner DC, Burgan CM. Wandering spleen: an overview. *Curr Probl Diagn Radiol*. 2018;47:68–70, <http://dx.doi.org/10.1067/j.cpradiol.2017.02.007>.
3. Awan M, Gallego JL, Al Hamadi A, Vinod VC. Torsion of wandering spleen treated by laparoscopic splenopexy: a case report. *Int J Surg Case Rep*. 2019;62:58–61, <http://dx.doi.org/10.1016/j.ijscr.2019.06.040>.