



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Letter to the Editor

Pediatric tubular acute lymphangitis caused by *Rickettsia sibirica mongolitimonae*: Case report and literature review



Linfangitis tubular aguda pediátrica por *Rickettsia sibirica mongolitimonae*: reporte de caso y revisión bibliográfica

Dear Editor,

In a recent report, Salazar Alarcón et al.¹ described a case of lymphangitis-associated rickettsiosis (LAR) caused by *Rickettsia sibirica mongolitimonae* (*R. sibirica mongolotimonae*), with no cases in the pediatric population previously reported in the literature. In this letter, we report another pediatric case of LAR by *R. sibirica mongolitimonae* in southern Spain, which confirms an expanded distribution of this agent in our country. Moreover, we aim to highlight the potential risks associated with its main vector, the species of the genus *Hyalomma*.² Human transmission by *Hyalomma* ticks of other life-threatening zoonotic agents such as Crimean-Congo hemorrhagic fever virus (CCHFV) has already been reported.³

We report the case of a 4-year-old boy with fever of up to 39.7 °C for 5 days. In the 24 h before the onset of fever, he presented a lesion compatible with a bite on the inner side of the right thigh. Physical examination revealed a circular lesion (5 mm) with a necrotic eschar, surrounded by a slightly raised inflammatory halo (Fig. 1). He associated rope-like lymphangitis running from the eschar up to the inguinal region, where a painful adenopathy (2 cm × 1 cm) was found. The rest of the physical examination was normal. A shave biopsy was performed to determine a Polymerase chain reaction (PCR) for *Rickettsia*. Analytical control was performed with blood count: 10,180 leukocytes (70.8% N, 16.4% L, 12.6% M); Red and platelet series, biochemical profile, and basic coagulation were normal. CRP 24.9 mg/L. He began treatment with oral azithromycin and intravenous amoxicillin-clavulanate. Serological studies of *Rickettsia conorii*, *Borrelia Burgdoferi*, and *Bartonella henselae* were negative. He showed a favorable evolution and was discharged on the fifth day. Follow-up confirmed good evolution and no seroconversion. The diagnosis was confirmed with the results of PCR in the eschar sample performed at the National Center for Microbiology: PCR *Borrelia sp.* negative and positive for *Rickettsia sibirica mongolotimonae*. Bands compatible with *Rickettsia* (SFG) infection were detected, using as targets fragments of the ompA and ompB genes (conventional PCR), and 23S rRNA (real-time PCR). Definitive identification was obtained by molecular sequencing of ompA and ompB genes (491 and 464 bp, respectively) that revealed 100% identity with *Rickettsia sibirica subsp. mongolitimonae*.

Different clinical manifestations of the infection have been described, such as Dermacentor-borne necrosis erythema and lymphadenopathy (DEBONEL) and mainly, lymphangitis-associated rickettsiosis (LAR).² The few reported cases of *R. sibirica mongolo-*



Fig. 1. Necrotic eschar on the inner side of the right thigh, surrounded by a slightly raised inflammatory halo.

timonae infection in children^{1,4–6} are presented in Table 1. The median age of LAR cases was 5 y.o. The three of them were detected in late spring and early summer. In all cases fever was present and just one case presented a generalized skin rash. In the DEBONEL pictures, the presence of eschar and adenopathy also stands out, generally in the upper half of the body and scalp; fever may not always be present and usually occur in cold months.² In adults, severe manifestations have been described,⁷ such as septic shock, acute kidney injury, myopericarditis, retinal vasculitis, and neurological alterations, including encephalitis, among others.

One of the limitations in the study of these cases is the difficulty in identifying the species of tick involved, unknown as in our case. Nevertheless, although it has been isolated in *Rhipicephalus pusillus* and *R. bursa*, its main vector is considered to be the species of the genus *Hyalomma*.² These species are characterized by their aggressive host-seeking behavior⁸ (unlike other ticks utilizing a passive ambush strategy as they wait in vegetation), at a surprising speed for its size.⁸ Perfectly adapted to semi-desert climates,

Table 1
Infections by *Rickettsia sibirica mongolitimonae*^a in children.

Case Year	Geographic location	Month	Risk factors	Age (years)	Sex	Tick bite site	Fever	Rash	Diagnostic	Serology	Reference
5 2020	Alcorcón (Madrid – Spain)	July	Rural dwelling. Tick infested dog contact	11	Men	Upper extremity	Yes	No	LAR ^b	Negative for <i>Rickettsia conorii</i>	Echevarría-Zubero et al. ⁴
4 2020	Illescas (Madrid – Spain)	June	Rural dwelling. Tick infested dog contact	5	Men	Genital area	Yes	Yes	LAR	Negative for <i>Rickettsia conorii</i> and <i>Borrelia burgdorferi</i>	Salazar Alarcón et al. ¹
3 2020	Alcalá la Real (Jaén-Spain)	May	Rural dwelling. Tick infested dog contact	4	Men	Lower extremity	Yes	No	LAR	Negative for <i>Rickettsia conorii</i> , <i>Bartonella Henselae</i> ; and <i>Borrelia Burgdorferi</i>	Case presented in this article
2 2015	Reus (Tarragona-Spain)	Outside the summer season (N.S.) ^d	Rural dwelling	6	N.S.	Head	Yes	Yes	DEBONEL ^c	Negative for <i>Rickettsia conorii</i>	Monterde-Álvarez et al. ⁶
1 2009	Arganda del Rey (Madrid-Spain)	October	Rural dwelling Farm animals	9	Men	Head	Yes	No	DEBONEL ^c	Negative for <i>Rickettsia conorii</i> and <i>Borrelia Burgdorferi</i>	Morales et al. ⁵

^a In all cases, the diagnosis was confirmed with the results of polymerase chain reaction (PCR) in the biopsy of the eschar sample. DNA samples were tested by PCR targeting fragments of the rickettsial genes ompA and ompB (conventional PCR). Definitive identification was obtained by molecular sequence of ompA and ompB, what showed a 100% identity with *Rickettsia sibirica subsp. Mongolitimonae*.

^b Lymphangitis-associated rickettsiosis.

^c Dermacentor-borne necrosis erythema and lymphadenopathy.

^d Not specified.

they usually have maximum activity in the hot and dry months.⁸ They are not very anthropophilic, although, in Spain, a progressive increase in their population is reported^{3,9} along with a high proportion of CCHFV infected ticks collected from wildlife.^{3,9} The change in climatic conditions seems to play an important role in this increment.^{9,10} The fragmentation of the plant habitat and the abandonment of farmland would also be also decisive in tick populations and their hosts, causing increased contact rates between humans and infected ticks.¹⁰ These potential risks emphasize the importance of the development of new prevention strategies and the evaluation of this threat to public health.³

Informed consent

The family of the patient described in this case report gave their informed consent for the inclusion in this publication.

Conflict of interest

The authors declare no conflict of interest in this article.

References

- Salazar Alarcón E, Guillén-Martín S, Callejas-Caballero I, Valero-Arenas A. A propósito de un caso: no toda rickettsiosis es fiebre botonosa mediterránea. *Enferm Infecc Microbiol Clin*. 2021; <http://dx.doi.org/10.1016/j.eimc.2021.01.002>.
- Faccini-Martínez AA, García-Álvarez L, Hidalgo M, Oteo JA. Syndromic classification of rickettsioses: an approach for clinical practice. *Int J Infect Dis*. 2014;28:126–39. <http://dx.doi.org/10.1016/j.ijid.2014.05.025>.
- Portillo A, Palomar AM, Santibáñez P, Oteo JA. Epidemiological aspects of Crimean-Congo hemorrhagic fever in Western Europe: what about the future? *Microorganisms*. 2021;9:649. <http://dx.doi.org/10.3390/microorganisms9030649>.

- Echevarría-Zubero R, Porras-López E, Campelo-Gutiérrez C, Rivas-Crespo JC, Lucas AM, Cobo-Vázquez E. Lymphangitis-associated rickettsiosis by *Rickettsia sibirica mongolitimonae*. *J Pediatric Infect Dis Soc*. 2021;10:797–9. <http://dx.doi.org/10.1093/jpids/piab018>.
- Morales V, García Acebes CR, Miguelez Hernandez AP, Alfagueme Roldán F, Rodríguez Albarrán A, Alins Sahun Y, et al. Infección por *Rickettsia sibirica subsp. mongolitimonae*. *Piel*. 2011;26:224–6. <http://dx.doi.org/10.1016/j.piel.2011.01.005>.
- Monterde-Álvarez ML, Calbet-Ferré C, Rius-Gordillo N, Pujol-Bajador I, Ballester-Bastardie F, Escribano-Subías J. Rickettsiosis tras la picadura de una garrapata: una clínica sutil en muchas ocasiones, debemos estar atentos. *Enferm Infecc Microbiol Clin*. 2017;35:100–3. <http://dx.doi.org/10.1016/j.eimc.2016.01.013>.
- Loarte MDC, Melenotte C, Cassir N, et al. *Rickettsia mongolitimonae* Encephalitis Southern France, 2018. *Emerg Infect Dis*. 2020;26:362–4. <http://dx.doi.org/10.3201/eid2602.181667>.
- Anderson JF, Magnarelli LA. Biology of ticks. *Infect Dis Clin North Am*. 2008;22. <http://dx.doi.org/10.1016/j.idc.2007.12.006>, 195–v.
- Moraga-Fernández A, Ruiz-Fons F, Habela MA, et al. Detection of new Crimean-Congo haemorrhagic fever virus genotypes in ticks feeding on deer and wild boar, Spain. *Transbound Emerg Dis*. 2021;68:993–1000. <http://dx.doi.org/10.1111/tbed.13756>.
- Estrada-Peña A, Ayllón N, de la Fuente J. Impact of climate trends on tick-borne pathogen transmission. *Front Physiol*. 2012;3:64. <http://dx.doi.org/10.3389/fphys.2012.00064>.

Álvaro Vázquez-Pérez^{a,*}, Javier Rodríguez-Granger^b, Elizabeth Calatrava-Hernández^b, Juan Luis Santos-Pérez^a

^a Department of Pediatrics, Virgen de las Nieves Children's Hospital, Granada, Spain

^b Department of Microbiology, Virgen de las Nieves University Hospital, Granada, Spain

* Corresponding author.

E-mail address: avzqpz@gmail.com (Á. Vázquez-Pérez).