




# NEUROLOGY PERSPECTIVES

[www.journals.elsevier.com/neurology-perspectives](http://www.journals.elsevier.com/neurology-perspectives)



## SCIENTIFIC LETTER

### HaNDL syndrome after COVID-19

### Síndrome de HaNDL tras infección por COVID-19

J.C.N. García<sup>a</sup>, M.T.A. Isasi<sup>a,\*</sup>, C. M.Parada<sup>b</sup>, J.V. Lorenzo<sup>a</sup>

<sup>a</sup> *Sección de Neurología, Servicio de Medicina Interna, Hospital Virgen del Castillo, Yecla, Murcia, Spain*

<sup>b</sup> *Residente de Medicina Familiar y Comunitaria, Unidad Docente Multiprofesional de Atención Familiar y Comunitaria de la Vega Media del Segura, Vega Alta del Segura y Altiplano de Murcia, Spain*

Received 26 April 2022; accepted 9 June 2022

Available online 18 July 2022



Dear Editor:

Headache and neurological deficits with cerebrospinal fluid (CSF) lymphocytosis, also known as HaNDL syndrome, is characterised by episodes of migraine-like headache associated with motor, sensory, or language deficits lasting several hours and recurring within less than 3 months.<sup>1</sup> Diagnosis is confirmed by predominantly lymphocytic pleocytosis and normal results in other complementary tests.<sup>2</sup>

The aetiopathogenic mechanism of HaNDL syndrome is unclear, although the condition is thought to be a transient inflammatory process caused by an immune response to viral infection (e.g., HHV-7, EBV, CMV, HHV-6)<sup>3–6</sup> or other pathogens; this would trigger the production of antibodies and/or inflammatory cytokines, which would act on leptomeningeal vessels,<sup>1</sup> causing vasoconstriction and vasodilation.<sup>7</sup>

We present the case of a 14-year-old boy with no relevant personal or family history who visited the emergency department due to an episode of sudden-onset, moderate-intensity holocranial pulsatile headache, associated with nausea, photo- and phonophobia, and language impairment in the form of paraphasia and blocking. Symptoms persisted for over 4 h. He had not presented fever or any other systemic symptoms in the previous days, but in the previous month he had presented a mild upper respiratory tract infection, with odynophagia and mild fever, and a PCR test had yielded positive results for SARS-CoV-2.

The patient was asymptomatic and haemodynamically stable, and presented no fever. Blood analysis did not reveal elevated levels of acute-phase reactants or blood count alterations, and a urine toxicology screening test yielded negative results. Eye fundus examination did not reveal signs of intracranial hypertension, and a head CT scan detected no relevant alterations. CSF opening pressure was 20 cm H<sub>2</sub>O, and a biochemical analysis revealed pleocytosis (525 cells/ $\mu$ L, mostly lymphocytes [99%]), elevated protein levels (98.2 mg/dL), and lack of glucose consumption. A CSF culture did not detect bacterial growth. We started empirical treatment with intravenous aciclovir and lacosamide, and the patient was admitted to the neurology ward.

During hospitalisation, the patient presented 2 additional episodes of headache with the same characteristics, accompanied by language alterations and sensory alterations affecting the left limbs, after which he recovered completely. Complementary analyses showed normal thyroid, liver, and kidney function; a normal lipid profile; and normal autoimmune test results. Serology tests for HCV, EBV, CMV, HSV, VZV, HHV-6, HHV-7, HHV-8, syphilis, *Rickettsia*, *Toxoplasma gondii*, *Borrelia*, and *Brucella* yielded negative results, as did CSF PCR assays for viral infection. Magnetic resonance angiography revealed no parenchymal alterations or intravenous contrast uptake. A video EEG study did not detect focal slowing or epileptogenic foci. In view of the negative viral PCR test results and the lack of additional episodes, antiviral and antiepileptic treatment was discontinued. A second CSF analysis performed before discharge

\* Corresponding author.

E-mail address: [mtalbaisasi@gmail.com](mailto:mtalbaisasi@gmail.com) (M.T.A. Isasi).

**Table 1** International Classification of Headache Disorders (ICHD-3) diagnostic criteria for HaNDL syndrome.

Syndrome of transient headache and neurological deficits with cerebrospinal fluid lymphocytosis (HaNDL syndrome).

A) Episodes of migraine-like headache fulfilling criteria B and C.

B) Both of the following:

1. Accompanied or shortly preceded by onset of at least one of the following transient neurological deficits lasting >4 h:

a) hemiparaesthesia

b) dysphasia

c) hemiparesis

2. Associated with CSF lymphocytic pleocytosis (>15 white cells/ $\mu$ L), with negative aetiological studies

C) Evidence of causation demonstrated by either or both of the following:

1. Headache and transient neurological deficits have developed or significantly worsened in temporal relation to onset or worsening of the CSF lymphocytic pleocytosis, or led to its discovery.

2. Headache and transient neurological deficits have significantly improved in parallel with improvement in the CSF lymphocytic pleocytosis.

D) Not better accounted for by another ICHD-3 diagnosis.

showed improvements, with a leukocyte count of 150 cells/ $\mu$ L and normal protein levels.

HaNDL syndrome is included in the International Classification of Headache Disorders<sup>8</sup>; the diagnostic criteria for this rare entity are listed in [Table 1](#).

In younger patients, the condition may be misdiagnosed as hemiplegic migraine (associated with a mutation in *CACNA1A*)<sup>2</sup> or familial hemiplegic migraine type 1.<sup>1</sup> In fact, the mechanism believed to cause the neurological symptoms is cortical spreading depression, similarly to what occurs in migraine aura.<sup>9</sup> In older individuals, the condition may present as a stroke mimic, which, when in doubt, is managed with fibrinolytic treatment.<sup>1</sup> Differential diagnosis of HaNDL syndrome includes a wide range of diseases; in addition to the conditions mentioned previously, it also includes CNS vasculitis, infectious or inflammatory encephalitis, neoplastic arachnoiditis, Mollaret meningitis, and posterior reversible encephalopathy.<sup>9</sup>

Although diagnosis is established by exclusion, other complementary findings have been identified in recent years, including leptomeningeal contrast uptake on brain MRI, focal or diffuse slowing of EEG activity,<sup>2,7,9</sup> cerebral hypoperfusion predominantly in the temporal lobe,<sup>10,11</sup> and vasospasm in transcranial Doppler ultrasound studies.<sup>7</sup> Given its excellent prognosis without specific treatment and the monophasic course of the disease,<sup>1,2</sup> recurrence should lead us to reconsider the diagnosis of HaNDL syndrome.

As occurs with other neurotropic viruses, SARS-CoV-2 may cause HaNDL syndrome due to its capacity to trigger a "cytokine storm,"<sup>12</sup> resulting in the vasomotor alterations that are believed to cause the syndrome. History of SARS-CoV-2 infection should therefore be taken into account when HaNDL syndrome is suspected.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## Ethical considerations

The authors observed their centre's protocols for the publication of patient data. The patient gave informed consent for the publication of this case report.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neuro.2022.06.007>.

## References

- Filina T, Feja KN, Tolan Jr RW. An adolescent with pseudomigraine, transient headache, neurological deficits, and lymphocytic pleocytosis (HaNDL Syndrome): case report and review of the literature. *Clin Pediatr (Phila)*. 2013 Jun;52(6):496–502. <https://doi.org/10.1177/0009922813483358> Epub 2013 Apr 3. PMID: 23559488.
- Babi MA, Applebee A, Shapiro R, Waheed W. Syndrome of transient headache and neurologic deficits with cerebrospinal fluid lymphocytosis presenting as acute neurological emergencies. *Cephalalgia*. 2017 Mar;37(3):284–9. <https://doi.org/10.1177/0333102416642733> Epub 2016 Sep 29. PMID: 27036641.
- Stelten BM, Venhovens J, van der Velden LB, Meulstee J, Verhagen WJ. Syndrome of transient headache and neurological deficits with cerebrospinal fluid lymphocytosis (HaNDL): A case report with serial electroencephalography (EEG) recordings. Is there an association with human herpes virus type 7 (HHV-7) infection? *Cephalalgia*. 2016 Nov;36(13):1296–301. <https://doi.org/10.1177/0333102415618616> Epub 2016 Jul 20. PMID: 26682576.
- Apetse K, Breynaert L, Butaud C, Beschet A, Blanc-Lasserre K, Ribouillard L, Chan V. Transient headache and neurological deficits with cerebrospinal fluid lymphocytosis associated with IgM antibodies to the Epstein-Barr virus viral capsid antigen. *Case Rep Neurol Med*. 2013;2013:975709. <https://doi.org/10.1155/2013/975709> Epub 2013 Apr 17. PMID: 23691380; PMCID: PMC3652098.
- Verentzioti A, Tavernarakis A, Mamali M, Siatouni A, Gatzonis S. Pseudomigraine with transient neurological deficits and cerebrospinal fluid lymphocytosis or HaNDL syndrome: a case report with confusion and positive IgM antibodies to CMV in serum. *Cephalalgia*. 2017 Jan;37(1):99–100. <https://doi.org/10.1177/0333102416629808> Epub 2016 Sep 29. PMID: 26848127.
- Emond H, Schnorf H, Poloni C, Vuillimoz S, Lalive PH. Syndrome of transient headache and neurological deficits with CSF lymphocytosis (HaNDL) associated with recent human herpesvirus-6 infection. *Cephalalgia*. 2009 Apr;29(4):487–91. <https://doi.org/10.1111/j.1468-2982.2008.01763.x> Epub 2009 Jan 8. PMID: 19170703.
- Hidalgo de la Cruz M, Domínguez Rubio R, Luque Buzo E, Díaz Otero F, Vázquez Alén P, Orcajo Rincón J, Prieto Montalvo J,

- Contreras Chicote A, Grandas Pérez F. Syndrome of transient headache and neurological deficits with cerebrospinal fluid lymphocytosis (HaNDL) in a patient with confusional symptoms, diffuse EEG abnormalities, and bilateral vasospasm in transcranial Doppler ultrasound: a case report and literature review. *Neurología (English Edition)*. October 2019;34(8):536–42.
8. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. 2013;33:629–808. <https://doi.org/10.1177/0333102413485658>.
  9. Armstrong-Javors A, Krishnamoorthy K. HaNDL syndrome: case report and literature review. *J Child Neurol*. 2019 Mar;34(3):161–7. <https://doi.org/10.1177/0883073818811546> Epub 2018 Dec 5. PMID: 30514135.
  10. García-Espérón C, Lozano M, Escudero D. Focal leptomeningeal uptake, a new radiological finding in pseudomigraine with pleocytosis. *Neurología (English Edition)*. 2017;32(1):63–5 Febrero.
  11. Ho BL, Lai CL, Hsu CY. Acute confusion in headache with neurologic deficits and cerebrospinal fluid lymphocytosis syndrome. *Am J Emerg Med*. 2012 Nov;30(9):2078.e7–8. <https://doi.org/10.1016/j.ajem.2011.10.011> Epub 2011 Dec 6. PMID: 22154160.
  12. Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19. *J Infect*. 2020;80:607–13. <https://doi.org/10.1016/j.jinf.2020.03.037> 0163-4453.