

CLINICAL RESEARCH

In-hospital outcome in patients with ST elevation myocardial infarction and right bundle branch block. A sub-study from RENASICA II, a national multicenter registry

Úrsulo Juárez-Herrera,¹ Carlos Jerjes Sánchez,² Héctor González-Pacheco,¹ Carlos Martínez-Sánchez.³

¹ Staff Coronary Care Unit of Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico

² Head of the Emergency Room Department, Hospital de Cardiología N° 34, IMSS. Monterrey, NL. Mexico

³ Head of the Coronary Care Unit. Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico.

Received in June 1, 2009; accepted in February 15, 2010.

KEY WORDS

Acute coronary syndrome;
ST elevation myocardial
infarction;
Bundle branch block;
Right bundle branch block;
Mortality; Mexico.

Abstract

Objective: Compare in-hospital outcome in patients with ST-elevation myocardial infarction with right versus left bundle branch block.

Methods: RENASICA II, a national mexican registry enrolled 8098 patients with final diagnosis of acute coronary syndrome secondary to ischemic heart disease. In 4555 STEMI patients, 545 had bundle branch block, 318 (58.3%) with right and 225 patients with left (41.6%). Both groups were compared in terms of in-hospital outcome through major cardiovascular adverse events; (cardiovascular death, recurrent ischemia and reinfarction). Multivariable analysis was performed to identify in-hospital mortality risk among right and left bundle branch block patients.

Results: There were not statistical differences in both groups regarding baseline characteristics, time of ischemia, myocardial infarction location, ventricular dysfunction and reperfusion strategies. In-hospital outcome in bundle branch block group was characterized by a high incidence of major cardiovascular adverse events with a trend to higher mortality in patients with right bundle branch block (OR 1.70, CI 1.19 - 2.42, $p < 0.003$), compared to left bundle branch block patients. **Conclusion:** In this sub-study right bundle branch block accompanying ST-elevation myocardial infarction of any location at emergency room presentation was an independent predictor of high in-hospital mortality.

PALABRAS CLAVE

Síndrome coronario
agudo; Infarto del
miocardio con elevación
del ST; Bloqueo de rama;
Bloqueo de rama derecha;
Mortalidad; México.

Evolución intrahospitalaria en pacientes con infarto agudo del miocardio con elevación del segmento ST y bloqueo de rama derecha. Sub-estudio de RENASICA II, un registro nacional multicéntrico

Resumen

Objetivo: Comparar la evolución hospitalaria en pacientes con infarto agudo del miocardio con bloqueo de rama derecha versus bloqueo de rama izquierda. **Método:** El Registro Nacional de Síndromes Coronarios Agudos II, incluyó 8098 pacientes con síndrome coronario agudo, de los

Corresponding author: Carlos Martínez-Sánchez. Coronary Care Unit of Instituto Nacional de Cardiología Ignacio Chávez. Juan Ba-
diano No. 1. Tlalpan. Mexico City. Col Secc. XVI ZC 14080 Tel-Fax (+52+55) 54 85-99-99. E-mail: carlos.martinez@cardiologia.org.
mx

PALABRAS CLAVE

Síndrome coronario agudo; Infarto del miocardio con elevación del ST; Bloqueo de rama; Bloqueo de rama derecha; Mortalidad; México.

cuales 4555 corresponden a infarto con elevación del segmento ST. De ellos, se demostró en 545 bloqueo de rama: 318 (58.3%) tuvieron bloqueo de rama derecha y 227 (41.6%) bloqueo de rama izquierda. Fueron comparados en términos de mortalidad hospitalaria y eventos cardiovasculares mayores adversos. Se realizó un análisis multivariado para identificar mortalidad hospitalaria a través de eventos mayores entre pacientes con ambos bloqueos de rama.

Resultados: No hubo deferencia estadísticamente significativa en ambos grupos en relación con características basales, tiempo de isquemia, localización del infarto, disfunción ventricular o estrategia de reperfusión utilizada. Los pacientes con infarto agudo del miocardio de cualquier localización y bloqueo de rama derecha tuvieron mayor tendencia para mortalidad hospitalaria y eventos cardiovasculares mayores (OR 1.70, IC 1.19-2.42, $p < 0.003$) vs. pacientes con bloqueo de rama izquierda.

Conclusión: En el infarto agudo del miocardio con elevación del segmento ST, el bloqueo de rama derecha fue un predictor independiente de alta mortalidad hospitalaria.

Introduction

Data coming from reperfusion era showed,¹ that bundle branch block (BBB) in early acute phases of ST-elevation myocardial infarction (STEMI) had close relationship with high mortality.² Although, the evidences³ suggesting that the real incidence of a BBB is variable, higher incidence of right bundle branch block (RBBB) over left bundle branch block (LBBB) has been identified. In patients with RBBB extensive jeopardized myocardium involving interventricular septum, could induce a worse in-hospital outcome. In addition, on ECG bases it is not possible to separate new or presumably new conduction disturbances from a pre-existing condition. The purpose of this *post-hoc* substudy was to identify in-hospital outcome and prognostic value of RBBB in STEMI from the largest Mexican acute coronary syndromes registry.⁴

Methods

The characteristics of RENASICA II registry has been previously published,⁴ in brief: it is a prospective and observational registry of the Mexican Cardiology Society that included 8098 patients with final diagnosis of acute coronary syndrome. The main target was to identify a representative population to know diagnosis, stratification and treatment trends. Patients were enrolled in primary and tertiary hospitals. To improve the quality of data, criteria of Alpert were used. The hospital varied in terms of access to on-site cardiac catheterization, number of acute care beds and the type of practice setting. Patients were selected during the hospital admission if they had high clinical suspicion of acute coronary syndromes with symptoms and signs of acute ischemia, with or without electrocardiographic changes, with necrosis or not and proved ischemic heart disease by invasive or non invasive test at discharge. Patients with symptoms precipitated by anemia, hypertension, and heart failure or another secondary condition were not considered. On admission and discharge nomenclature with or without ST elevation was standardized. All treatment decisions were made at discretion of treating physicians. In RENASICA tertiary hospitals with capabilities for coronary arteriography, percutaneous transluminal coronary angioplasty (PTCA) and coronary artery bypass graft (CABG) surgery enrolled 90% of the patients.

In this substudy, STEMI patients plus LBBB or RBBB were compared in terms of in-hospital outcome and MACE (cardiovascular death, recurrent ischemia and reinfarction).


Clinical and electrocardiographic criteria: 1) Characteristics and definitions of STEMI patients in RENASICA II have been previously published,⁴ 2) LBBB under universal criteria with ≥ 120 msc QRS duration. Exclusion criteria: a) acute ischemia precipitated by anemia, hypertension, heart failure or another secondary condition, b) well identified previous BBB and pacemaker rhythm.

RBBB definition: QRS duration ≥ 120 msc, with an rsr' , rsR' in V_1 or V_2 leads, R wave notched in V_1 with prolonged R wave peak time > 50 msc in V_1 and normal peak time in V_5 and V_6 . Leads DI and V_6 had to show a QRS complex with a wide S wave $> R$ duration or > 40 msc.⁵ ST segment deviation was measured at maximum J point. Abnormal Q waves were interpreted according to the European Society of Cardiology / American College of Cardiology consensus.⁶

Statistical Analysis: To analyze the clinical characteristics through non-parametric and parametric variables *chi-squared* test and *Student t* test were used. To determine a normal or abnormal distribution, Wilcoxon rank sum test was considered. To analyze the relationship among mortality and mortality markers Pearson's correlation analysis was performed. Through univariate, logistic and multivariate regression analysis, we examined the relationship between variables for atherosclerosis (smoking habit, diabetes mellitus, systemic hypertension and dislipidemia, etc.) and bad outcome (> 60 years of age, diabetes mellitus, anterior or extensive infarct, LBBB, ventricular dysfunction, and expulsion fraction $< 40\%$) with mortality. To evaluate the relationship between each marker and other variables a multivariate Cox proportional hazard model was used. A $p < 0.05$ was considered statistically significant. Data are presented in percentages, mean standard deviation or median values, odds ratio (OR) and confidence intervals 95% (CI).

Results

RENASICA II registry enrolled a total of 8098 patients, 4555 with final diagnosis of STEMI, of these; 4010 patients without BBB (88%) and 545 patients with BBB (11.9%) were analyzed in this sub-study (Figure 1). In this group, 318 (58.3%) had RBBB and 227 (41.6%) LBBB. Table 1 shows



STEMI= ST elevation acute myocardial infarction, BBB= bundle branch block, RBBB= right bundle branch block, LBBB= left bundle branch block

Figure 1. RENASICA II Substudy patients recruited.

demographic characteristics in both groups, AMI location and Killip & Kimbal status. Table 2 shows reperfusion strategy and adjunctive treatment. Regarding age, major risk factors, AMI location, clinical expression of left ventricular dysfunction, reperfusion strategies and adjunctive treatment no statistical differences were observed. A trend for male gender, incidence of anterior or inferior AMI, pharmacological or mechanical reperfusion was observed in patients with RBBB compared to LBBB group. Low use of reperfusion strategies and optimal treatment was observed in both groups. In terms of mortality no statistically significant difference was observed among RBBB group compared with LBBB patients (20% vs. 18%, $p = ns$). Among 4555 STEMI patients, 1685 (37%) were under fibrinolytic therapy, the time to onset symptoms and drug administration was < 2 hours in 31%, from 2 to 4 hours in 36%, 4 to 6 hours in 19% and finally > 6 hours in 15%. Primary or facilitated PTCA were performed in 15%. A logistic regression analysis performed to assess the effect of independent variables on mortality while adjusting for potentially confounding factors identified ECG findings as in-hospital mortality predictors (Figure 2) including RBBB (OR 1.70, 95% CI 1.19 - 2.4, $p < 0.0003$), LBBB (OR 1.7, 95% CI 1.1 - 2.5, $p = 0.007$), AV block high-degree (OR 2.99, 95% CI 1.9-3.1) and STEMI plus ST depression > 3 ECG leads (OR 3.43, 95% CI 1.51 - 7.78, $p = 0.003$). Several multivariate regression models were applied to identify stronger correlation and a higher statistical significance for mortality. In-hospital outcome and MACE are shown in Figure 3. A high incidence of cardiovascular mortality and MACE were observed in both groups.

Discussion

Our data show four important findings. First, in-hospital outcome of STEMI patients with BBB was characterized by a high mortality. Second, as compared with LBBB, RBBB ECG finding was a mortality predictor too. Third, focus in low use of reperfusion strategies and optimal adjunctive treatment is mandatory. Fourth, this could be the first evidence about the outcome of STEMI patients complicated with BBB coming from a Mexican representative population.⁴

Table 1. Demographic characteristics, acute myocardial infarction location and Killip & Kimbal status.

Variable	RBBB n= 318 (%)	LBBB n= 227 (%)	$p <$
Age (years, median)	66.7	67.3	ns
Men	76	71	ns
Hypertension	55	59	ns
Hyperlipidemia	27	26	ns
Current Smoker	63	66	ns
Diabetes	48	47	ns
Previous AMI	32	35	ns
Anterior	32	23	ns
Inferior	23	16	ns
KK I	74	74	ns
KK II	17	16	ns
KK III	5	5	ns
KK IV	4	4	ns

RBBB= right bundle branch block, LBBB= left bundle branch block, AMI= acute myocardial infarction, ns= no significance

In the setting of STEMI, the clinical relevance of BBB has been established before and after reperfusion era.⁷ In the Fibrinolytic Therapy Trialist's,⁸ the mortality with or without thrombolysis was 19% and 24% respectively in the first 35 days. The studies did not establish any distinction between RBBB or LBBB and did not specify if BBB were new or not. Different types of BBB occurring during the initial hours of AMI with different prognostic implications that is independent of other prognostic factors.

In our substudy, patients with RBBB had trend to major incidence of anterior or inferior infarction compared to LBBB group. In these patients ECG criteria for diagnosis of STEMI with LBBB possibly were not used. An analysis based on simple ST segment changes, may help identify patients with acute myocardial infarction who can then receive an appropriate reperfusion treatment.⁹

In the earliest phase of STEMI, independently of other prognostic factors, several types of BBB occurred with different prognostic implications. In our study it was not possible to identify the new appearance of BBB. This abnormal conduction disturbance in a patient with acute chest pain is highly suggestive of infarction. Ischemic changes superimposed on a pattern of chronic LBBB are easy to recognize when a previous ECG is available for comparison. The timely availability of a previous ECG, however, is the exception rather than the rule. An important issue of this study is that physicians at emergency room department have to decide to administer appropriate reperfusion strategy on the basis of only the most recent ECG.⁹

An interesting finding in this study was the relationship among STEMI, RBBB and high incidence of MACE. Several anatomic conditions could explain this outcome. a)


Table 2. Reperfusion strategy and adjunctive treatment in patients with right and left bundle branch block.

Variable	RBBB n= 318 (%)	LBBB n= 227 (%)	p <
Reperfusion approach			
Lytic	32	23	ns
Primary PTCA	23	20	ns
Adjunctive treatment			
Aspirin	88	89	ns
Clopidogrel	44	38	ns
UFH/LMWH	47/34	48/32	ns
Beta-blockers	51	51	ns
ACEI/ARB	64	59	ns
Statins	14	13	ns


RBBB= right bundle branch block, LBBB= left bundle branch block, AMI= acute myocardial infarction, PTCA= percutaneous coronary angioplasty, UFH/LMWH = unfractionated heparin/ low molecular weight heparin, ACEI / ARB= angiotensin conversion enzyme inhibitors / angiotensin receptor blockers, ns= no significance

Abnormal conduction disturbance traverses the interventricular septum toward the cardiac apex; b) in anterior acute myocardial infarction, the proximal occlusion of the left anterior descending artery produces ischemia of the septum and RBBB; c) sometimes, RBBB and LBBB divide into multiple portions with an atrio-ventricular branch, constituting a true bifascicular conduction abnormality, with a distribution network of anterosuperior and central fibers.¹⁰ This variable anatomy in the conduction system produces variations when ischemia or necrosis occurs and causes conduction disturbances in STEMI, explaining the variable incidences and prognostic meanings of the left fascicular blocks occurring simultaneously to RBBB.¹¹ The contributions from the Mexican school of electrocardiography in the diagnosis of RBBB has been described in three elegant terms: a) without “jumping-wave” phenomenon or minor grade block, b) with “jumping-wave” phenomenon, and c) with limited “jumping-wave” phenomenon or intermediate grade block. The last two types are associated to ischemic conditions and dead septal tissue.^{12, 13}

In the present study patients with RBBB with anterior location AMI had a trend to high in-hospital mortality compared to LBBB patients. This mortality in STEMI and RBBB may be explained by septal ischemia from a more proximal left anterior descending artery occlusion (before the



STEMI = ST elevation acute myocardial infarction, ECG= Electrocardiogram, LBBB = left bundle branch block, RBBB = right bundle branch block, AV = atrio ventricular, OR = odds ratio, CI = confidence interval

Figure 2. ECG findings and logistic regression in hospital mortality predictors in ST elevation acute myocardial infarction.

MACE = major adverse cardiovascular events, LBBB = left bundle branch block, RBBB = right Bundle branch block, ns = no significance

Figure 3. Outcomes and in hospital comparison of MACE in both bundle branch block.

large septal branch). It is important to emphasize that RBBB was a consistent risk marker, as strong as, LBBB for in-hospital mortality. A wider QRS duration (≥ 160 msec) during anterior STEMI and RBBB may reflect more extensive ischemia in the conduction system; currently this ECG finding has been used in the risk stratification.¹⁴ In addition, the observed mortality rates in STEMI were higher than expected and higher than reported previously,⁴ this was not unexpected since a substantial proportion of patients had not access to reperfusion facilities. Left ventricular dysfunction was the most important MACE and the strongest mortality predictor, which is in line with the low incidence of reperfusion approaches and possible long ischemia time. New pharmacological reperfusion directions including FT bolus, as is currently used in several countries, might improve even further STEMI outcome. Current evidence to use an optimal treatment (clopidogrel, enoxaparin and statins) was obtained after RENASICA.¹⁵⁻¹⁷

Clinical Implications: Abnormal conduction disturbances plus ST depression > 3 leads, in the setting of STEMI allowed to identify a high risk group to in-hospital mortality (Figure 2). Prompt recognition and reperfusion strategies should improve survival. In addition, physicians have

to have in mind that the timely availability of a previous ECG is the exception rather than the rule. At emergency room department, physicians in charge have to decide reperfusion strategies on the basis of only the most recent ECG. Considering the significant proportion of patients without any reperfusion strategy, new directions from the Health System and Mexican Cardiology Society are required to improve quality of care.

Limitations: As in all clinical trials, a selection bias could have occurred in RENASICA II resulting in underrepresentation of very high-risk patients, including those with RBBB accompanying anterior STEMI in the trial cohort. In addition, the ECG interpretation was not performed in all centers by experts in electrocardiography. Thus, it is not possible to study the different types of RBBB. As this was a transversal study, it was not possible to identify if BBB was a new condition.

Conclusion

In STEMI RBBB was an independent predictor of high in-hospital mortality and had at least the same risk implication than LBBB. Both should be considered in risk stratification to identify high risk patients.

References

- Norris RM, Croxson MS: Bundle branch block in acute myocardial infarction. *Am Heart J*.1970;79:728-733.
- Melgarejo A, Galcerá J, García A, et al. Incidencia, características clínicas y significación pronóstica del bloqueo de rama izquierda asociado a infarto agudo de miocardio. *Rev Esp Cardiol* 1999;52:245-252.
- Hindman MC, Wagner GS, JaRo M, et al. The clinical significance of bundle branch block complicating acute myocardial infarction. Clinical characteristics, hospital mortality, and one-year follow-up. *Circulation* 1978;58:679-688.
- Castillo GA, Jerjes-Sánchez C, Martínez BP, et al. por los Investigadores del Registro Nacional de Síndromes Coronarios Agudos II. Registro Nacional de Síndromes Coronarios Agudos. RENASICA II. *Arch Cardiol Mex* 2005;75:S6-S19.
- Willems JL, Robles de Medina EO, Bernard R, et al. Criteria for intraventricular conduction disturbances and pre-excitation: World Heart organization / International Society and Federation for Cardiology Task Force. *J Am Coll Cardiol*. 1985;5:1261-1275.
- The Joint European Society of Cardiology / American College of Cardiology Committee Myocardial Infarction Redefined: a consensus document for the Joint European Society of Cardiology/ American College of Cardiology Committee for the redefinition of Myocardial infarction. *J Am Coll Cardiol*. 2000;36:959-969.
- Sgarbossa EB, Pinski SL, Topol EJ, et al, for the GUSTO Investigators. Acute myocardial infarction and complete bundle branch block at hospital admission: clinical characteristics and outcome in the thrombolytic era. *J Am Coll Cardiol* 1998;31:105-110.
- Wong CK, Stewart RA, Gao W, et al. For the Hirulog and Early Reperfusion or Occlusion (HERO-2) Trial Investigators. Prognostic differences between different types of bundle branch block during the early phase of acute myocardial infarction: insights from the Hirulog and Early reperfusion or Occlusion (Hero-2) trial. *Eur Heart J* 2006;27:21-28.
- Sgarbossa EB, Pinski SL, Barbagelata A, et al. for the GUSTO Investigators. Electrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle-branch block. *N Engl J Med* 1996;334:481-487.
- Rubart M, Zipes DP. Genesis of cardiac arrhythmias: electrophysiological considerations. In: Zipes DP, Libby P, Bonow RO, Braunwald E. editors. *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. 7th ed. Philadelphia, Pa: Saunders;2005:653-688.
- Ricou F, Nicod P, Gilpin E, et al. Influence of right bundle branch block on short- and long- term survival after acute anterior myocardial infarction. *J Am Coll Cardiol* 1991;17:858-863.
- De Micheli A, Medrano GA. Enfoque electrofisiológico del diagnóstico de bloqueos ventriculares derechos no complicados y complicados. *Arch Inst Cardiol Mex* 1996; 66:269-281.
- De Micheli, Aranda A, Medrano GA. Aspectos eléctricos de la asociación de miocardio inactivable y bloqueos tronculares de grado intermedio. *Arch Cardiol Mex* 2007;77:67-73.
- Wong CK, Wanzhen G, Ralph AH, et al. On behalf of Hero 2 Investigators. Risk stratification of patients with acute anterior myocardial infarction and right bundle branch block. Importance of QRS duration and early ST segment resolution after fibrinolytic therapy. *Circulation* 2006;114:783-789.
- Sabatine MS, Cannon CP, Gibson CM, et al. for the CLARITY-TIMI 28 Investigators. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med* 2005;352:1179-1189.
- Antman EM, Morrow DA, McCabe CH, et al. for the EXTRACT - TIMI 25 Investigators. Enoxaparin versus unfractionated heparin for ST - elevation myocardial infarction. *N Engl J Med* 2006;354:1477-1488.
- Cannon CP, Braunwald E, McCabe CH, et al. for de Pravastatin or Atorvastatin Evaluation and Infection TIMI 22 Investigators. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350:1495-1504.